

Examples of Spherical Varieties

Spherical varieties seminar - August 3rd 2018

references: • "Lectures on spherical and wonderful varieties" by Guido Pezzini
• "Intro to spherical varieties" by Boris Pasquier
• "Frobenius splitting Methods in Geometry + Rep Theory" by Michel Brion + Shrawan Kumar

GOAL: Discuss various examples of spherical, horospherical, and wonderful varieties, as well as any related definitions and theorems that we haven't seen in this seminar.

Notation/Defn Reminder

Def. • A G -variety is **spherical** if it is normal and has an open B -orbit.

• (X, x) is an **embedding** of G/H if X is spherical, the G -orbit Gx is open in X , and H is the stabilizer of $x \in X$. We call an embedding **simple** if it has a unique closed G -orbit

↳ It's also useful to recall that any spherical G -variety admits a cover by open G -stable simple spherical varieties - so we really only need to worry about simple guys.

Def. • $\mathcal{C}(X)^{(B)} = \{f \in \mathcal{C}(X) \mid bf = \chi(b)f \ \forall b \in B \ \nexists \text{ some } \chi: B \rightarrow \mathbb{C}^* \} = "$ B -Eigenvectors" or **B semi-invariant functions**

• $\Lambda(X) = \{ \chi_f \mid f \in \mathcal{C}(X)^{(B)} \} =$ set of all **B -weights**

• A **color** is a B -stable prime divisor that is not G -stable. We call the set of all colors the **palette**, which we denote by $\Delta(X)$.

• Let (X, x) a simple embedding of G/H . Define $\mathcal{C}(X) \subseteq N(X)$ to be the convex cone generated by $\rho_x(\Delta(X))$ and by all of the G -invariant valuations associated to G -stable prime divisors of X . The pair $(\mathcal{C}(X), \Delta(X))$ is the **painted cone** of X .

• Given an embedding (X, x) of G/H , we define its **painted fan** as:

$$\mathcal{F}(X) = \{ \text{colored cones associated to } X_{y,G} \text{ for any } G\text{-orbit } Y \text{ of } X \}$$

$$\text{with } X_{y,G} = \{ x \in G \mid \overline{G \cdot x} \supseteq Y \}$$

Notation. • $\rho_D =$ valuation associated to the prime divisor D .

$$\bullet N(X) := \text{Hom}_{\mathbb{Z}}(\Lambda(X), \mathbb{Q})$$

• $\mathcal{D}(X, Y) =$ colors of X that contain the closed orbit Y .

• $\mathcal{C}(X) =$ cone (convex) generated by $\rho_x(\mathcal{D}(X))$ \nexists weights associated to G -stable prime divisors of X .

• $\mathcal{U}(X) =$ set of G -invariant valuations on X .

Pezzini's notes (and Rob's!) mention the following:

To classify ALL spherical G -varieties, we can look at the following:

- Fix a spherical subgroup $H \leq G$ and study all embeddings X of G/H
- Study all spherical subgroups $H \leq G$

(Recall: When G/H is a spherical variety, we call H a **spherical subgroup**.)

Revisiting examples of Spherical varieties

AND THEIR PAINTED FANS/CONES

First, I want to fully flesh out an example that we've already seen.

Example When Tracy talked about spherical embeddings, she gave the example of G/H with $G = \mathrm{SL}_2$ and $H = T$. (It's also in Rob's notes from last week) During her talk, we saw that the homogeneous space G/H admits only one nontrivial embedding: $X = \mathbb{P}^1 \times \mathbb{P}^1$. We'll (try) to construct the painted fan for X .

Recall our choice of max torus $T = H$ and Borel subset B :

$$T = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \mid a \neq 0 \right\} \quad \& \quad B = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \mid a \neq 0 \right\}$$

The B -orbit:

$$\begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} a \\ 0 \end{pmatrix} \sim \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} b \\ a^{-1} \end{pmatrix} \quad \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} a+b \\ a^{-1} \end{pmatrix}$$

Performing a change of variables as in Tracy's talk, we see that the B -orbit is iso to

$$\{ (p, q) \in \mathbb{P}^1 \times \mathbb{P}^1 \text{ with } p \neq q \text{ and } p, q \neq \begin{pmatrix} 1 \\ 0 \end{pmatrix} \}$$

(you can also check that this is open)

B -stable divisors: (that aren't G -stable)

From Tracy's talk, we have: $D^+ = \mathbb{P}^1 \times \{[1, 0]\}$ & $D^- = \{[1, 0]\} \times \mathbb{P}^1$

Notice also that the closed G -orbit is $Z = \mathrm{diag}(\mathbb{P}^1)$. We also have the following B -stable affine open set:

$$X_{z,B} = X \setminus (D^+ \cup D^-) = \{[x, 1], [y, 1]\} \cong \mathbb{A}^2$$

From Rob's talk, recall that the function field of $\mathbb{P}^1 \times \mathbb{P}^1 - \Delta(\mathbb{P}^1)$ is the same as that of $\mathbb{A}^2 = \mathbb{C}(x, y)$

In $X_{z,B}$, we know a local equation for Z : $f(x, y) = (x - y)^{-1}$. This is a B -eigenvector of weight $-\alpha_1$.

Thus, we have the following: $\langle \rho(Z), \alpha_1 \rangle = -1$, and $\mathcal{V}(X) = \mathbb{Q}_{\geq 0} v_Z$.

Notice also that $f(x, y)$ has poles of order 1 along D^+ and D^- , so that

$$\langle \rho(D^+), \alpha_1 \rangle = \langle \rho(D^-), \alpha_1 \rangle = 1.$$

$D_1 =$ vanishing of x_2

$D_2 =$ vanishing of x_1

$$\rho(v_{D_1}) = v_b(f) = 1$$


$$v_{D_1}(x_2)$$

$$\text{and}$$

$$\rho(v_{D_2}) = v_{b_2}(f) = 1$$

Finally, from Rob's notes we have that $\Lambda(X) \cong \mathbb{Z}$, so that $N(X) = \mathrm{Hom}_{\mathbb{Z}}(\Lambda(X), \mathbb{Q}) \cong \mathbb{Q}$.

Using this, we get the following painted cone:



Horospherical varieties

Def. A homogeneous space G/H is called **horospherical** if one of the following holds:

- It is a torus bundle over a flag variety G/P
- H contains the unipotent radical of a Borel subgroup
- H is the kernel of characters of a parabolic subgroup P of G

} These are all equivalent

→ This is uniquely defined as the normalizer of H in G

Theorem X is horospherical if and only if $\mathcal{V}(X) = N(X)$.

Example Let $G = SL_2$ again. We'll take $H = U =$ set of unipotent upper Δ matrices in G .

$$H = \left\{ \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \right\} \quad \text{and} \quad B = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \mid a \neq 0 \right\}$$

NOTICE: $G/H = SL_2/U = \mathbb{C}^* \setminus \{(0,0)\}$

• The only color D is given by the equation $\{y=0\}$.

$$\Rightarrow \Lambda(SL_2/U) \cong \mathbb{Z}w_1$$

• $\langle \rho(D), w_1 \rangle = v_b(f_{w_1}) = v_b(y) = 1$

• Notice: $\mathcal{V}(SL_2/U) \cong N(SL_2/U)$.


We have the following nontrivial simple embeddings:



• $(\mathcal{L}_1, \mathcal{D}_1) = (\mathbb{Q}_{\leq 0} \rho(D), \emptyset)$. According to Pezzini, this gives the embedding $G/H \cup \{\text{line @ } \infty\} \dots (\mathbb{P}^1 \setminus \{0\})$




• $(\mathcal{L}_2, \mathcal{D}_2) = (\mathbb{Q}_{\geq 0} \rho(D), \{D\})$. This gives the embedding $G/H \cup \{(0,0)\} \cong \mathbb{C}^2$.




• $(\mathcal{L}_3, \mathcal{D}_3) = (\mathbb{Q}_{\geq 0} \rho(D), \emptyset)$. This gives us $B_0 \mathbb{C}^2$.

This is possibly a stupid question, but how do these painted fans compare to their toric variety counterparts? (ie $B_0 \mathbb{C}^2$... as a toric variety)

ie:



$B_0 \mathbb{C}^2$



\mathbb{C}^2

(The shaded regions are only to indicate the maximal cones of the fan)

... What about the nonsimple embeddings of SL_2/U ? There are two:

↳ painted fan instead of a painted cone!

- $\mathcal{F} = \{(\{0\}, \emptyset), (C_1, D_1), (C_2, D_2)\}$ gives embedding \mathbb{P}^2 . The fan is the following:



- $\mathcal{F} = \{(\{0\}, \emptyset), (C_1, D_1), (C_3, D_3)\}$ which gives $Bl_0 \mathbb{P}^2$



Wonderful Varieties

Def. Let X be a G -variety. We call X wonderful if:

- (i) X is smooth and complete
- (ii) X contains an open G -orbit X_G° whose complement is the union of smooth G -stable prime divisors $X_1^{(u)}, \dots, X_r^{(u)}$ which have normal crossings and nonempty intersection.

(iii) For all $x, y \in X$ we have:

$$Gx = Gy \text{ if and only if } \{i \mid x \in X_i^{(u)}\} = \{j \mid y \in X_j^{(u)}\}$$

The number r (of divisors from part (ii)) is the rank of X , and $\bigcup_{i=1}^r X_i^{(u)}$ (union of G -stable prime divisors) is the boundary of X , which we denote as ∂X .

A note on "normal crossing": I interpret this to mean that the $X_i^{(u)}$ intersect like the coordinate hyperplanes in \mathbb{C}^n . As in Pezzini's notes, any intersection of them will give a wonderful subvariety.

Theorem (due to Luna in 1996) Wonderful varieties are projective and spherical.

Def Let $z \in X$ be the unique point fixed by B . It lies on $Z = Gz =$ unique closed G -orbit. Consider the vector space $T_z X / T_z Z$. (this is naturally a T -module) Its T -weights are called spherical roots of X , and we denote the set of these by Σ_X .

If X is the wonderful embedding of a S.H.S. G/H of rank r , then the spherical roots are in bijection with codim 1 B -stable closed subvarieties, and with codim $r-1$ G -stable closed subvarieties.

Facts (That I found interesting)

- Flag varieties are wonderful of rank 0.
- The only wonderful variety that is toric and wonderful is the point.
- A spherical variety is wonderful if + only if its the canonical embedding of its open G -orbit and the embedding is smooth.

Thus: a spherical homogeneous space admits at most one wonderful embedding

- Wonderful varieties are classified by their associated root system
- A natural question: when does a spherical homog. space admit a wonderful embedding?

~> Classification of such spaces is not yet complete.

Although, there is a necessary condition: $N_B H / H$ must be finite.

The general situation: The homogeneous spaces $G/H = H \times H / \text{diag}(H)$ for H semisimple and adjoint admit wonderful compactification X . X has spherical roots $\sigma_i = \alpha_i + \alpha'_i$, where α_i and α'_i are simple roots for each copy of H . There are also n colors, & we can determine the values of their functionals on each σ_i via the cartan matrix assoc. to H .

According to Pezzini...

Note. SL_2 admits a wonderful compactification. (its the only non-adjoint simple group that does) it is:

$$X = \{ad \cdot bc = t^2\} \subset \mathbb{P}(M_{2 \times 2} \oplus \mathbb{C})$$

Example. Let $G = \mathrm{PGL}_2$. We consider $G \ltimes G$, with $X \cong \mathbb{P}(\mathcal{M}_{2 \times 2}) \cong \mathbb{P}^3$

A wonderful compactification

\mathcal{B} = Borel subgroup of $G \ltimes G$. so we'll take $\mathcal{B} = B \ltimes B$, with:

$$B^- = \left\{ \begin{pmatrix} a & 0 \\ b & c \end{pmatrix} \right\} \quad \text{and} \quad B = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \right\}. \quad \text{Notice that } B \cap B^- = T.$$

• What is the \mathcal{B} -orbit of X ? (How does \mathcal{B} act on the identity?)

$$\begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} c^{-1} & 0 \\ -d & c \end{pmatrix} = \begin{pmatrix} ac^{-1}bd & bc \\ -a^{-1}d & ac \end{pmatrix} \quad \text{Need } a \neq 0.$$

(Also.. didn't Keller mention this in his talk?
... except with $\mathrm{SL}_2 \times \mathrm{SL}_2$?)

so the open \mathcal{B} -orbit will be: $X_B^\circ = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a \neq 0, ad-bc \neq 0 \right\}$

• What is the G -orbit of X ?

This should just be: $\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid ad-bc \neq 0 \right\}$ Now we can figure out what our boundary divisors should be.

so, the open $G \ltimes G$ -orbit of X will be: $X^\circ = \mathbb{P}(\mathcal{M}_{2 \times 2}) \setminus \mathbb{Z}(ad-bc)$

then.. $X \setminus X^\circ$ ought to give us the boundary divisors. Notice $\mathbb{P}(\mathcal{M}_{2 \times 2}) \setminus (\mathbb{P}(\mathcal{M}_{2 \times 2}) \setminus \mathbb{Z}(ad-bc)) =$

$\mathbb{Z}(ad-bc)$
↓
boundary divisor

• What about the colors?

Recall from the defn: $X^\circ \setminus X_B^\circ$ = union of prime B -stable divisors = union of colors

$$\text{so.. } X^\circ \setminus X_B^\circ = (\mathbb{P}^3 \setminus \mathbb{Z}(ad-bc)) \setminus (\mathbb{P}^3 \setminus (\mathbb{Z}(ad-bc) \cup \mathbb{Z}(a))) = \boxed{\mathbb{Z}(a)} \quad \text{one color!}$$

(See "Total coord ring of a Wonderful variety" by Brion)

• Spherical roots (w.r.t $B \ltimes B$) are pairs $(-\alpha_i, \alpha_i)$, where α_i are the simple roots of the root system associated to PGL_2 ... which is A_1 . A choice of simple root for A_1 is $e_1 - e_2$.

So.. $\Lambda(X) \cong$ root lattice of A_1 , and $N(X) = \mathrm{Hom}_{\mathbb{Z}}(\Lambda(X), \mathbb{Q}) \cong$ weight lattice of A_1 . (tensored w/ \mathbb{Q} ?)

Finally... $\mathcal{V}(X)$ = negative Weyl chamber. (I think Blake talked about this?)

an attempt at a painted fan:

