

2.1 and 2.6: Graphs and Graph Theory

Question 1.


(a) Sketch a simple diagram showing the countries Russia, Estonia, Latvia, Lithuania, Belarus, Ukraine, and Poland. Your diagram should show which countries border each other, but otherwise it should be as simple as possible

(b) Is it possible to plan a round trip (on land) through all these countries that visits each country exactly once? Is there more than one way?

(c) How many pairs of countries share a border? Is it possible to plan a trip through all these countries that visits each border crossing exactly once? Find one, or explain why it is impossible.



Example 1. Represent the following diagram as a graph. Can we walk around the city and cross each bridge exactly once?



Terminology: A graph is a collection of vertices, represented by dots, and edges, represented by lines or curves. The edges can be directed or undirected with directed edges being represented by adding an arrow to the edge. The degree of a vertex is the number of times an edge touches it. That means *loops* add two to the degree of a vertex. For directed graphs, we can also speak of *in-degree* and *out-degree*. A path is a sequence

$$v_0, e_1, v_1, e_2, v_2, \dots, v_{n-1}, e_n, v_n$$

such that the edge e_j connects to v_{j-1} and v_j for all $1 \leq j \leq n$. A cycle (or circuit) is a closed path; i.e. $v_0 = v_n$. A graph is said to be connected if there is a path between any two vertices. A *weighted graph or network* is a graph in which the edges have numerical values, usually representing distance, frequency or flow. A coloring of a graph is a way to color all the vertices of the graph such that no edge has the same colors at both ends.

Observations by Euler of the bridges of Königsberg Let $G = (V, E)$ be a graph.

Theorem 1. *Then the sum of the degrees of the vertices is equal to twice the number of edges; i.e.*

$$\sum_{v \in V} \deg(v) = 2|E|.$$

Theorem 2. *If more than two vertices has odd degree, then a path which crosses each edge exactly once (an Euler path) is impossible.*

Theorem 3. *If exactly two vertices have odd degree, then an Euler path may be possible, but you must start at one of the two vertices and end at the other.*

Theorem 4. *If no vertices have odd degree, then a closed Euler path (an Euler circuit) is possible starting at any vertex.*

Example 2. A college registrar's office needs to schedule the following courses: Physics, Computer Science, Chemistry, Calculus, Discrete Math, Biology, and Psychology. The following pairs of classes always have students in common, so they can't be scheduled in the same time slot:


- Physics and Computer Science
- Physics and Chemistry
- Calculus and Chemistry
- Calculus and Physics
- Calculus and Computer Science
- Calculus and Discrete Math
- Calculus and Biology
- Discrete Math and Computer Science
- Discrete Math and Biology
- Psychology and Biology
- Psychology and Chemistry

What is the fewest number of time slots needed to schedule all these classes without conflict?

Binary Search Tree: A graph with no circuits is called a tree. Equivalently, a tree is a connected graph such that there is a unique path between any two vertices. Trees are extremely useful in computer science and writing algorithms. Consider the following *binary search tree* containing the numbers

123, 684, 121, 511, 602, 50, 43.

- (a) What is the *height* of this tree?
- (b) Can we rearrange the numbers so that the height is smaller? What is the minimum height?



Theorem 5. Let T be a tree with n vertices. Then T has $n - 1$ edges.

Proof.

Isomorphism of Graphs: Two graphs $G = (V_G, E_G)$ and $H = (V_H, E_H)$ are isomorphic if they are in one-to-one correspondence. That is to say, if there are maps

$$\alpha : V_G \rightarrow V_H \quad \text{and} \quad \beta : E_G \rightarrow E_H$$

such that, for any edge $e \in E_G$,

$$e \text{ joins vertex } v \text{ to } w \iff \beta(e) \text{ joins vertex } \alpha(v) \text{ to vertex } \alpha(w).$$

In this case, we write $G \cong H$.

Theorem 6. Let $G = (V_G, E_G)$ and $H = (V_H, E_H)$ be graphs without multiple edges. Notice that the collection of edges $E_G \subseteq V_G \times V_G$ is an equivalence relation. If there is a one-to-one correspondence $f : V_G \rightarrow V_H$ with the property that

$$(x, y) \in E_G \iff (f(x), f(y)) \in E_H$$


for all $x, y \in V_G$, then $G \cong H$.

Proof.

G Example 2. Show that the following two graphs are isomorphic.



H



Homework. (Due Oct 31, 2018) Section 2.1: 8, 20; Section 2.6: 11-13

Practice Problems. Section 2.1: 1-4, 9, 10, 17-19; Section 2.6: 3, 5, 9, 10, 14-15, 28-30