

MARK BOX		
PROBLEM	POINTS	
1 a – y	25	
2 a – o	15	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
9	10	
10	10	
11	10	
12	10	
13	10	
14	10	
15	10	
16	10	
17 a – f	10	
18	10	
TOTAL	200	

NAME: \_\_\_\_\_

please check the box of your section

Section 005 (WF 8:00 am)

or

Section 006 (WF 9:05 am)

**Problem Inspiration:**

**1.–3.** Fill in blanks & True/False

**4.–5.** Ch. 7

**6.–11.** Ch. 8

**12.–16.** § 10.1 – 10.6 and 10.8

**17.–18.** § 10.7, 10.9, 10.10

**INSTRUCTIONS:**

- (1) To receive credit you must:
  - (a) **work in a logical fashion, show all your work, indicate your reasoning;**  
no credit will be given for an answer that *just appears*;  
such explanations help with partial credit
  - (b) if a line/box is provided, then:
    - show you work **BELOW** the line/box
    - put your answer on/in the line/box
  - (c) if no such line/box is provided, then box your answer
- (2) The MARK BOX indicates the problems along with their points.  
 Check that your copy of the exam has all of the problems.
- (3) You may **not** use a calculator, books, personal notes.
- (4) During this exam, do not leave your seat. If you have a question, raise your hand. When you finish: turn your exam over, put your pencil down, and raise your hand.
- (5) This exam covers (from *Calculus* by Anton, Bivens, Davis 8<sup>th</sup> ed.):  
 Sections: 7.1 - 7.4, 7.6, 7.7, 8.1 - 8.5, 8.8, 10.1 - 10.10 .

**1. Fill in the blanks.**

**1a.**  $\int \frac{du}{u} = \underline{\hspace{2cm}} |u| + C$

**1b.** If  $a$  is a constant and  $a > 0$  but  $a \neq 1$ , then  $\int a^u du = \underline{\hspace{2cm}} + C$

**1c.**  $\int \cos u du = \underline{\hspace{2cm}} + C$

**1d.**  $\int \sec^2 u du = \underline{\hspace{2cm}} + C$

**1e.**  $\int \sec u \tan u du = \underline{\hspace{2cm}} + C$

**1f.**  $\int \sin u du = \underline{\hspace{2cm}} + C$

**1g.**  $\int \csc^2 u du = \underline{\hspace{2cm}} + C$

**1h.**  $\int \csc u \cot u du = \underline{\hspace{2cm}} + C$

**1i.**  $\int \tan u du = \underline{\hspace{2cm}} + C$

**1j.**  $\int \cot u du = \underline{\hspace{2cm}} + C$

**1k.**  $\int \sec u du = \underline{\hspace{2cm}} + C$

**1l.**  $\int \csc u du = \underline{\hspace{2cm}} + C$

**1m.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{\sqrt{a^2-u^2}} du = \underline{\hspace{2cm}} + C$

**1n.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{a^2+u^2} du = \underline{\hspace{2cm}} + C$

**1o.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{u\sqrt{u^2-a^2}} du = \underline{\hspace{2cm}} + C$

**1p.** Partial Fraction Decomposition. If one wants to integrate  $\frac{f(x)}{g(x)}$  where  $f$  and  $g$  are polynomials

and  $[\text{degree of } f] \geq [\text{degree of } g]$ , then one must first do  $\underline{\hspace{2cm}}$

**1q.** Integration by parts formula:  $\int u dv = \underline{\hspace{2cm}}$

**1r.** Trig substitution: (recall that the *integrand* is the function you are integrating)  
if the integrand involves  $a^2-u^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1s.** Trig substitution:  
if the integrand involves  $a^2+u^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1t.** Trig substitution:  
if the integrand involves  $u^2-a^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1u.** trig formula ... your answer should involve trig functions of  $\theta$ , and not of  $2\theta$ :  $\sin(2\theta) = \underline{\hspace{2cm}}$ .

**1v.** trig formula ...  $\cos(2\theta)$  should appear in the numerator:  $\cos^2(\theta) = \frac{\underline{\hspace{2cm}}}{2}$ .

**1w.** trig formula ...  $\cos(2\theta)$  should appear in the numerator:  $\sin^2(\theta) = \frac{\underline{\hspace{2cm}}}{2}$ .

**1x.** trig formula ... since  $\cos^2 \theta + \sin^2 \theta = 1$ , we know that the corresponding relationship between tangent (i.e., tan) and secant (i.e., sec) is  $\underline{\hspace{2cm}}$ .

**1y.**  $\arctan(-\sqrt{3}) = \underline{\hspace{2cm}}$  **RADIANS**, not degrees.

2. Fill-in-the blanks/boxes. All series  $\sum$  are understood to be  $\sum_{n=1}^{\infty}$ .

**Hint: I do NOT want to see the words absolute nor conditional on this page!**

**2a. Sequences** Let  $-\infty < r < \infty$ . (Fill-in-the blanks with *exists* or *does not exist*, i.e. *DNE*)

- If  $|r| < 1$ , then  $\lim_{n \rightarrow \infty} r^n$  \_\_\_\_\_
- If  $|r| > 1$ , then  $\lim_{n \rightarrow \infty} r^n$  \_\_\_\_\_
- If  $r = 1$ , then  $\lim_{n \rightarrow \infty} r^n$  \_\_\_\_\_
- If  $r = -1$ , then  $\lim_{n \rightarrow \infty} r^n$  \_\_\_\_\_

**2b. Geometric Series** where  $-\infty < r < \infty$ . The series  $\sum r^n$

- converges if and only if  $|r|$  \_\_\_\_\_
- diverges if and only if  $|r|$  \_\_\_\_\_

**2c.  $p$ -series** where  $0 < p < \infty$ . The series  $\sum \frac{1}{n^p}$

- converges if and only if  $p$  \_\_\_\_\_
- diverges if and only if  $p$  \_\_\_\_\_

**2d. Integral Test** for a positive-termed series  $\sum a_n$  where  $a_n \geq 0$ .

Let  $f: [1, \infty) \rightarrow \mathbb{R}$  be so that

- $a_n = f(\text{_____})$  for each  $n \in \mathbb{N}$
- $f$  is a \_\_\_\_\_ function
- $f$  is a \_\_\_\_\_ function
- $f$  is a \_\_\_\_\_ function .

Then  $\sum a_n$  converges if and only if \_\_\_\_\_ converges.

**2e. Comparison Test** for a positive-termed series  $\sum a_n$  where  $a_n \geq 0$ .

- If  $0 \leq a_n \leq b_n$  for all  $n \in \mathbb{N}$  and  $\sum b_n$  \_\_\_\_\_, then  $\sum a_n$  \_\_\_\_\_.
- If  $0 \leq b_n \leq a_n$  for all  $n \in \mathbb{N}$  and  $\sum b_n$  \_\_\_\_\_, then  $\sum a_n$  \_\_\_\_\_.

**2f. Limit Comparison Test** for a positive-termed series  $\sum a_n$  where  $a_n \geq 0$ .

Let  $b_n > 0$  and  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L$ .

If \_\_\_\_\_  $< L <$  \_\_\_\_\_, then  $\sum a_n$  converges if and only if \_\_\_\_\_ .

**2g. Ratio and Root Tests** for a positive-termed series  $\sum a_n$  where  $a_n \geq 0$ .

Let  $\rho = \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$  or  $\rho = \lim_{n \rightarrow \infty} (a_n)^{\frac{1}{n}}$ .

- If  $\rho$  \_\_\_\_\_ then  $\sum a_n$  converges.
- If  $\rho$  \_\_\_\_\_ then  $\sum a_n$  diverges.
- If  $\rho$  \_\_\_\_\_ then the test is inconclusive.

**2h. Alternating Series Test** for an alternating series  $\sum (-1)^n a_n$  where  $a_n > 0$  for each  $n \in \mathbb{N}$ .

If

- $a_n$  \_\_\_\_\_  $a_{n+1}$  for each  $n \in \mathbb{N}$
- $\lim_{n \rightarrow \infty} a_n =$  \_\_\_\_\_

then  $\sum (-1)^n a_n$  \_\_\_\_\_

**2i.  $n^{\text{th}}$ -term test** for an arbitrary series  $\sum a_n$ .

If  $\lim_{n \rightarrow \infty} a_n \neq 0$  or  $\lim_{n \rightarrow \infty} a_n$  does not exist, then  $\sum a_n$  \_\_\_\_\_ .


2j. By definition, for an arbitrary series  $\sum a_n$ , (fill in the blanks with converges or diverges).

- $\sum a_n$  is absolutely convergent if and only if  $\sum |a_n|$  \_\_\_\_\_
- $\sum a_n$  is conditionally convergent if and only if  $\sum a_n$  \_\_\_\_\_ and  $\sum |a_n|$  \_\_\_\_\_
- $\sum a_n$  is divergent if and only if  $\sum a_n$  \_\_\_\_\_

2k. Fill in the 3 blank lines, with

**absolutely convergent, conditional convergent, or divergent,**

on the following FLOW CHART for class used to determine if a series  $\sum_{n=17}^{\infty} a_n$  is: absolutely convergent, conditional convergent, or divergent.



2l. Fill in the blank with: **perpendicular or parallel**.

If you revolve a graph of a function about an axis of revolution and you want to find the volume of the resulting solid of revolution using the **disk or washer method**, then you draw in your typical rectangles (used to form your typical elements) \_\_\_\_\_ to the axis of revolution.

2m. Fill in the blank with a formula involving *some of*:

$2, \pi, \text{radius}, \text{radius}_{\text{big}}, \text{radius}_{\text{little}}, \text{average radius}, \text{height}, \text{and/or thickness}.$

In problem 2l above, if you use the **washer method**, then the volume of a typical washer is:

\_\_\_\_\_ .

2n. Fill in the blank with: **perpendicular or parallel**.

If you revolve a graph of a function about an axis of revolution and you want to find the volume of the resulting solid of revolution using the **shell method**, then you draw in your typical rectangles (used to form your typical elements) \_\_\_\_\_ to the axis of revolution.

2o. Fill in the blank with a formula involving *some of*:

$2, \pi, \text{radius}, \text{radius}_{\text{big}}, \text{radius}_{\text{little}}, \text{average radius}, \text{height}, \text{and/or thickness}.$

In problem 2n above, if you use the **shell method**, then the volume of a typical shell is:

\_\_\_\_\_ .

3. Circle T if the statement is (always) TRUE. Circle F if the statement is (sometimes) FALSE.

- |   |   |   |
|---|---|---|
| T | F | If $f(n) = a_n$ for $n = 1, 2, 3, \dots$ and if $\lim_{x \rightarrow \infty} f(x) = L$ then $\lim_{n \rightarrow \infty} a_n = L$ .   |
| T | F | If $f(n) = a_n$ for $n = 1, 2, 3, \dots$ and if $\lim_{n \rightarrow \infty} a_n = L$ , then $\lim_{x \rightarrow \infty} f(x) = L$ . |
| T | F | If $0 < a_n < 1$ , then $\lim_{n \rightarrow \infty} a_n$ exists.   |
| T | F | If $0 < a_n < 1$ , then $\sum a_n$ exists.  |
| T | F | If $\lim_{n \rightarrow \infty} a_n = 0$ , then $\sum a_n$ converges  |
| T | F | If $\sum a_n$ converges, then $\lim_{n \rightarrow \infty} a_n = 0$ .   |
| T | F | If $\sum (a_n + b_n)$ converges, then $\sum a_n$ converges and $\sum b_n$ converge.   |
| T | F | If $\sum a_n$ converges and $\sum b_n$ converge, then $\sum (a_n + b_n)$ converges.   |
| T | F | If $S_N = \sum_{n=2}^{N-1} r^n$ , then $S_N = \frac{r^2 - r^N}{1 - r}$ . Notice that the sum is from <b>2</b> to <b>N-1</b>           |
| T | F | Calculus is fun! (each <b>honest</b> solution is correct!)  |

**For problem 4 and 5:** Let  $R$  be the region in the first quadrant of the  $xy$ -plane enclosed by:

- the parabola  $y = 3x^2$
- the  $x$ -axis
- the vertical line  $x = 2$ .

Let  $V$  be the volume of the solid obtained by revolving the region  $R$  about the  $y$ -axis.

**Please remind Prof. Girardi to sketch  $R$  on the board for you.**

4. Using the **disk or washer method**, express the volume  $V$  as an integral.

$V =$

You do not have to evaluate the integral nor do lot of algebra to the integrand.

5. Using the **shell method**, express the volume  $V$  as an integral.

$V =$

You do not have to evaluate the integral nor do lot of algebra to the integrand.

6.

$$\int \sec^4 x \tan^4 x \, dx =$$

$+C$

Hint: problem 1x.

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

7.

$$\int \frac{dx}{(4+x^2)^2} =$$

$+C$

Hint: problem 1s, 1u, 1v, 1w.

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).



8.

$$\int \frac{2x^2 - 2x - 1}{x^2(x-1)} dx =$$

$+C$

Hint:  $x^2 = (x - 0)^2 = (\text{a linear term})^2 \neq (\text{an irreducible quadratic})^1$ .

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

9.

$$\int x^2 e^x dx =$$

$+C$

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

10.

$$\int \ln(x + 17) dx =$$

$+C$

Hint: some cleverness in your choice of  $v$  can make life easier.

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

11.

$$\int e^{(2x)} \cos(3x) dx = \qquad \qquad \qquad +C$$

Hint: bring to the other side idea.

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

12. Sequences (not series)

12a.

$$\lim_{n \rightarrow \infty} \frac{\ln(n^2)}{n} =$$

12b.

$$\lim_{n \rightarrow \infty} \frac{\sqrt{4n^3 + 7n + 5}}{7n^{\frac{3}{2}} + 8} =$$

**NOW Series - NOT sequences**

**13.** Check the correct box and then indicate your reasoning below. Specifically specify what test(s) you are using. A correctly checked box without appropriate explanation will receive no points.

$$\sum_{n=17}^{\infty} \frac{(-1)^n}{n}$$

absolutely convergent

conditionally convergent

divergent

14. Check the correct box and then indicate your reasoning below. Specifically specify what test(s) you are using. A correctly checked box without appropriate explanation will receive no points.

$$\sum_{n=17}^{\infty} \frac{1}{\sqrt{(n-1)(n-2)(n-3)}}$$

absolutely convergent

conditionally convergent

divergent

15. Consider the formal power series

$$\sum_{n=1}^{\infty} \frac{(2x - 12)^n}{n}$$

Hint:  $(2x - 12)^n = [2(x - 6)]^n = 2^n (x - 6)^n$  .

The center is  $x_0 =$  \_\_\_\_\_ and the radius of convergence is  $R =$  \_\_\_\_\_ .

As we did in class, make a number line indicating where the power series is:

**absolutely convergent, conditionally convergent, and divergent.**

Indicate your reasoning and specifically specify what test(s) you are using. Don't forget to check the endpoints, if there are any.





16. Consider the formal power series

$$\sum_{n=2}^{\infty} \frac{(n!) x^n}{2^n}.$$

The center is  $x_0 =$  \_\_\_\_\_ and the radius of convergence is  $R =$  \_\_\_\_\_.

As we did in class, make a number line indicating where the power series is:

**absolutely convergent, conditionally convergent, and divergent.**

Indicate your reasoning and specifically specify what test(s) you are using. Don't forget to check the endpoints, if there are any.



17. Do parts (a) - (f) for the following:

$$f(x) = xe^x \quad x_0 = 0 \quad J = (-17, 2)$$

You might find it easier to do problems (a) - (f) in a different order. Just do what you find easiest.

Use only:

- the definition of Taylor polynomial
- the definition of Taylor series
- the theorem/error-estimate on the  $N^{\text{th}}$ -Remainder term for Taylor polynomials.

Do **NOT** use a known Taylor Series (i.e., do not use methods from section 10.10).

17a Find the following. Note the 1<sup>st</sup> column are functions of  $x$  and the 2<sup>nd</sup> and 3<sup>rd</sup> columns are numbers

$f^{(0)}(x) =$	$f^{(0)}(x_0) =$	$c_0 =$
$f^{(1)}(x) =$	$f^{(1)}(x_0) =$	$c_1 =$
$f^{(2)}(x) =$	$f^{(2)}(x_0) =$	$c_2 =$
$f^{(3)}(x) =$	$f^{(3)}(x_0) =$	$c_3 =$
$f^{(4)}(x) =$	$f^{(4)}(x_0) =$	$c_4 =$
$f^{(5)}(x) =$	nothing for here	nothing for here

17b Find the  $N^{\text{th}}$ -order Taylor polynomial of  $y = f(x)$  about  $x_0$  in OPEN form for  $N = 0, 1, 2, 3, 4$ .

$P_0(x) =$
$P_1(x) =$
$P_2(x) =$
$P_3(x) =$
$P_4(x) =$

**17c.** Find the Taylor series of  $y = f(x)$  about  $x_0$  in OPEN form.

$$P_{\infty}(x) =$$

**17d.** Find the Taylor series of  $y = f(x)$  about  $x_0$  in CLOSED form.

$$P_{\infty}(x) =$$

**17e.** Consider the given interval  $J$ . Find an upper bound for the maximum of  $|f^{(5)}(x)|$  on the interval  $J$ .

You answer should be a number. You answer cannot have an:  $N, x, x_0, c$ .

$$\max_{c \in J} |f^{(5)}(c)| \leq$$

**17f.** Consider the given interval  $J$ . Using Taylor's Remainder Theorem (i.e., Taylor's Big Theorem), find an upper bound for the maximum of  $|R_4(x)|$  on the interval  $J$ . You answer should be a number.

You answer cannot have an:  $N, x, x_0, c$ .

$$\max_{x \in J} |R_4(x)| \leq$$

18. Using the fact that

$$\frac{1}{1-r} = \sum_{n=0}^{\infty} r^n \quad \text{when} \quad |r| < 1, \quad (*)$$

find a power series expansion of

$$\frac{x}{4+100x^2}$$

and state when it is valid. Simplify your answer so that your power series has the form

$\sum_{n=0}^{\infty} c_n x^{\text{some power}}$  for some constants  $c_n$ .

$$\frac{x}{4+100x^2} = \sum_{n=0}^{\infty} c_n x^n$$

valid when  $|x| < \frac{1}{10}$