

MARK BOX		
PROBLEM	POINTS	
1	25	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
take home	10	
Extra Credit	5	
%	100	

NAME: _____

class PIN: _____

(*) Extra Credit: 5 point for knowing your PIN number.

INSTRUCTIONS:

- (1) To receive credit you must:
 - (a) **work in a logical fashion, show all your work, indicate your reasoning; no credit will be given for an answer that just appears;**
such explanations help with partial credit
 - (b) if a line/box is provided, then:
 - show you work BELOW the line/box
 - put your answer on/in the line/box
 - (c) if no such line/box is provided, then box your answer
- (2) The MARK BOX indicates the problems along with their points. Check that your copy of the exam has all of the problems.
- (3) You may **not** use an electronic device, a calculator, books, personal notes.
- (4) During this exam, do not leave your seat. If you have a question, raise your hand. When you finish: turn your exam over, put your pencil down, and raise your hand.
- (5) This exam covers (from *Calculus* (ET) by Stewart 6th ed.):
Sections 7.1 – 7.5, 7.8. 11.1 .

Problem Inspiration:

1. You were warned.
2. example in class
3. homework problem § 7.5 # 9
4. homework problem § 7.5 # 15
5. homework problem § 7.5 # 21
6. Handout of 100 integrals # 35
7. homework problem § 7.5 # 41

Hints:

- (1) **You can check your answers to the indefinite integrals by differentiating.**
- (2) **For more partial credit, box your $u - du$ substitutions.**

Honor Code Statement

I understand that it is the responsibility of every member of the Carolina community to uphold and maintain the University of South Carolina's Honor Code.

As a Carolinian, I certify that I have neither given nor received unauthorized aid on this exam.

Furthermore, I have not only read but will also follow the above Instructions.

Signature : _____

1. Fill in the blanks (each worth 1 point).

- $\int \frac{du}{u} = \underline{\hspace{2cm}} |u| + C$
- If a is a constant and $a > 0$ but $a \neq 1$, then $\int a^u du = \underline{\hspace{2cm}} + C$
- $\int \cos u du = \underline{\hspace{2cm}} + C$
- $\int \sin u du = \underline{\hspace{2cm}} + C$
- $\int \tan u du = \underline{\hspace{2cm}} + C$
- $\int \cot u du = \underline{\hspace{2cm}} + C$
- $\int \sec u du = \underline{\hspace{2cm}} + C$
- $\int \csc u du = \underline{\hspace{2cm}} + C$
- $\int \sec^2 u du = \underline{\hspace{2cm}} + C$
- $\int \sec u \tan u du = \underline{\hspace{2cm}} + C$
- $\int \csc^2 u du = \underline{\hspace{2cm}} + C$
- $\int \csc u \cot u du = \underline{\hspace{2cm}} + C$
- If a is a constant and $a > 0$ then $\int \frac{1}{a^2+u^2} du = \underline{\hspace{2cm}} + C$
- If a is a constant and $a > 0$ then $\int \frac{1}{\sqrt{a^2-u^2}} du = \underline{\hspace{2cm}} + C$
- If a is a constant and $a > 0$ then $\int \frac{1}{u\sqrt{u^2-a^2}} du = \underline{\hspace{2cm}} + C$
- Partial Fraction Decomposition. If one wants to integrate $\frac{f(x)}{g(x)}$ where f and g are polynomials and $[\text{degree of } f] \geq [\text{degree of } g]$, then one must first do $\underline{\hspace{2cm}}$
- Integration by parts formula: $\int u dv = \underline{\hspace{2cm}}$
- Trig substitution: (recall that the *integrand* is the function you are integrating) if the integrand involves a^2+u^2 , then one makes the substitution $u = \underline{\hspace{2cm}}$
- Trig substitution: if the integrand involves a^2-u^2 , then one makes the substitution $u = \underline{\hspace{2cm}}$
- Trig substitution: if the integrand involves u^2-a^2 , then one makes the substitution $u = \underline{\hspace{2cm}}$
- trig formula ... your answer should involve trig functions of θ , and not of 2θ : $\sin(2\theta) = \underline{\hspace{2cm}}$.
- trig formula ... your answer should have $\cos(2\theta)$ in it: $\cos^2(\theta) = \frac{1}{2} (\underline{\hspace{2cm}})$.
- trig formula ... your answer should have $\cos(2\theta)$ in it: $\sin^2(\theta) = \frac{1}{2} (\underline{\hspace{2cm}})$.
- trig formula ... since $\cos^2 \theta + \sin^2 \theta = 1$, we know that the corresponding relationship between tangent (i.e., tan) and secant (i.e., sec) is $\underline{\hspace{2cm}}$.
- $\arctan(-1) = \underline{\hspace{2cm}}$ **RADIANS**. (your answer should be an angle)

2.

$$\int \sin^2 x \, dx =$$

+ C

Hint: trig formulas from problem 1 come in handy.

3.

$$\int x^{\frac{3}{2}} \ln x \, dx =$$

+ C

4.

$$\int \frac{x-1}{x^2+2x} dx = \qquad \qquad \qquad + C$$

Hint: $x^2 + 2x = x(x + 2) = (x - 0)(x + 2)$

5a. Complete the square. The two lines should have numbers on them. The box should have a plus or minus sign in it.

$$x^2 - 4x = (x - \underline{\hspace{2cm}})^2 \boxed{\hspace{1cm}} \underline{\hspace{2cm}}$$

5b.

$\int \frac{1}{\sqrt{x^2 - 4x}} dx =$	$+ C$
---------------------------------------	-------

6.

$$\int \sec^3 x \tan^3 x \, dx =$$

+ C

7.

$$\int_1^{\infty} \frac{1}{(2x+1)^3} dx =$$

Warning: write your solution in proper form.