

MARK BOX		
PROBLEM	POINTS	
1	25	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
take home	10	
Extra Credit	5	
%	100	

NAME: _____

Soln Key

class PIN: _____

17

(*) Extra Credit: 5 point for knowing your PIN number.

INSTRUCTIONS:

- (1) To receive credit you must:
 - (a) **work in a logical fashion, show all your work, indicate your reasoning;**
no credit will be given for an answer that just appears;
such explanations help with partial credit
 - (b) if a line/box is provided, then:
 - show you work BELOW the line/box
 - put your answer on/in the line/box
 - (c) if no such line/box is provided, then box your answer
- (2) The MARK BOX indicates the problems along with their points.
Check that your copy of the exam has all of the problems.
- (3) You may **not** use an electronic device, a calculator, books, personal notes.
- (4) During this exam, do not leave your seat. If you have a question, raise your hand. When you finish: turn your exam over, put your pencil down, and raise your hand.
- (5) This exam covers (from *Calculus* (ET) by Stewart 6th ed.):
Sections 7.1 – 7.5, 7.8. 11.1 .

Problem Inspiration:

1. You were warned.
2. example in class
3. homework problem § 7.5 # 9
4. homework problem § 7.5 # 15
5. homework problem § 7.5 # 21
6. Handout of 100 integrals # 35
7. homework problem § 7.5 # 41

Hints:

- (1) **You can check your answers to the indefinite integrals by differentiating.**
- (2) **For more partial credit, box your $u - du$ substitutions.**

Honor Code Statement

I understand that it is the responsibility of every member of the Carolina community to uphold and maintain the University of South Carolina's Honor Code.

As a Carolinian, I certify that I have neither given nor received unauthorized aid on this exam.

Furthermore, I have not only read but will also follow the above Instructions.

Signature : _____


You were warned on the first day of the semester.

1. Fill in the blanks (each worth 1 point).

- $\int \frac{du}{u} = \ln |u| + C$
- If a is a constant and $a > 0$ but $a \neq 1$, then $\int a^u du = \frac{a^u}{\ln a} + C$
- $\int \cos u du = \sin u + C$
- $\int \sin u du = -\cos u + C$
- $\int \tan u du = \ln |\sec u|$ or $-\ln |\cos u| + C$
- $\int \cot u du = \ln |\sin u|$ or $-\ln |\csc u| + C$
- $\int \sec u du = \ln |\sec u + \tan u|$ or $-\ln |\sec u - \tan u| + C$
- $\int \csc u du = \ln |\csc u - \cot u|$ or $-\ln |\csc u + \cot u| + C$
- $\int \sec^2 u du = \tan u + C$
- $\int \sec u \tan u du = \sec u + C$
- $\int \csc^2 u du = -\cot u + C$
- $\int \csc u \cot u du = -\csc u + C$
- If a is a constant and $a > 0$ then $\int \frac{1}{a^2+u^2} du = \frac{1}{a} \tan^{-1} \frac{u}{a} + C$
- If a is a constant and $a > 0$ then $\int \frac{1}{\sqrt{a^2-u^2}} du = \sin^{-1} \frac{u}{a} + C$
- If a is a constant and $a > 0$ then $\int \frac{1}{u\sqrt{u^2-a^2}} du = \frac{1}{a} \sec^{-1} \frac{|u|}{a} + C$
- Partial Fraction Decomposition. If one wants to integrate $\frac{f(x)}{g(x)}$ where f and g are polynomials and $[\text{degree of } f] \geq [\text{degree of } g]$, then one must first do long division
- Integration by parts formula: $\int u dv = uv - \int v du$
- Trig substitution: (recall that the *integrand* is the function you are integrating) if the integrand involves a^2+u^2 , then one makes the substitution $u = a \tan \theta$
- Trig substitution: if the integrand involves a^2-u^2 , then one makes the substitution $u = a \sin \theta$
- Trig substitution: if the integrand involves u^2-a^2 , then one makes the substitution $u = a \sec \theta$
- trig formula ... your answer should involve trig functions of θ , and not of 2θ : $\sin(2\theta) = 2 \sin \theta \cos \theta$
- trig formula ... your answer should have $\cos(2\theta)$ in it: $\cos^2(\theta) = \frac{1}{2} (1 + \cos 2\theta)$
- trig formula ... your answer should have $\cos(2\theta)$ in it: $\sin^2(\theta) = \frac{1}{2} (1 - \cos 2\theta)$
- trig formula ... since $\cos^2 \theta + \sin^2 \theta = 1$, we know that the corresponding relationship between tangent (i.e., \tan) and secant (i.e., \sec) is $1 + \tan^2 \theta = \sec^2 \theta$
- $\arcsin \frac{1}{2} = \frac{\pi}{6}$ RADIANS. (your answer should be an angle)

$s + c$

$$\arctan(-1) = -\frac{\pi}{4} \neq 1$$



$$\frac{c \phi + a}{n}$$

2.

$$\int \sin^2 x \, dx = \frac{x}{2} - \frac{\sin(2x)}{4} + C$$

Hint: trig formulas from problem 1 come in handy.

$$\int \sin^2 x \, dx = \frac{1}{2} \int (1 - \cos 2x) \, dx$$

$$= \frac{1}{2} \int 1 \, dx - \frac{1}{2} \int \cos 2x \, dx$$

$$= \frac{1}{2} \int 1 \, dx - \frac{1}{2} \cdot \frac{1}{2} \int \cos 2x (2 \, dx)$$

$$= \frac{1}{2} \cdot x - \frac{1}{4} \sin(2x) + C$$

3.

$$\int x^{3/2} \ln x \, dx = \frac{2x^{5/2} \ln x}{5} - \frac{4}{25} x^{5/2} + C$$

$$\begin{array}{l} u = \ln x \\ du = \frac{1}{x} dx \end{array} \quad \begin{array}{l} dv = x^{3/2} dx \\ v = \frac{2}{5} x^{5/2} \end{array}$$

$$\int x^{3/2} \ln x \, dx = \frac{2x^{5/2} \ln x}{5} - \frac{2}{5} \int x^{5/2} \cdot x^{-1} dx$$

$$= \frac{2x^{5/2} \ln x}{5} - \frac{2}{5} \int x^{3/2} dx$$

$$= \frac{2x^{5/2} \ln x}{5} - \frac{2}{5} \cdot \frac{2}{5} x^{5/2} + C$$

4.

$$\int \frac{x-1}{x^2+2x} dx = -\frac{1}{2} \ln|x| + \frac{3}{2} \ln|x+2| + C$$

Hint: $x^2 + 2x = x(x+2) = (x-0)(x+2)$

$$\frac{x-1}{x(x+2)} = \frac{A}{x} + \frac{B}{x+2} = \frac{A(x+2) + Bx}{x(x+2)}$$

$$\Rightarrow x-1 = A(x+2) + Bx \quad \begin{cases} x=0 & -1 = 2A \\ x=-2 & -3 = -2B \end{cases} \Rightarrow \begin{cases} A = -\frac{1}{2} \\ B = \frac{3}{2} \end{cases}$$

or can equate coefficients

$$x: 1 = A + B$$

$$\text{constant: } -1 = 2A$$

$$\text{So } \int \frac{x-1}{x^2+2x} dx = \int \frac{-1/2}{x} dx + \int \frac{3/2}{x+2} dx$$

$$= -\frac{1}{2} \ln|x| + \frac{3}{2} \ln|x+2| + C$$

$$\text{or } \ln|x|^{-1/2} + \ln|x+2|^{3/2} + C$$

$$\text{or } \ln \left[|x|^{-1/2} |x+2|^{3/2} \right] + C$$

$$\text{or } \ln \left[\frac{|x+2|^{3/2}}{|x|^{1/2}} \right] + C \quad \text{or etc....}$$

5a. Complete the square. The two lines should have numbers on them. The box should have a plus or minus sign in it.

$$x^2 - 4x = (x - \underline{2})^2 \boxed{-} \underline{4}$$

↓
x - 4x + 4

5b.


$$\int \frac{1}{\sqrt{x^2 - 4x}} dx = \ln \left| \frac{x-2}{2} + \frac{\sqrt{x^2 - 4x}}{2} \right| + C$$

$$\int \frac{dx}{\sqrt{x^2 - 4x}} = \int \frac{dx}{\sqrt{(x-2)^2 - 2^2}} = \int \frac{2 \sec \theta \tan \theta d\theta}{2 \tan \theta}$$

$$= \int \sec \theta d\theta$$

$$= \ln |\sec \theta + \tan \theta| + C$$

$x - 2 = 2 \sec \theta$
 $dx = 2 \sec \theta \tan \theta d\theta$
 $\sqrt{(x-2)^2 - 2^2} = \sqrt{4 \sec^2 \theta - 4}$
 $= 2 \sqrt{\sec^2 \theta - 1}$
 $= 2 \tan \theta$
 $\sec \theta = \frac{x-2}{2}$



6.

$$\int \sec^3 x \tan^3 x \, dx =$$

+ C

$$\textcircled{35} \int \sec^3 x \tan^3 x \, dx = \int \sec^2 x \tan^2 x (\sec x \tan x \, dx)$$

$$\begin{array}{l} u = \tan x \\ du = \sec^2 x \, dx \end{array}$$

get in terms of v
or

$$\begin{array}{l} v = \sec x \\ dv = \sec x \tan x \, dx \end{array}$$

$$= \int \sec^2 x (\sec^2 x - 1) (\sec x \tan x \, dx)$$

$$= \int v^2 (v^2 - 1) \, dv = \int (v^4 - v^2) \, dv = \frac{v^5}{5} - \frac{v^3}{3} + C$$

$$= \frac{\sec^5 x}{5} - \frac{\sec^3 x}{3} + C$$

7.

$$\int_1^{\infty} \frac{1}{(2x+1)^3} dx = \frac{1}{36}$$

Warning: write your sol'n in proper form.

$$\int_1^{\infty} \frac{dx}{(2x+1)^3} = \lim_{b \rightarrow \infty} \int_{x=1}^{x=b} \frac{1}{2} (2x+1)^{-3} (2dx)$$

$$= \lim_{b \rightarrow \infty} \frac{1}{2} \frac{(2x+1)^{-2}}{-2} \Big|_{x=1}^{x=b}$$

$$= -\frac{1}{4} \lim_{b \rightarrow \infty} \frac{1}{(2x+1)^2} \Big|_{x=1}^{x=b}$$

$$= -\frac{1}{4} \left[\lim_{b \rightarrow \infty} \left[\frac{1}{(2b+1)^2} - \frac{1}{3^2} \right] \right]$$

$$= -\frac{1}{4} \left[0 - \frac{1}{3^2} \right]$$

$$= \frac{1}{4} \cdot \frac{1}{9}$$