

If you do not make at least a 50% on this exam's *first few problems*, then your **total** exam score will be only as many points as you managed to get on the *first few problems*. Here are some typical *first few problems*.

1. Fill-in-the blanks/boxes. All series \sum are understood to be $\sum_{n=1}^{\infty}$.

Hint: You should NOT write the words absolute nor conditional on Problem 1!

- 1a. **Sequences** (Afterall, this is needed for Geometric Series!)

Let $-\infty < r < \infty$. (Fill-in-the blanks with *exists* or *does not exist*, i.e. *DNE*)

- If $|r| < 1$, then $\lim_{n \rightarrow \infty} r^n$ exists (= 0)
- If $|r| > 1$, then $\lim_{n \rightarrow \infty} r^n$ DNE
- If $r = 1$, then $\lim_{n \rightarrow \infty} r^n$ exists (= 1)
- If $r = -1$, then $\lim_{n \rightarrow \infty} r^n$ DNE (osc.)

- 1b. **Geometric Series** where $-\infty < r < \infty$. The series $\sum r^n$

- converges if and only if $|r|$ < 1
- diverges if and only if $|r|$ ≥ 1

- 1c. **p-series** where $0 < p < \infty$. The series $\sum \frac{1}{n^p}$

- converges if and only if p > 1
- diverges if and only if p ≤ 1

- 1d. **Integral Test** for a positive-termed series $\sum a_n$ where $a_n \geq 0$.

Let $f: [1, \infty) \rightarrow \mathbb{R}$ be so that

- $a_n = f(\underline{n})$ for each $n \in \mathbb{N}$
- f is a positive function
- f is a continuous function
- f is a decreasing (or non increasing) function.

Then $\sum a_n$ converges if and only if $\int_1^{\infty} f(x) dx$ converges.

- 1e. **Comparison Test** for a positive-termed series $\sum a_n$ where $a_n \geq 0$.

- If $0 \leq a_n \leq b_n$ for all $n \in \mathbb{N}$ and $\sum b_n$ converges, then $\sum a_n$ converges.
- If $0 \leq b_n \leq a_n$ for all $n \in \mathbb{N}$ and $\sum b_n$ diverges, then $\sum a_n$ diverges.

- 1f. **Limit Comparison Test** for a positive-termed series $\sum a_n$ where $a_n \geq 0$.

Let $b_n > 0$ and $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L$.

If 0 < L < ∞ , then $\sum a_n$ converges if and only if $\sum b_n$ converges

- 1g. **Ratio and Root Tests** for a positive-termed series $\sum a_n$ where $a_n \geq 0$.

Let $\rho = \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$ or $\rho = \lim_{n \rightarrow \infty} (a_n)^{\frac{1}{n}}$.

- If ρ < 1 then $\sum a_n$ converges.
- If ρ > 1 then $\sum a_n$ diverges.
- If ρ = 1 then the test is inconclusive.

1h. **Alternating Series Test** for an alternating series $\sum (-1)^n a_n$ where $a_n > 0$ for each $n \in \mathbb{N}$.

If

- $a_n > a_{n+1}$ for each $n \in \mathbb{N}$ (decreasing)
- $\lim_{n \rightarrow \infty} a_n = 0$

then $\sum (-1)^n a_n$ converges

1i. **n^{th} -term test** for an arbitrary series $\sum a_n$.

If $\lim_{n \rightarrow \infty} a_n \neq 0$ or $\lim_{n \rightarrow \infty} a_n$ does not exist, then $\sum a_n$ diverges.

1j. By definition, for an arbitrary series $\sum a_n$, (fill in the blanks with converges or diverges).

- $\sum a_n$ is absolutely convergent if and only if $\sum |a_n|$ converges
- $\sum a_n$ is conditionally convergent if and only if $\sum a_n$ converges and $\sum |a_n|$ diverges
- $\sum a_n$ is divergent if and only if $\sum a_n$ diverges

1k. If a power series in $x - x_0$ has radius of convergence R where $0 < R < \infty$, then the power series is:

- absolutely convergent for $x_0 - R < x < x_0 + R$
- divergent for $x < x_0 - R$ and $x_0 + R < x$

1l. Consider a **function** $y = f(x)$ where $f: [1, \infty) \rightarrow \mathbb{R}$.

Next consider the corresponding **sequence** $\{a_n\}_{n=1}^{\infty}$ where $a_n \stackrel{\text{def.}}{=} f(n)$.

- If the limit of the **function** $y = f(x)$ as $x \rightarrow \infty$ is L ,

then the limit of the corresponding **sequence** $\{a_n\}_{n=1}^{\infty}$ as $n \rightarrow \infty$ is L .

- If $\lim_{n \rightarrow \infty} a_n = L$, is it necessarily true that $\lim_{x \rightarrow \infty} f(x) = L$? Circle: **Yes** or **(No)**

2. Fill in the 3 blank lines (with absolutely convergent, conditional convergent, or divergent) on the following FLOW CHART for class used to determine if a series $\sum_{n=17}^{\infty} a_n$ is: absolutely convergent, conditional convergent, or divergent.

Does $\sum |a_n|$ converge?
Since $|a_n| \geq 0$, use a positive term test:
integral test, CT, LCT, ratio/root test.

if YES \downarrow

$\sum a_n$ is absolutely convergent

if NO \Rightarrow

Does $\lim_{n \rightarrow \infty} |a_n| = 0$?

if NO \Rightarrow

$\sum a_n$ is divergent

if YES \downarrow

Is $\sum a_n$ an alternating series?

if YES \downarrow

Does $\sum a_n$ satisfy the conditions of the Alternating Series Test?

if YES \downarrow

$\sum a_n$ is conditionally convergent

3. Circle T if the statement is TRUE. Circle F if the statement is FALSE. To be more specific: circle T if the statement is always true and circle F if the statement is NOT always true.
Scoring: 2 pts for a correct answer, 1 pt for a blank answer, 0 pts for an incorrect answer.

- (T) F If $\sum a_n$ converges, then $\lim_{n \rightarrow \infty} a_n = 0$.
- T (F) If $\lim_{n \rightarrow \infty} a_n = 0$, then $\sum a_n$ converges. (eg, $a_n = \frac{1}{n}$)
- T (F) If a sequence $\{a_n\}_{n=1}^{\infty}$ satisfies that $\lim_{n \rightarrow \infty} a_n = L$ and $f: [0, \infty) \rightarrow \mathbb{R}$ is a function satisfying that $f(n) = a_n$ for each natural number n , then $\lim_{x \rightarrow \infty} f(x) = L$. Consider $f(x) = \sin(\pi x)$.
- (T) F If a function $f: [0, \infty) \rightarrow \mathbb{R}$ satisfies that $\lim_{x \rightarrow \infty} f(x) = L$ and $\{a_n\}_{n=1}^{\infty}$ is a sequence satisfying that $f(n) = a_n$ for each natural number n , then $\lim_{n \rightarrow \infty} a_n = L$.
- (T) F If $\sum a_n$ converges and $\sum b_n$ converge, then $\sum(a_n + b_n)$ converges.
- T (F) If $\sum(a_n + b_n)$ converges, then $\sum a_n$ converges and $\sum b_n$ converge. e.g. $\begin{cases} a_n = \frac{1}{n} \\ b_n = -\frac{1}{n} \end{cases}$
- (T) F If $r \neq 1$ and $S_N = \sum_{n=17}^N r^n$, then $S_N = \frac{r^{17} - r^{N+1}}{1 - r}$ for each $N > 17$.
NOTICE, the above sum starts at $n = 17$, not at $n = 0$.

$$\begin{aligned} 1 S_N &= r^{17} + r^{18} + r^{19} + \dots + r^N \\ r S_N &= \phantom{r^{17}} + r^{18} + r^{19} + \dots + r^N + r^{N+1} \end{aligned}$$

subtract

$$(1-r) S_N = r^{17} - r^{N+1}$$

MORE \Rightarrow

4. Geometric Series. Let, for $N \geq 102$,

$$s_N = \sum_{n=102}^N 3 \frac{2^n}{7^{n+1}}.$$

4a. Do some algebra to write s_N as $\sum_{n=102}^N c r^n$ for an appropriate constant c and ratio r .

$$s_N = \sum_{n=102}^N \frac{3}{7} \left(\frac{2}{7}\right)^n$$

$$3 \frac{2^n}{7^{n+1}} = 3 \cdot \frac{2^n}{7^n \cdot 7^1} = \frac{3}{7} \frac{2^n}{7^n} = \frac{3}{7} \left(\frac{2}{7}\right)^n$$

4b. Using the method from class (rather than some formula), find an expression for s_N in closed form (i.e. without a summation \sum sign nor some dots ...).

$$s_N = \frac{3}{5} \left[\left(\frac{2}{7}\right)^{102} - \left(\frac{2}{7}\right)^{N+1} \right]$$

$$\begin{aligned} 1 s_N &= \frac{3}{7} \left[\left(\frac{2}{7}\right)^{102} + \left(\frac{2}{7}\right)^{103} + \dots + \left(\frac{2}{7}\right)^N \right] \quad \text{or} \quad \frac{3}{7} \left(\frac{2}{7}\right)^{102} + \frac{3}{7} \left(\frac{2}{7}\right)^{103} + \dots + \frac{3}{7} \left(\frac{2}{7}\right)^N \\ \frac{2}{7} s_N &= \frac{3}{7} \left[\left(\frac{2}{7}\right)^{103} + \dots + \left(\frac{2}{7}\right)^N + \left(\frac{2}{7}\right)^{N+1} \right] \quad \text{or} \quad \frac{3}{7} \left(\frac{2}{7}\right)^{103} + \dots + \frac{3}{7} \left(\frac{2}{7}\right)^N + \frac{3}{7} \left(\frac{2}{7}\right)^{N+1} \end{aligned}$$

$$\left(1 - \frac{2}{7}\right) s_N = \frac{3}{7} \left[\left(\frac{2}{7}\right)^{102} - \left(\frac{2}{7}\right)^{N+1} \right] \quad \text{or} \quad \frac{3}{7} \left(\frac{2}{7}\right)^{102} - \frac{3}{7} \left(\frac{2}{7}\right)^{N+1}$$

$$\frac{5}{7} \Rightarrow s_N = \frac{7}{5} \cdot \frac{3}{7} \left[\left(\frac{2}{7}\right)^{102} - \left(\frac{2}{7}\right)^{N+1} \right]$$

4c. Does $\sum_{n=102}^{\infty} 3 \frac{2^n}{7^{n+1}}$ converge or diverge? If it converges, find its sum. Justify your answer.

$$\sum_{n=102}^{\infty} 3 \frac{2^n}{7^{n+1}} = \frac{3}{5} \left(\frac{2}{7}\right)^{102}$$

Geometric Series, ratio $r = \frac{2}{7}$, $|r| < 1 \Rightarrow$ converges

$$\sum_{n=102}^{\infty} 3 \cdot \frac{2^n}{7^{n+1}} \stackrel{\text{def}}{=} \lim_{N \rightarrow \infty} \sum_{n=102}^N 3 \cdot \frac{2^n}{7^{n+1}} \stackrel{(4b)}{=} \lim_{N \rightarrow \infty} \frac{3}{5} \left[\left(\frac{2}{7}\right)^{102} - \left(\frac{2}{7}\right)^{N+1} \right]$$

$$\stackrel{(1a)}{=} \frac{3}{5} \left[\left(\frac{2}{7}\right)^{102} - 0 \right]$$