

MARK BOX		
PROBLEM	POINTS	
1	25	
2	5	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
TOTAL	90	
%	100	

NAME: _____

class PIN: _____

INSTRUCTIONS:

- (1) To receive credit you must:
 - (a) **work in a logical fashion, show all your work, indicate your reasoning; no credit will be given for an answer that *just appears*; such explanations help with partial credit**
 - (b) if a line/box is provided, then:
 - show you work BELOW the line/box
 - put your answer on/in the line/box
 - (c) if no such line/box is provided, then box your answer
- (2) The MARK BOX indicates the problems along with their points. Check that your copy of the exam has all of the problems.
- (3) You may **not** use an electronic device, a calculator, books, personal notes.
- (4) During this exam, do not leave your seat unless you have permission. If you have a question, raise your hand. When you finish: turn your exam over, put your pencil down, and raise your hand.
- (5) If you do not make at least 12.5 out of 25 points on Problem 1, then your score for the entire exam will be whatever you made on Problem 1.
- (6) This exam covers (from *Calculus* (ET) by Stewart 6th ed.): Sections 7.1 – 7.5, 7.8, 11.1 .

Hints:

- (1) **You can check your answers to the indefinite integrals by differentiating.**
- (2) **For more partial credit, box your $u - du$ substitutions.**

Honor Code Statement

I understand that it is the responsibility of every member of the Carolina community to uphold and maintain the University of South Carolina's Honor Code.

As a Carolinian, I certify that I have neither given nor received unauthorized aid on this exam.

I understand that if it is determined that I used any unauthorized assistance or otherwise violated the University's Honor Code then I will receive a failing grade for this course and be referred to the academic Dean and the Office of Academic Integrity for additional disciplinary actions.

Furthermore, I have not only read but will also follow the above Instructions.

Signature : _____

1. Fill in the blanks (each worth 1 point).

1a. $\int \frac{du}{u} = \underline{\hspace{2cm}} |u| + C$

1b. If a is a constant and $a > 0$ but $a \neq 1$, then $\int a^u du = \underline{\hspace{2cm}} + C$

1c. $\int \cos u du = \underline{\hspace{2cm}} + C$

1d. $\int \sec^2 u du = \underline{\hspace{2cm}} + C$

1e. $\int \sec u \tan u du = \underline{\hspace{2cm}} + C$

1f. $\int \sin u du = \underline{\hspace{2cm}} + C$

1g. $\int \csc^2 u du = \underline{\hspace{2cm}} + C$

1h. $\int \csc u \cot u du = \underline{\hspace{2cm}} + C$

1i. $\int \tan u du = \underline{\hspace{2cm}} + C$

1j. $\int \cot u du = \underline{\hspace{2cm}} + C$

1k. $\int \sec u du = \underline{\hspace{2cm}} + C$

1l. $\int \csc u du = \underline{\hspace{2cm}} + C$

1m. If a is a constant and $a > 0$ then $\int \frac{1}{\sqrt{a^2-u^2}} du = \underline{\hspace{2cm}} + C$

1n. If a is a constant and $a > 0$ then $\int \frac{1}{a^2+u^2} du = \underline{\hspace{2cm}} + C$

1o. If a is a constant and $a > 0$ then $\int \frac{1}{u\sqrt{u^2-a^2}} du = \underline{\hspace{2cm}} + C$

1p. Partial Fraction Decomposition. If one wants to integrate $\frac{f(x)}{g(x)}$ where f and g are polynomials

and $[\text{degree of } f] \geq [\text{degree of } g]$, then one must first do $\underline{\hspace{2cm}}$

1q. Integration by parts formula: $\int u dv = \underline{\hspace{2cm}}$

1r. Trig substitution: (recall that the *integrand* is the function you are integrating)
if the integrand involves a^2-u^2 , then one makes the substitution $u = \underline{\hspace{2cm}}$

1s. Trig substitution:
if the integrand involves a^2+u^2 , then one makes the substitution $u = \underline{\hspace{2cm}}$

1t. Trig substitution:
if the integrand involves u^2-a^2 , then one makes the substitution $u = \underline{\hspace{2cm}}$

1u. trig formula ... your answer should involve trig functions of θ , and not of 2θ : $\sin(2\theta) = \underline{\hspace{2cm}}$.

1v. trig formula ... $\cos(2\theta)$ should appear in the numerator: $\cos^2(\theta) = \frac{1}{2} \underline{\hspace{2cm}}$.

1w. trig formula ... $\cos(2\theta)$ should appear in the numerator: $\sin^2(\theta) = \frac{1}{2} \underline{\hspace{2cm}}$.

1x. trig formula ... since $\cos^2 \theta + \sin^2 \theta = 1$, we know that the corresponding relationship between tangent (i.e., tan) and secant (i.e., sec) is $\underline{\hspace{2cm}}$.

1y. $\arcsin\left(-\frac{\sqrt{2}}{2}\right) = \underline{\hspace{2cm}}$ **RADIANS.** (your answer should be an angle)

2.

$$\int (\sin x) (\sec x) dx =$$

+ C

3.

$$\int x \tan^2 x \, dx =$$

+ C

4.

$$\int \ln(1+x) dx =$$

+ C

Hint: bring to the other side idea.

5.

$$\int \frac{x^2}{\sqrt{4-x^2}} dx =$$

+ C

6.

$$\int \frac{x^4 + 2x + 2}{x^4(x + 1)} dx =$$

+ C

Hint $x^4 = (x - 0)^4$

7.

$$\int_1^{\infty} \frac{1}{(3x+1)^4} dx =$$

Warning: write your solution in proper form.

8. For the following **SEQUENCES**:

- if the limit exists, find it
- if the limit does not exist, then say that it DNE.

Put your ANSWER IN the box and show your WORK BELOW the box.

8a.

$$\lim_{n \rightarrow \infty} \frac{5n^2 + 4\sqrt{n}}{6n^3 + 7n^2 + 1} =$$

8b.

$$\lim_{n \rightarrow \infty} \frac{5n^8 + 4\sqrt{n}}{6n^3 + 7n^2 + 1} =$$

8c.

$$\lim_{n \rightarrow \infty} \frac{5n^3 + 4\sqrt{n}}{6n^3 + 7n^2 + 1} =$$