

Indication of Solutions.

$$1. \int_0^5 \frac{1}{x^2+25} dx = \frac{1}{5} \tan^{-1} \frac{x}{5} \Big|_{x=0}^{x=5} = \frac{1}{5} [\tan^{-1} 1 - \tan^{-1} 0]$$

$$= \frac{1}{5} \left[\frac{\pi}{4} - 0 \right] = \frac{\pi}{20}$$

$$2. \int \frac{x}{x^2+25} dx = \frac{1}{2} \int \frac{du}{u} = \frac{1}{2} \ln |u| + C = \frac{1}{2} \ln |x^2+25| + C$$

$u = x^2 + 25$
 $du = 2x dx$

$$\stackrel{\text{do}}{=} \int_0^5 \frac{x dx}{x^2+25} = \frac{1}{2} \ln |x^2+25| \Big|_0^5 = \frac{1}{2} [\ln 50 - \ln 25]$$

$$\hookrightarrow = \frac{1}{2} \ln \frac{50}{25} = \frac{1}{2} \ln 2 = \ln 2^{1/2} = \ln \sqrt{2}$$

$$3. \int \frac{x^2}{x^2+25} dx \stackrel{\uparrow}{=} \int \left[1 - 5^2 \frac{1}{x^2+5^2} \right] dx = x - 5^2 \cdot \frac{1}{5} \tan^{-1} \frac{x}{5} + C = x - 5 \tan^{-1} \frac{x}{5} + C$$

do not have (strictly) bigger bottoms so need to do long division, which we "fake"

$$\frac{x^2}{x^2+25} = \frac{x^2+25}{x^2+25} + \frac{-25}{x^2+25} = 1 - 25 \frac{1}{x^2+25}$$

$$\text{So } \int_0^5 \frac{x^2}{x^2+25} dx = \left(x - 5 \tan^{-1} \frac{x}{5} \right) \Big|_{x=0}^{x=5} = [5 - 5 \tan^{-1} 1] - [0 - 5 \tan^{-1} 0]$$

$$= [5 - 5(\frac{\pi}{4})] - [0] = \frac{20 - 5\pi}{4}$$

$$4. \int \frac{1-x+2x^2-x^3}{x(x^2+1)^2} dx = \ln |x| - \frac{1}{2} \ln |x^2+1| - \tan^{-1} x - \frac{1}{2(x^2+1)} + C$$

see textbook, § 7.4, Example 8, p 480. Do with PFD.

$$\hookrightarrow \ln \left(\frac{|x|}{(x^2+1)^{1/2}} \right) - \tan^{-1} x - \frac{1}{2(x^2+1)} + C$$

2

5. $\int \ln x \, dx = x \ln x - x + C$. See textbook, §7.1, Example 2, p. 454.

So $\int_1^e \ln x \, dx = [x \ln x - x]_1^e = [e \ln e - e] - [\ln 1 - 1] = [e - e] - [-1] = 1$.

6. $\int x^2 e^x \, dx = x^2 e^x - 2x e^x + 2e^x + C = e^x (x^2 - 2x + 2) + C$.

See textbook, §7.1, Example 3, p. 455.

So $\int_0^1 x^2 e^x \, dx = [e^x (x^2 - 2x + 2)]_0^1 = [e(1 - 2 + 2)] - [e^0(0 - 0 + 2)] = [e(1)] - [1(2)] = e - 2$.

7. $\int e^x \sin x \, dx = \frac{e^x(\sin x - \cos x)}{2} + C$ (see textbook, §7.1, Example 4, p. 455-6)

So $\int_0^{\pi/2} e^x \sin x \, dx = \frac{e^{\pi/2}(\sin \frac{\pi}{2} - \cos \frac{\pi}{2})}{2} - \frac{e^0(\sin 0 - \cos 0)}{2}$
 $= \frac{e^{\pi/2}(1 - 0)}{2} - \frac{1(0 - 1)}{2} = \frac{e^{\pi/2}}{2} + \frac{1}{2}$
 $= \frac{1}{2}(1 + e^{\pi/2})$

8. $\int \sin^2 x \, dx = \left[\frac{x}{2} - \frac{1}{4} \sin(2x) \right] + C$. (See textbook, §7.2, Example 3, p. 461)


So $\int_0^{\pi/4} \sin^2 x \, dx = \left[\frac{1}{2} \left(\frac{\pi}{4} \right) - \frac{1}{4} \sin \frac{\pi}{2} \right] - \left[\frac{0}{2} - \frac{1}{4} \sin 0 \right] = \frac{\pi}{8} - \frac{1}{4}$

9. $\int \frac{1}{[x^2 + 2x + 2]^2} \, dx = \int \frac{1}{[(x+1)^2 + 1]^2} \, dx = \int \frac{\sec^2 \theta}{(\sec^2 \theta)^2} \, d\theta = \int \cos^2 \theta \, d\theta$

$x+1 = \tan \theta$
 $dx = \sec^2 \theta \, d\theta$
 $(x+1)^2 + 1 = \tan^2 \theta + 1 = \sec^2 \theta$

$= \frac{1}{2} \int (1 + \cos 2\theta) \, d\theta = \frac{1}{2} \left[\theta + \frac{\sin(2\theta)}{2} \right] + C$
 $= \frac{1}{2} \theta + \frac{1}{2} \frac{2 \cos \theta \sin \theta}{2} + C = \frac{1}{2} \theta + \frac{1}{2} \cos \theta \sin \theta + C$

$= \frac{1}{2} \arctan(x+1) + \frac{1}{2} \frac{x+1}{x^2 + 2x + 2} + C$



2


3

So

$$\int_{-1}^0 \frac{dx}{(x^2+2x+2)^2} = \left(\frac{1}{2} \tan^{-1}(x+1) + \frac{1}{2} \frac{x+1}{x^2+2x+2} \right) \Big|_{x=-1}^{x=0}$$

$$= \left[\frac{1}{2} \underbrace{\tan^{-1}(1)}_{=\frac{\pi}{4}} + \frac{1}{2} \frac{1}{2} \right] - \left[\frac{1}{2} \underbrace{\tan^{-1}(0)}_{=0} + \frac{1}{2} \cdot 0 \right] = \frac{\pi}{8} + \frac{1}{4} \checkmark$$

10. $\int \frac{-2}{(x-11)^3} dx = \int -2(x-11)^{-3} dx = \frac{-2(x-11)^{-2}}{-2} + C = \frac{1}{(x-11)^2} + C \checkmark$



$$\int_{10}^{12} \frac{-2}{(x-11)^3} dx = \left[\lim_{a \rightarrow 11^-} \int_{10}^a \frac{-2}{(x-11)^3} dx \right] + \left[\lim_{b \rightarrow 11^+} \int_b^{12} \frac{-2}{(x-11)^3} dx \right]$$

$$= \left[\lim_{a \rightarrow 11^-} \frac{1}{(x-11)^2} \Big|_{x=10}^{x=a} \right] + \left[\lim_{b \rightarrow 11^+} \frac{1}{(x-11)^2} \Big|_{x=b}^{x=12} \right]$$

$$= \left[\lim_{a \rightarrow 11^-} \left(\frac{1}{(a-11)^2} - 1 \right) \right] + \left[\lim_{b \rightarrow 11^+} \left(1 - \frac{1}{(b-11)^2} \right) \right]$$

$$= \left[\lim_{a \rightarrow 11^-} \frac{1}{(a-11)^2} \right] - 1 + 1 - \left[\lim_{b \rightarrow 11^+} \frac{1}{(b-11)^2} \right]$$

$$= \left[\lim_{a \rightarrow 11^-} \frac{1}{(a-11)^2} \right] - \left[\lim_{b \rightarrow 11^+} \frac{1}{(b-11)^2} \right]$$

$$\left[\begin{array}{ccc} \infty & - & \infty \end{array} \right] \checkmark$$


indeterminate form, DNE.

The region R for problems 11-14


Let R be the region in the first quadrant enclosed by

$$y = 7 - x \quad \text{and} \quad y = 0 \quad \text{and} \quad x = 3.$$

Note that R is the triangle with vertices: $(3, 0)$ and $(3, 4)$ and $(7, 0)$.



11.




With respect to $y \Rightarrow$ partition y axis & all expressed in terms of y
 Area of typical rectangle = (height)(base) = $[(7-y) - 3] \Delta y$
 Area of $R = \int_{y=0}^{y=4} [(7-y) - 3] dy$

11. Let's check answer since we know formula for area of a triangle.

$$\int_0^4 [(7-y) - 3] dy = \int_0^4 (4-y) dy = (4y - \frac{y^2}{2}) \Big|_0^4 = (16 - 8) - 0 = 8,$$

and area of triangle = $\frac{1}{2}(\text{base})(\text{height}) = \frac{1}{2}(7-3)(4-0) = \frac{1}{2} \cdot 4 \cdot 4 = 8.$

12.



wrt $y \Rightarrow$ partition y -axis & all expressed in terms of y
 \downarrow rotate abt y -axis
 Disk or washer
 \hookrightarrow has hole

Volume of typical washer = $\pi [(\text{big radius})^2 - (\text{little radius})^2] (\text{height})$

$$= \pi [(7-y)^2 - (3)^2] \Delta y$$

$$\text{Volume of solid} = \pi \int_{y=0}^{y=4} [(7-y)^2 - (3)^2] dy$$

Check: Basic calculus gives = $\frac{208\pi}{3}$


Volume cone = $\frac{1}{3} \pi r^2 h$
 Volume Cylinder = $\pi r^2 h$

\uparrow $h = \text{height}$
 $r = \text{radius of base}$

Volume = Volume of cone $\left\langle \begin{matrix} h=7 \\ r=7 \end{matrix} \right\rangle$ - Volume of cone $\left\langle \begin{matrix} h=3 \\ r=3 \end{matrix} \right\rangle$ - Volume cylinder $\left\langle \begin{matrix} h=4 \\ r=3 \end{matrix} \right\rangle$

$$= \pi \left[\frac{1}{3} 7^2 \cdot 7 - \frac{1}{3} 3^2 \cdot 3 - 3^2 \cdot 4 \right] = \frac{\pi}{3} [343 - 27 - 108] = \frac{208\pi}{3}.$$

B.



wrt $y \Rightarrow$ partition y -axis

\downarrow rotate abt a line parallel to y -axis

Disk or Washer
 \rightarrow no hole


Volume of typical disk = $\pi (\text{radius})^2 (\text{height})$
 $= \pi ((7-y) - 3)^2 (\Delta y)$

Volume of solid = $\pi \int_{y=0}^{y=4} ((7-y) - 3)^2 dy$

Check: Basic calculus gives $= \frac{64\pi}{3}$

Volume of cone $\left\{ \begin{array}{l} h = 4 \\ r = 7-3 = 4 \end{array} \right. = \frac{1}{3} \pi r^2 h = \frac{1}{3} \pi 4^2 \cdot 4 = \frac{64\pi}{3}$

14.



wrt $y \Rightarrow$ partition y -axis

\downarrow rotate abt x -axis
Shell Method.

Volume of typical shell = $2\pi (\text{avg radius}) (\text{height}) (\text{thickness})$
 $= 2\pi (y) [(7-y) - 3] (\Delta y)$

Volume of solid = $2\pi \int_{y=0}^{y=4} y [(7-y) - 3] dy$

Check: Basic calculus gives $= \frac{64\pi}{3}$

Volume of cone w/ height 4 & radius of base 4
 $= \frac{\pi}{3} \cdot 4 \cdot 4^2 = \frac{64\pi}{3}$

15. $r = \tan \theta \sec \theta \iff r = \frac{\sin \theta}{\cos \theta} \cdot \frac{1}{\cos \theta} \iff r = \frac{\sin \theta}{\cos^2 \theta}$

$\implies r \cos^2 \theta = \sin \theta \iff (r \cos \theta)^2 = r \sin \theta \iff \boxed{x^2 = y}$


Note that $(*) \implies$ can be reversed to \leftarrow

Since $\nexists r \cos^2 \theta = \sin \theta$ then $\cos^2 \theta \neq 0$ always.

Indeed $\nexists r \cos^2 \theta = \sin \theta$ and $\cos^2 \theta = 0$
 then $0 = \sin \theta$ so $\cos^2 \theta + \sin^2 \theta = 0$,
 which cannot be.

16. $r = 3 \cos \theta$, $\frac{1}{4}$ (period of $\cos \theta$) = $\frac{1}{4}(2\pi) = \frac{\pi}{2}$.

θ	$r = 3 \cos \theta$
$0 \xrightarrow{\textcircled{1}} \frac{\pi}{2}$	$3 \rightarrow 0$
$\frac{\pi}{2} \xrightarrow{\textcircled{2}} \pi$	$0 \rightarrow -3$
$\pi \xrightarrow{\textcircled{3}} \frac{3\pi}{2}$	$-3 \rightarrow 0$
$\frac{3\pi}{2} \xrightarrow{\textcircled{4}} 2\pi$	$0 \rightarrow 3$



$A = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta = \frac{1}{2} \int_0^{\pi} [3 \cos \theta]^2 d\theta$

17. $\frac{\sqrt{25n^3 + 4n^2 + n - 5}}{7n^{3/2} + 6n - 1}$
 \div num & dem. by $n^{3/2} = \sqrt{n^3}$

$\frac{\sqrt{25 + \frac{4}{n} + \frac{1}{n^2} - \frac{5}{n^3}}}{7 + \frac{6}{n^{1/2}} - \frac{1}{n^{3/2}}} \xrightarrow{n \rightarrow \infty} \frac{\sqrt{25}}{7} = \frac{5}{7}$

18 Want $\sum_{n=2}^{\infty} r^n = \frac{1}{12}$. We know that we must have $|r| < 1$ (geom. series) 7

Let $S_N = \sum_{n=2}^N r^n$, So

$$\begin{aligned} 1 \quad S_N &= r^2 + r^3 + r^4 + \dots + r^N \\ \text{subtract } r S_N &= r^3 + r^4 + \dots + r^N + r^{N+1} \end{aligned}$$

$$(1-r) S_N = r^2 - r^{N+1}$$

$$S_N \stackrel{r \neq 1}{=} \frac{r^2 - r^{N+1}}{1-r} \xrightarrow[N \rightarrow \infty]{\text{if } |r| < 1} \frac{r^2}{1-r}$$

So want $r \in \mathbb{R}$ so that $|r| < 1$ and $\frac{r^2}{1-r} = \frac{1}{12}$

$$\frac{r^2}{1-r} = \frac{1}{12} \Leftrightarrow 12r^2 = 1-r \Leftrightarrow 12r^2 + r - 1 = 0$$

$$\begin{aligned} \Leftrightarrow r &= \frac{-1 \pm \sqrt{1^2 - 4(12)(-1)}}{2(12)} = \frac{-1 \pm \sqrt{1+48}}{2 \cdot 12} = \frac{-1 \pm 7}{2 \cdot 12} \\ &= \begin{cases} \frac{-1-7}{2 \cdot 12} = \frac{-8}{2 \cdot 12} = -\frac{2 \cdot 4}{2 \cdot 4 \cdot 3} = -\frac{1}{3} \\ \frac{-1+7}{2 \cdot 12} = \frac{6}{2 \cdot 6 \cdot 2} = \frac{1}{4} \end{cases} \end{aligned}$$

19. Know (A) $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (p-series, $p=1$, $p \leq 1$) (or, also, harmonic series)
 (B) $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges (just apply the AST).

Now, look at the choices:


20 $\frac{1}{\sqrt{(n+2)(n+7)}}$ $\stackrel{n \text{ big}}{\sim} \frac{1}{\sqrt{n \cdot n}} = \frac{1}{n}$. Now look at the choices.

21. $\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)!}{[3(n+1)]!} \cdot \frac{(3n)!}{n!} = \frac{(n!)(n+1)}{(n!)} \cdot \frac{(3n)!}{[(3n)!](3n+1)(3n+2)(3n+3)}$
 $= \frac{n+1}{(3n+1)(3n+2)(3n+3)} \xrightarrow{n \rightarrow \infty} 0.$

22. $\sum \frac{(5x+15)^n}{4^n} = \sum \frac{[5(x+3)]^n}{4^n} = \sum \left(\frac{5}{4}\right)^n (x+3)^n \Rightarrow \text{center} =$

$\left[\left| \frac{(5x+15)^n}{4^n} \right| \right]^{1/n} = \left| \frac{5x+15}{4} \right| = \frac{5}{4} |x+3| \xrightarrow{n \rightarrow \infty} \frac{5}{4} |x+3|$

$\frac{5}{4} |x+3| < 1 \Leftrightarrow |x+3| < \frac{4}{5} \Rightarrow \text{rad. of conv. is } \frac{4}{5}$



Check endpoints...

$x = -\frac{11}{5} : \sum \frac{(5x+15)^n}{4^n} = \sum \frac{4^n}{4^n} = \sum_{n=1}^{\infty} 1 = \text{divg (to } \infty)$

$x = -\frac{19}{5} : \sum \frac{(5x+15)^n}{4^n} = \sum \frac{(-4)^n}{4^n} = \sum_{n=1}^{\infty} (-1)^n = \text{divg (osc.)}$

23. $\ln(10-x) = \ln(1+(9-x)) \stackrel{(*)}{=} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(9-x)^n}{n} = \sum$

$= \sum_{n=1}^{\infty} (-1)^{n+1} \frac{((-1)(x-9))^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} (-1)^n (x-9)^n}{n}$

$= \sum_{n=1}^{\infty} (-1)^{2n+1} \frac{(x-9)^n}{n} = \sum_{n=1}^{\infty} -\frac{1}{n} (x-9)^n$

valid $(*) \quad -1 < 9-x \leq 1 \Leftrightarrow -1 \leq x-9 < 1 \Leftrightarrow 8 \leq x < 10.$

24. The computations below show that the 3rd order Taylor polynomial, about the center $x_0 = 1$, for the function $f(x) = x^5 - x^2 + 5$ is $p_3(x) = 5 + 3(x - 1) + 9(x - 1)^2 + 10(x - 1)^3$.

we were given $x_0 = 1$			
n	$f^{(n)}(x)$	$f^{(n)}(x_0)$	$\frac{f^{(n)}(x_0)}{n!}$
0	$x^5 - x^2 + 5$	5	$\frac{5}{0!} = \frac{5}{1} = 5$
1	$5x^4 - 2x$	$5 - 2 = 3$	$\frac{3}{1!} = \frac{3}{1} = 3$
2	$5 \cdot 4x^3 - 2$	$20 - 2 = 18$	$\frac{18}{2!} = \frac{18}{2} = 9$
3	$5 \cdot 4 \cdot 3$	$(5)(4)(3)$	$\frac{(5)(4)(3)}{3!} = \frac{(5)(4)(3)}{(3)(2)} = \frac{(5)(4)}{2} = 10$

25. Consider the function $f(x) = e^{-x}$ as well as the interval $(7, 9)$.

The 5th order Taylor polynomial of $y = f(x)$ about the center $x_0 = 0$ is

$$P_5(x) = \sum_{n=0}^5 \frac{(-x)^n}{n!} = 1 - x + \frac{x^2}{2!} - \frac{x^3}{3!} + \frac{x^4}{4!} - \frac{x^5}{5!}.$$

The 5th order Remainder term $R_5(x)$ is defined by $R_5(x) = f(x) - P_5(x)$ and so $f(x) \approx P_5(x)$ where the approximation is within an error of $|R_5(x)|$. Using Taylor's (BIG) Theorem, find a good upper bound for $|R_5(x)|$ that is valid for each $x \in (7, 9)$.

By Taylor's Remainder Theorem, for each $x \in (7, 9)$, there exists c between x and 0 so that

$$R_5(x) = \frac{f^{(6)}(c) (x - 0)^6}{6!}.$$

Note that if $x \in (7, 9)$ and c is between x and 0, then $c \in (0, 9)$. So for each $x \in (7, 9)$,

$$|R_5(x)| = \left| \frac{f^{(6)}(c) (x - 0)^6}{6!} \right| = \frac{|f^{(6)}(c)| |x|^6}{6!} = \frac{e^{-c} |x|^6}{6!} \leq \frac{e^{-c} 9^6}{6!} \leq \frac{e^{-0} 9^6}{6!} = \frac{9^6}{6!}.$$

If you prefer, you can also think of the above line as:

$$|R_5(x)| = \left| \frac{f^{(6)}(c) (x - 0)^6}{6!} \right| = \frac{|f^{(6)}(c)| |x|^6}{6!} = \frac{e^{-c} |x|^6}{6!} = \frac{|x|^6}{e^c 6!} \leq \frac{9^6}{e^c 6!} \leq \frac{9^6}{e^0 6!} = \frac{9^6}{6!}.$$