

HAND IN PART

Prof. Girardi

Math 142

Spring 2013

03.26.2013

Exam 2

MARK BOX		
PROBLEM	POINTS	
0	30	
1	10	
2	10	
3	10	
4	10	
5–10	30	
%	100	

NAME: _____

PIN: _____

INSTRUCTIONS

- (1) The MARK BOX above indicates the problems along with their points.
Check that your copy of the exam has all of the problems.
- (2) You may **not** use an electronic device, a calculator, books, personal notes.
- (3) On Problem 0, fill in the blanks. As you were warned, if you do not make at least half of the points on Problem 0, then your score for the entire exam will be whatever you made on Problem 0.
- (4) For Problems 1–4, to receive credit you must:
 - (a) **work in a logical fashion, show all your work, indicate your reasoning;**
no credit will be given for an answer that *just appears*;
such explanations help with partial credit
 - (b) if a line/box is provided, then:
 - show you work BELOW the line/box
 - put your answer on/in the line/box
 - (c) if no such line/box is provided, then box your answer.
- (5) Problems 5–10 are multiple choice.
 - First, indicate (to yourself) your answers directly on the STATEMENT OF MULTIPLE CHOICE PROBLEMS part.
 - Once finished with the multiple choice problems, go back to the HAND IN PART and indicate your answers on the table provided.
 - Hand in the HAND IN PART. You can take the STATEMENT OF MULTIPLE CHOICE PROBLEMS part home with you (so you can check your answers once the solutions are posted).
- (6) During this exam, do not leave your seat unless you have permission. If you have a question, raise your hand. When you finish: put your pencil down and raise your hand.
- (7) This exam covers (from *Calculus* by Stewart, 6th ed., ET):
§ 11.2 – 11.8 .

0. Fill-in-the blanks. All series \sum are understood to be $\sum_{n=1}^{\infty}$, unless otherwise indicated.

0a. n^{th} -term test for an arbitrary series $\sum a_n$.

If $\lim_{n \rightarrow \infty} a_n \neq 0$ or $\lim_{n \rightarrow \infty} a_n$ does not exist, then $\sum a_n$ _____ .

0b. **Geometric Series** where $-\infty < r < \infty$. The series $\sum r^n$

- converges if and only if $|r|$ _____
- diverges if and only if $|r|$ _____

0c. p -series where $0 < p < \infty$. The series $\sum \frac{1}{n^p}$

- converges if and only if p _____
- diverges if and only if p _____

0d. **Integral Test** for a positive-termed series $\sum a_n$ where $a_n \geq 0$.

Let $f: [1, \infty) \rightarrow \mathbb{R}$ be so that

- $a_n = f(\text{_____})$ for each $n \in \mathbb{N}$
- f is a _____ function
- f is a _____ function
- f is a _____ function .

Then $\sum a_n$ converges if and only if _____ converges.

0e. **Comparison Test** for a positive-termed series $\sum a_n$ where $a_n \geq 0$. (Fill in the blanks with a_n and/or b_n .)

- If $0 \leq a_n \leq b_n$ for all $n \in \mathbb{N}$ and \sum _____ converge, then \sum _____ converge.
- If $0 \leq b_n \leq a_n$ for all $n \in \mathbb{N}$ and \sum _____ diverge, then \sum _____ diverge.

0f. **Limit Comparison Test** for a positive-termed series $\sum a_n$ where $a_n \geq 0$.

Let $b_n > 0$ and $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L$.

If _____ $< L <$ _____, then $\sum a_n$ converges if and only if _____ .

0g. **Ratio and Root Tests** for arbitrary-termed series $\sum a_n$ with $-\infty < a_n < \infty$.

Let $\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$ or $\rho = \lim_{n \rightarrow \infty} |a_n|^{\frac{1}{n}}$.

- If ρ _____ then $\sum a_n$ converges absolutely.
- If ρ _____ then $\sum a_n$ diverges.
- If ρ _____ then the test is inconclusive.

0h. **Alternating Series Test** for an alternating series $\sum (-1)^n a_n$ where $a_n > 0$ for each $n \in \mathbb{N}$.

If

- a_n _____ a_{n+1} for each $n \in \mathbb{N}$
- $\lim_{n \rightarrow \infty} a_n =$ _____

then $\sum (-1)^n a_n$ _____

0i. By definition, for an arbitrary series $\sum a_n$, (fill in the blanks with converges or diverges).

- $\sum a_n$ is absolutely convergent if and only if $\sum |a_n|$ _____
- $\sum a_n$ is conditionally convergent if and only if $\sum a_n$ _____ and $\sum |a_n|$ _____
- $\sum a_n$ is divergent if and only if $\sum a_n$ _____

2. Check the correct box and then indicate your reasoning below. Specifically specify what test(s) you are using. A correctly checked box without appropriate explanation will receive no points.

$$\sum_{n=1}^{\infty} (-1)^n \frac{n^3}{3n^3 + 2n^2}$$

absolutely convergent

conditionally convergent

divergent

3. Let

$$a_n = \frac{n!}{(2n-1)!}$$

3a. Find an expression for $\frac{a_{n+1}}{a_n}$ that does NOT have a factorial sign (that is a ! sign) in it.

$\frac{a_{n+1}}{a_n} =$

3b. Check the correct box and then indicate your reasoning below. Specifically specify what test(s) you are using. A correctly checked box without appropriate explanation will receive no points.

$$\sum_{n=1}^{\infty} (-1)^n \frac{n!}{(2n-1)!}$$

absolutely convergent

conditionally convergent

divergent

4. Consider the formal power series

$$\sum_{n=2}^{\infty} \frac{x^n}{(\ln n)^n}.$$

Hint 1: $\frac{x^n}{(\ln n)^n} = \left[\frac{x}{\ln n}\right]^n$ so would you rather use the root test or the ratio test?

Hint 2: $\ln(a^r) = r \ln(a)$ but $(\ln(a))^r \neq r \ln(a)$

The center is $x_0 =$ _____ and the radius of convergence is $R =$ _____. As we did in class, make a number line indicating where the power series is: absolutely convergent, conditionally convergent, and divergent. Indicate your reasoning and specifically specify what test(s) you are using. Don't forget to check the endpoints, if there are any.



TABLE FOR YOUR ANSWERS TO MULTIPLE CHOICE PROBLEMS 5 – 10

Instructions.

- Indicate (by circling) your solution to each problem.
- You may choose up to 3 answers for each problem. The scoring is as follows. For a problem with precisely one answer marked and the answer is correct, 5 points. For a problem with precisely two answers marked, one of which is correct, 3 points. For a problem with precisely three answers marked, one of which is correct, 1 point. All other cases, 0 points.

Your Solutions						
PROBLEM						points
5	5a	5b	5c	5d	5e	
6	6a	6b	6c	6d	6e	
7	7a	7b	7c	7d	7e	
8	8a	8b	8c	8d	8e	
9	9a	9b	9c	9d	9e	
10	10a	10b	10c	10d	10e	

STATEMENT OF MULTIPLE CHOICE PROBLEMS

Instructions.

- Indicate (by circling) your solution to each problem on the Hand In Part of the exam.
- You may choose up to 3 answers for each problem. The scoring is as follows. For a problem with precisely one answer marked and the answer is correct, 5 points. For a problem with precisely two answers marked, one of which is correct, 3 points. For a problem with precisely three answers marked, one of which is correct, 1 point. All other cases, 0 points.

5. Evaluate

$$\sum_{n=17}^{\infty} 5 \frac{(-2)^n}{3^{2n+1}}$$

- a. $\left(\frac{7}{11}\right)\left(\frac{5}{3}\right)\left(\frac{-2}{9}\right)^{18}$ b. $\left(\frac{9}{11}\right)\left(\frac{-2}{9}\right)^{17}$ c. $\left(\frac{7}{11}\right)\left(\frac{5}{3}\right)\left(\frac{-2}{9}\right)^{17}$ d. $\left(\frac{9}{11}\right)\left(\frac{5}{3}\right)\left(\frac{-2}{9}\right)^{17}$ e. None of these

6. Evaluate

$$\sum_{n=1}^{\infty} \frac{4}{(4n-3)(4n+1)}$$

Hint: Telescoping Series, use PFD.

- a. 2 b. $\frac{1}{2}$ c. 1 d. 4 e. None of these

7. Consider the following two series.

Series A is $\sum_{n=1}^{\infty} \frac{1}{n}$

Series B is $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$.

- a. both series converge absolutely b. both series diverge
 c. series A converges conditionally and series B diverges
 d. series A diverges and series B converges conditionally e. none of these
8. Consider the formal series $\sum_{n=1}^{\infty} a_n$ where

$$a_n = (-1)^n \frac{(n+1)!}{(2n)!}$$

and let

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|.$$

- a. $\sum_{n=1}^{\infty} a_n$ converges absolutely because $\rho = \frac{1}{2}$. b. $\sum_{n=1}^{\infty} a_n$ converges absolutely because $\rho = 0$.
 c. $\rho = 1$ so the Ratio Test fails for $\sum_{n=1}^{\infty} a_n$ d. $\sum_{n=1}^{\infty} a_n$ diverges e. none of these
9. What is the LARGEST interval for which the formal power series

$$\sum_{n=1}^{\infty} \frac{(2x+6)^n}{4^n}$$

is absolutely convergent?

- a. (1, 5) b. (-4, -2) c. (-5, -1) d. [-5, -1] e. none of these
10. Suppose that the radius of convergence of a power series $\sum_{n=0}^{\infty} c_n x^n$ is 16. What is the radius of convergence of the power series $\sum_{n=0}^{\infty} c_n x^{2n}$?
- a. 256 b. 4 c. 1 d. 16 e. none of these