**0.** Fill-in-the boxes. All series  $\sum$  are understood to be  $\sum_{n=1}^{\infty}$  , unless otherwise indicated.

Sequences

0.1. Practice taking basic limits of sequences. (Important, e.g., for Ratio and Root Tests.) Can you do similar problems?

$$\bullet \lim_{n \to \infty} \frac{5n^{17} + 6n^2 + 1}{7n^{18} + 9n^3 + 5} =$$

$$\bullet \lim_{n \to \infty} \sqrt{\frac{36n^{17} - 6n^2 - 1}{4n^{17} + 9n^3 + 5}} = \boxed{}$$

$$\bullet \lim_{n \to \infty} \frac{-5n^{18} + 6n^2 + 1}{7n^{17} + 9n^3 + 5} = \boxed{}$$

$$\bullet \lim_{n \to \infty} n^{\frac{1}{n}} =$$

**0.2.** Geometric Sequence. Fill in the boxes with with the proper range of  $r \in \mathbb{R}$ . (Needed for Geometric Series!)

•  $\lim_{n\to\infty} r^n = 0$  if and only if r satisfies

•  $\lim_{n\to\infty} r^n = 1$  if and only if r satisfies

• the sequence  $\{r^n\}_{n=1}^{\infty}$  diverges to  $\infty$  if and only if r satisfies

• the sequence  $\{r^n\}_{n=1}^{\infty}$  diverges but does not diverge to  $\infty$  if and only if r satisfies

**0.3. Commonly Occurring Limits of Sequences**. Here,  $c \in \mathbb{R}$  is a constant.  $\langle$  Thomas Book §10.1, Theorem 5

 $(1) \lim_{n \to \infty} \frac{\ln n}{n} = \boxed{}$ 

$$(2) \lim_{n \to \infty} \sqrt[n]{n} =$$

 $(3) \lim_{n \to \infty} c^{1/n} = \boxed{ (c > 0)}$ 

$$(4) \lim_{n \to \infty} c^n = \boxed{ (|c| < 1)}$$

 $(5) \lim_{n \to \infty} \left( 1 + \frac{c}{n} \right)^n = \boxed{ (c \in \mathbb{R})}$ 

(6)  $\lim_{n \to \infty} \frac{x^n}{n!} = \boxed{ (c \in \mathbb{R})}$ 

Series

For a formal series $\sum_{n=1}^{\infty} a_n$ , where each $a_n \in \mathbb{R}$ , consider the corresponding sequence $\{s_n\}_{n=1}^{\infty}$ of									
partial sums, so $s_n = \sum_{k=1}^n a_k$ . Then the formal series $\sum a_n$ :									
• converges if and only if									
$ullet$ converges to $L \in \mathbb{R}$ if and only if									
• diverges if and only if									
Now assume, furthermore, that $a_n \geq 0$ for each $n$ . Then the sequence $\{s_n\}_{n=1}^{\infty}$ of partial sums									
$\underbrace{ ext{either}}$									
• <u>is</u> bounded above (by some finite number), in which case the <u>series</u> $\sum a_n$									
$ \circ_{\!$									
• is not bounded above (by some finite number), in which case the series $\sum a_n$									
Fix $r \in \mathbb{R}$ with $r \neq 1$ . For $N \geq 50$ , let $s_N = \sum_{n=50}^{N} r^n$ . (Note the sum starts at 50.) For each $N \geq 50$ ,									
the partial sums $s_N$ can be written as: (your answer should NOT contain a "" nor a " $\sum$ " sign)									
$s_N = $ .									
<b>Geometric</b> Series. Fill in the boxes with the proper range of $r \in \mathbb{R}$ .									
• The series $\sum r^n$ converges if and only if $r$ satisfies $\square$ .									

**0.8.** *p*-series. Fill in the boxes with the proper range of  $p \in \mathbb{R}$ .

**0.7.** State the  $n^{\text{th}}$ -term test for an arbitrary series  $\sum a_n$ .

• The series  $\sum \frac{1}{n^p}$  converges if and only if

Tests for Positive-Termed Series
(so for $\sum a_n$ where $a_n \geq 0$ )

0.9.	State the	Integral	Test	with	Remainder	Estimate for	rai	positive-termed	series	$\sum a_n$ .
	~ cccc c crrc						_ ~	0 0 0 1 0 1 0 0 0 1 1 1 1 0 0 0	NOT TON	/ / 0011.

Let  $f: [1, \infty) \to \mathbb{R}$  be so that

- (1)  $a_n = f(n)$  for each  $n \in \mathbb{N}$
- (2) f is a function
- (3) f is a function
- (4) f is a function.

Then

- $\sum a_n$  converges if and only if converges.
- and if  $\sum a_n$  converges, then

$$0 \le \left(\sum_{k=1}^{\infty} a_k\right) - \left(\sum_{k=1}^{N} a_k\right) \le$$

- **0.10.** State the **Direct Comparison Test** for a <u>positive</u>-termed series  $\sum a_n$ .
  - If when  $n \ge 17$  and , then  $\sum a_n$  converges.
  - If when  $n \ge 17$  and , then  $\sum a_n$  diverges.

Hint: sing the song to yourself.

## **0.11.** State the **Limit Comparison Test** for a <u>positive</u>-termed series $\sum a_n$ .

Let  $b_n > 0$  and  $L = \lim_{n \to \infty} \frac{a_n}{b_n}$ .

- If  $0 < L < \infty$ , then
- If L=0, then
- If  $L = \infty$ , then

Goal: cleverly pick positive  $b_n$ 's so that you know what  $\sum b_n$  does (converges or diverges) and the sequence  $\left\{\frac{a_n}{b_n}\right\}_n$  converges.

## **0.12. Helpful Intuition** Fill in the 3 boxes using: $e^x$ , $\ln x$ , $x^q$ . Use each once, and only once.

Consider a positive power q > 0. There is (some big number)  $N_q > 0$  so that if  $x \ge N_q$  then

$$\leq$$
  $\leq$   $.$ 

## Tests for Arbitrary-Termed Series (so for $\sum a_n$ where $-\infty < a_n < \infty$ )

**0.13.** By definition, for an arbitrary series  $\sum a_n$ , (fill in these 3 boxes with convergent or divergent).

- $\sum a_n$  is absolutely convergent if and only if  $\sum |a_n|$  is
- $\sum a_n$  is <u>conditionally convergent</u> if and only if

 $\sum a_n$  is and  $\sum |a_n|$  is

•  $\sum a_n$  is divergent if and only if  $\sum a_n$  is divergent.

**0.14.** State the Ratio and Root Tests for arbitrary-termed series  $\sum a_n$  with  $-\infty < a_n < \infty$ . Let

$$\rho = \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right| \quad \text{or} \quad \rho = \lim_{n \to \infty} |a_n|^{\frac{1}{n}}.$$

- If  $\sum a_n$  converges absolutely.
- If then  $\sum a_n$  diverges.
- If then the test is inconclusive.

0.15. State the Alternating Series Test (AST) & Alternating Series Estimation Theorem.

Let

- (1)  $u_n \ge 0$  for each  $n \in \mathbb{N}$
- $(2) \lim_{n\to\infty} u_n = \boxed{}$
- (3)  $u_n$   $u_{n+1}$  for each  $n \in \mathbb{N}$ .

Then

- •
- and we can estimate (i.e., approximate) the infinite sum  $\sum_{n=1}^{\infty} (-1)^n u_n$  by the finite sum  $\sum_{k=1}^{N} (-1)^k u_k$  and the error (i.e. remainder) satisfies

$$\left| \sum_{k=1}^{\infty} (-1)^k u_k - \sum_{k=1}^{N} (-1)^k u_k \right| \le \boxed{}.$$