

7.8 Improper Integrals

1. (a) Since $\int_1^\infty x^4 e^{-x^4} dx$ has an infinite interval of integration, it is an improper integral of Type I.

(b) Since $y = \sec x$ has an infinite discontinuity at $x = \frac{\pi}{2}$, $\int_0^{\pi/2} \sec x dx$ is a Type II improper integral.

(c) Since $y = \frac{x}{(x-2)(x-3)}$ has an infinite discontinuity at $x = 2$, $\int_0^2 \frac{x}{x^2 - 5x + 6} dx$ is a Type II improper integral.

(d) Since $\int_{-\infty}^0 \frac{1}{x^2 + 5} dx$ has an infinite interval of integration, it is an improper integral of Type I.

2. (a) Since $y = \frac{1}{2x-1}$ is defined and continuous on $[1, 2]$, $\int_1^2 \frac{1}{2x-1} dx$ is proper.

(b) Since $y = \frac{1}{2x-1}$ has an infinite discontinuity at $x = \frac{1}{2}$, $\int_0^{1/2} \frac{1}{2x-1} dx$ is a Type II improper integral.

(c) Since $\int_{-\infty}^{\infty} \frac{\sin x}{1+x^2} dx$ has an infinite interval of integration, it is an improper integral of Type I.

(d) Since $y = \ln(x-1)$ has an infinite discontinuity at $x = 1$, $\int_1^2 \ln(x-1) dx$ is a Type II improper integral.

3. The area under the graph of $y = 1/x^3 = x^{-3}$ between $x = 1$ and $x = t$ is



$A(t) = \int_1^t x^{-3} dx = \left[-\frac{1}{2}x^{-2} \right]_1^t = -\frac{1}{2}t^{-2} - \left(-\frac{1}{2} \right) = \frac{1}{2} - 1/(2t^2)$. So the area for $1 \leq x \leq 10$ is

$A(10) = 0.5 - 0.005 = 0.495$, the area for $1 \leq x \leq 100$ is $A(100) = 0.5 - 0.00005 = 0.49995$, and the area for

$1 \leq x \leq 1000$ is $A(1000) = 0.5 - 0.0000005 = 0.4999995$. The total area under the curve for $x \geq 1$ is

$$\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left[\frac{1}{2} - 1/(2t^2) \right] = \frac{1}{2}.$$

4. (a)



(b) The area under the graph of f from $x = 1$ to $x = t$ is

$$\begin{aligned} F(t) &= \int_1^t f(x) dx = \int_1^t x^{-1.1} dx = \left[-\frac{1}{0.1}x^{-0.1} \right]_1^t \\ &= -10(t^{-0.1} - 1) = 10(1 - t^{-0.1}) \end{aligned}$$

and the area under the graph of g is

$$G(t) = \int_1^t g(x) dx = \int_1^t x^{-0.9} dx = \left[\frac{1}{0.1}x^{0.1} \right]_1^t = 10(t^{0.1} - 1).$$

t	$F(t)$	$G(t)$
10	2.06	2.59
100	3.69	5.85
10^4	6.02	15.12
10^6	7.49	29.81
10^{10}	9	90
10^{20}	9.9	990

(c) The total area under the graph of f is $\lim_{t \rightarrow \infty} F(t) = \lim_{t \rightarrow \infty} 10(1 - t^{-0.1}) = 10$.

The total area under the graph of g does not exist, since $\lim_{t \rightarrow \infty} G(t) = \lim_{t \rightarrow \infty} 10(t^{0.1} - 1) = \infty$.

5. $I = \int_1^\infty \frac{1}{(3x+1)^2} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{(3x+1)^2} dx$. Now

$$\int \frac{1}{(3x+1)^2} dx = \frac{1}{3} \int \frac{1}{u^2} du \quad [u = 3x+1, du = 3dx] = -\frac{1}{3u} + C = -\frac{1}{3(3x+1)} + C,$$

$$\text{so } I = \lim_{t \rightarrow \infty} \left[-\frac{1}{3(3x+1)} \right]_1^t = \lim_{t \rightarrow \infty} \left[-\frac{1}{3(3t+1)} + \frac{1}{12} \right] = 0 + \frac{1}{12} = \frac{1}{12}. \quad \text{Convergent}$$

6. $\int_{-\infty}^0 \frac{1}{2x-5} dx = \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{2x-5} dx = \lim_{t \rightarrow -\infty} [\frac{1}{2} \ln |2x-5|]_t^0 = \lim_{t \rightarrow -\infty} [\frac{1}{2} \ln 5 - \frac{1}{2} \ln |2t-5|] = -\infty.$

Divergent

7. $\int_{-\infty}^{-1} \frac{1}{\sqrt{2-w}} dw = \lim_{t \rightarrow -\infty} \int_t^{-1} \frac{1}{\sqrt{2-w}} dw = \lim_{t \rightarrow -\infty} [-2\sqrt{2-w}]_t^{-1} \quad [u = 2-w, du = -dw]$
 $= \lim_{t \rightarrow -\infty} [-2\sqrt{3} + 2\sqrt{2-t}] = \infty. \quad \text{Divergent}$

8. $\int_0^\infty \frac{x}{(x^2+2)^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{x}{(x^2+2)^2} dx = \lim_{t \rightarrow \infty} \frac{1}{2} \left[\frac{-1}{x^2+2} \right]_0^t = \frac{1}{2} \lim_{t \rightarrow \infty} \left(\frac{-1}{t^2+2} + \frac{1}{2} \right)$
 $= \frac{1}{2} \left(0 + \frac{1}{2} \right) = \frac{1}{4}. \quad \text{Convergent}$

9. $\int_4^\infty e^{-y/2} dy = \lim_{t \rightarrow \infty} \int_4^t e^{-y/2} dy = \lim_{t \rightarrow \infty} [-2e^{-y/2}]_4^t = \lim_{t \rightarrow \infty} (-2e^{-t/2} + 2e^{-2}) = 0 + 2e^{-2} = 2e^{-2}.$

Convergent

10. $\int_{-\infty}^{-1} e^{-2t} dt = \lim_{x \rightarrow -\infty} \int_x^{-1} e^{-2t} dt = \lim_{x \rightarrow -\infty} [-\frac{1}{2}e^{-2t}]_x^{-1} = \lim_{x \rightarrow -\infty} [-\frac{1}{2}e^2 + \frac{1}{2}e^{-2x}] = \infty. \quad \text{Divergent}$

11. $\int_{-\infty}^\infty \frac{x dx}{1+x^2} = \int_{-\infty}^0 \frac{x dx}{1+x^2} + \int_0^\infty \frac{x dx}{1+x^2} \text{ and}$

$$\int_{-\infty}^0 \frac{x dx}{1+x^2} = \lim_{t \rightarrow -\infty} [\frac{1}{2} \ln(1+x^2)]_t^0 = \lim_{t \rightarrow -\infty} [0 - \frac{1}{2} \ln(1+t^2)] = -\infty. \quad \text{Divergent}$$

12. $I = \int_{-\infty}^\infty (2-v^4) dv = I_1 + I_2 = \int_{-\infty}^0 (2-v^4) dv + \int_0^\infty (2-v^4) dv$, but

$I_1 = \lim_{t \rightarrow -\infty} [2v - \frac{1}{5}v^5]_t^0 = \lim_{t \rightarrow -\infty} (-2t + \frac{1}{5}t^5) = -\infty$. Since I_1 is divergent, I is divergent, and there is no need to evaluate I_2 . Divergent

13. $\int_{-\infty}^\infty xe^{-x^2} dx = \int_{-\infty}^0 xe^{-x^2} dx + \int_0^\infty xe^{-x^2} dx$.

$$\int_{-\infty}^0 xe^{-x^2} dx = \lim_{t \rightarrow -\infty} (-\frac{1}{2}) \left[e^{-x^2} \right]_t^0 = \lim_{t \rightarrow -\infty} (-\frac{1}{2}) \left(1 - e^{-t^2} \right) = -\frac{1}{2} \cdot 1 = -\frac{1}{2}, \text{ and}$$

$$\int_0^\infty xe^{-x^2} dx = \lim_{t \rightarrow \infty} (-\frac{1}{2}) \left[e^{-x^2} \right]_0^t = \lim_{t \rightarrow \infty} (-\frac{1}{2}) \left(e^{-t^2} - 1 \right) = -\frac{1}{2} \cdot (-1) = \frac{1}{2}.$$

Therefore, $\int_{-\infty}^\infty xe^{-x^2} dx = -\frac{1}{2} + \frac{1}{2} = 0$. Convergent

$$14. \int_1^\infty \frac{e^{-\sqrt{x}}}{\sqrt{x}} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{e^{-\sqrt{x}}}{\sqrt{x}} dx = \lim_{t \rightarrow \infty} \int_1^{\sqrt{t}} e^{-u} (2 du) \quad \left[\begin{array}{l} u = \sqrt{x}, \\ du = dx/(2\sqrt{x}) \end{array} \right]$$

$$= 2 \lim_{t \rightarrow \infty} \left[-e^{-u} \right]_1^{\sqrt{t}} = 2 \lim_{t \rightarrow \infty} (-e^{-\sqrt{t}} + e^{-1}) = 2(0 + e^{-1}) = 2e^{-1}. \quad \text{Convergent}$$

15. $\int_{2\pi}^\infty \sin \theta d\theta = \lim_{t \rightarrow \infty} \int_{2\pi}^t \sin \theta d\theta = \lim_{t \rightarrow \infty} [-\cos \theta]_{2\pi}^t = \lim_{t \rightarrow \infty} (-\cos t + 1)$. This limit does not exist, so the integral is divergent. Divergent

16. $I = \int_{-\infty}^\infty \cos \pi t dt = I_1 + I_2 = \int_{-\infty}^0 \cos \pi t dt + \int_0^\infty \cos \pi t dt$, but $I_1 = \lim_{s \rightarrow -\infty} \left[\frac{1}{\pi} \sin \pi t \right]_s^0 = \lim_{s \rightarrow -\infty} \left(-\frac{1}{\pi} \sin \pi t \right)$ and this limit does not exist. Since I_1 is divergent, I is divergent, and there is no need to evaluate I_2 . Divergent

$$17. \int_1^\infty \frac{x+1}{x^2+2x} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{\frac{1}{2}(2x+2)}{x^2+2x} dx = \frac{1}{2} \lim_{t \rightarrow \infty} \left[\ln(x^2+2x) \right]_1^t = \frac{1}{2} \lim_{t \rightarrow \infty} [\ln(t^2+2t) - \ln 3] = \infty.$$

Divergent

$$18. \int_0^\infty \frac{dz}{z^2+3z+2} = \lim_{t \rightarrow \infty} \int_0^t \left[\frac{1}{z+1} - \frac{1}{z+2} \right] dz = \lim_{t \rightarrow \infty} \left[\ln \left(\frac{z+1}{z+2} \right) \right]_0^t$$

$$= \lim_{t \rightarrow \infty} \left[\ln \left(\frac{t+1}{t+2} \right) - \ln \left(\frac{1}{2} \right) \right] = \ln 1 + \ln 2 = \ln 2. \quad \text{Convergent}$$

$$19. \int_0^\infty se^{-5s} ds = \lim_{t \rightarrow \infty} \int_0^t se^{-5s} ds = \lim_{t \rightarrow \infty} \left[-\frac{1}{5}se^{-5s} - \frac{1}{25}e^{-5s} \right] \quad \left[\begin{array}{l} \text{by integration by} \\ \text{parts with } u = s \end{array} \right]$$

$$= \lim_{t \rightarrow \infty} \left(-\frac{1}{5}te^{-5t} - \frac{1}{25}e^{-5t} + \frac{1}{25} \right) = 0 - 0 + \frac{1}{25} \quad [\text{by l'Hospital's Rule}]$$

$$= \frac{1}{25}. \quad \text{Convergent}$$

$$20. \int_{-\infty}^6 re^{r/3} dr = \lim_{t \rightarrow -\infty} \int_t^6 re^{r/3} dr = \lim_{t \rightarrow -\infty} \left[3re^{r/3} - 9e^{r/3} \right]_t^6 \quad \left[\begin{array}{l} \text{by integration by} \\ \text{parts with } u = r \end{array} \right]$$

$$= \lim_{t \rightarrow -\infty} (18e^2 - 9e^2 - 3te^{t/3} + 9e^{t/3}) = 9e^2 - 0 + 0 \quad [\text{by l'Hospital's Rule}]$$

$$= 9e^2. \quad \text{Convergent}$$

$$21. \int_1^\infty \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[\frac{(\ln x)^2}{2} \right]_1^t \quad \left[\begin{array}{l} \text{by substitution with} \\ u = \ln x, du = dx/x \end{array} \right] = \lim_{t \rightarrow \infty} \frac{(\ln t)^2}{2} = \infty. \quad \text{Divergent}$$

22. $I = \int_{-\infty}^\infty x^3 e^{-x^4} dx = I_1 + I_2 = \int_{-\infty}^0 x^3 e^{-x^4} dx + \int_0^\infty x^3 e^{-x^4} dx$. Now

$$I_2 = \lim_{t \rightarrow \infty} \int_0^t x^3 e^{-x^4} dx = \lim_{t \rightarrow \infty} \int_0^{t^4} e^{-u} (\frac{1}{4} du) \quad \left[\begin{array}{l} u = x^4, \\ du = 4x^3 dx \end{array} \right]$$

$$= \frac{1}{4} \lim_{t \rightarrow \infty} \left[-e^{-u} \right]_0^{t^4} = \frac{1}{4} \lim_{t \rightarrow \infty} (-e^{-t^4} + 1) = \frac{1}{4}(0 + 1) = \frac{1}{4}.$$

Since $f(x) = x^3 e^{-x^4}$ is an odd function, $I_1 = -\frac{1}{4}$, and hence, $I = 0$. Convergent

23. $\int_{-\infty}^{\infty} \frac{x^2}{9+x^6} dx = \int_{-\infty}^0 \frac{x^2}{9+x^6} dx + \int_0^{\infty} \frac{x^2}{9+x^6} dx = 2 \int_0^{\infty} \frac{x^2}{9+x^6} dx$ [since the integrand is even].

$$\text{Now } \int \frac{x^2 dx}{9+x^6} \begin{bmatrix} u = x^3 \\ du = 3x^2 dx \end{bmatrix} = \int \frac{\frac{1}{3} du}{9+u^2} \begin{bmatrix} u = 3v \\ du = 3 dv \end{bmatrix} = \int \frac{\frac{1}{3}(3 dv)}{9+9v^2} = \frac{1}{9} \int \frac{dv}{1+v^2}$$

$$= \frac{1}{9} \tan^{-1} v + C = \frac{1}{9} \tan^{-1}\left(\frac{u}{3}\right) + C = \frac{1}{9} \tan^{-1}\left(\frac{x^3}{3}\right) + C,$$

$$\text{so } 2 \int_0^{\infty} \frac{x^2}{9+x^6} dx = 2 \lim_{t \rightarrow \infty} \int_0^t \frac{x^2}{9+x^6} dx = 2 \lim_{t \rightarrow \infty} \left[\frac{1}{9} \tan^{-1}\left(\frac{x^3}{3}\right) \right]_0^t = 2 \lim_{t \rightarrow \infty} \frac{1}{9} \tan^{-1}\left(\frac{t^3}{3}\right) = \frac{2}{9} \cdot \frac{\pi}{2} = \frac{\pi}{9}.$$

Convergent

24. $\int_0^{\infty} \frac{e^x}{e^{2x} + 3} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{e^x}{(e^x)^2 + (\sqrt{3})^2} dx = \lim_{t \rightarrow \infty} \left[\frac{1}{\sqrt{3}} \arctan \frac{e^x}{\sqrt{3}} \right]_0^t = \frac{1}{\sqrt{3}} \lim_{t \rightarrow \infty} \left(\arctan \frac{e^t}{\sqrt{3}} - \arctan \frac{1}{\sqrt{3}} \right)$

$$= \frac{1}{\sqrt{3}} \left(\frac{\pi}{2} - \frac{\pi}{6} \right) = \frac{1}{\sqrt{3}} \left(\frac{\pi}{3} \right) = \frac{\pi\sqrt{3}}{9}. \quad \text{Convergent}$$


25. $\int_e^{\infty} \frac{1}{x(\ln x)^3} dx = \lim_{t \rightarrow \infty} \int_e^t \frac{1}{x(\ln x)^3} dx = \lim_{t \rightarrow \infty} \int_1^{\ln t} u^{-3} du \begin{bmatrix} u = \ln x \\ du = dx/x \end{bmatrix} = \lim_{t \rightarrow \infty} \left[-\frac{1}{2u^2} \right]_1^{\ln t}$

$$= \lim_{t \rightarrow \infty} \left[-\frac{1}{2(\ln t)^2} + \frac{1}{2} \right] = 0 + \frac{1}{2} = \frac{1}{2}. \quad \text{Convergent}$$

26. $\int_0^{\infty} \frac{x \arctan x}{(1+x^2)^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{x \arctan x}{(1+x^2)^2} dx.$ Let $u = \arctan x, dv = \frac{x dx}{(1+x^2)^2}.$ Then $du = \frac{dx}{1+x^2},$

$$v = \frac{1}{2} \int \frac{2x dx}{(1+x^2)^2} = \frac{-1/2}{1+x^2}, \text{ and}$$

$$\begin{aligned} \int \frac{x \arctan x}{(1+x^2)^2} dx &= -\frac{1}{2} \frac{\arctan x}{1+x^2} + \frac{1}{2} \int \frac{dx}{(1+x^2)^2} \begin{bmatrix} x = \tan \theta, \\ dx = \sec^2 \theta d\theta \end{bmatrix} \\ &= -\frac{1}{2} \frac{\arctan x}{1+x^2} + \frac{1}{2} \int \frac{\sec^2 \theta d\theta}{(\sec^2 \theta)^2} \\ &= -\frac{1}{2} \frac{\arctan x}{1+x^2} + \frac{1}{2} \int \cos^2 \theta d\theta \\ &= -\frac{1}{2} \frac{\arctan x}{1+x^2} + \frac{\theta}{4} + \frac{\sin \theta \cos \theta}{4} + C \\ &= -\frac{1}{2} \frac{\arctan x}{1+x^2} + \frac{1}{4} \arctan x + \frac{1}{4} \frac{x}{1+x^2} + C \end{aligned}$$



It follows that

$$\begin{aligned} \int_0^{\infty} \frac{x \arctan x}{(1+x^2)^2} dx &= \lim_{t \rightarrow \infty} \left[-\frac{1}{2} \frac{\arctan x}{1+x^2} + \frac{1}{4} \arctan x + \frac{1}{4} \frac{x}{1+x^2} \right]_0^t \\ &= \lim_{t \rightarrow \infty} \left(-\frac{1}{2} \frac{\arctan t}{1+t^2} + \frac{1}{4} \arctan t + \frac{1}{4} \frac{t}{1+t^2} \right) = 0 + \frac{1}{4} \cdot \frac{\pi}{2} + 0 = \frac{\pi}{8}. \quad \text{Convergent} \end{aligned}$$

27. $\int_0^1 \frac{3}{x^5} dx = \lim_{t \rightarrow 0^+} \int_t^1 3x^{-5} dx = \lim_{t \rightarrow 0^+} \left[-\frac{3}{4x^4} \right]_t^1 = -\frac{3}{4} \lim_{t \rightarrow 0^+} \left(1 - \frac{1}{t^4} \right) = \infty. \quad \text{Divergent}$

28. $\int_2^3 \frac{1}{\sqrt{3-x}} dx = \lim_{t \rightarrow 3^-} \int_2^t (3-x)^{-1/2} dx = \lim_{t \rightarrow 3^-} \left[-2(3-x)^{1/2} \right]_2^t = -2 \lim_{t \rightarrow 3^-} (\sqrt{3-t} - \sqrt{1}) = -2(0-1) = 2.$

Convergent

29. $\int_{-2}^{14} \frac{dx}{\sqrt[4]{x+2}} = \lim_{t \rightarrow -2^+} \int_t^{14} (x+2)^{-1/4} dx = \lim_{t \rightarrow -2^+} \left[\frac{4}{3}(x+2)^{3/4} \right]_t^{14} = \frac{4}{3} \lim_{t \rightarrow -2^+} [16^{3/4} - (t+2)^{3/4}] \\ = \frac{4}{3}(8-0) = \frac{32}{3}. \quad \text{Convergent}$

30. $\int_6^8 \frac{4}{(x-6)^3} dx = \lim_{t \rightarrow 6^+} \int_t^8 4(x-6)^{-3} dx = \lim_{t \rightarrow 6^+} [-2(x-6)^{-2}]_t^8 = -2 \lim_{t \rightarrow 6^+} \left[\frac{1}{2^2} - \frac{1}{(t-6)^2} \right] = \infty. \quad \text{Divergent}$

31. $\int_{-2}^3 \frac{dx}{x^4} = \int_{-2}^0 \frac{dx}{x^4} + \int_0^3 \frac{dx}{x^4}, \text{ but } \int_{-2}^0 \frac{dx}{x^4} = \lim_{t \rightarrow 0^-} \left[-\frac{x^{-3}}{3} \right]_{-2}^t = \lim_{t \rightarrow 0^-} \left[-\frac{1}{3t^3} - \frac{1}{24} \right] = \infty. \quad \text{Divergent}$

32. $\int_0^1 \frac{dx}{\sqrt{1-x^2}} = \lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{\sqrt{1-x^2}} = \lim_{t \rightarrow 1^-} [\sin^{-1} x]_0^t = \lim_{t \rightarrow 1^-} \sin^{-1} t = \frac{\pi}{2}. \quad \text{Convergent}$

33. There is an infinite discontinuity at $x = 1$. $\int_0^{33} (x-1)^{-1/5} dx = \int_0^1 (x-1)^{-1/5} dx + \int_1^{33} (x-1)^{-1/5} dx$. Here

$$\int_0^1 (x-1)^{-1/5} dx = \lim_{t \rightarrow 1^-} \int_0^t (x-1)^{-1/5} dx = \lim_{t \rightarrow 1^-} \left[\frac{5}{4}(x-1)^{4/5} \right]_0^t = \lim_{t \rightarrow 1^-} \left[\frac{5}{4}(t-1)^{4/5} - \frac{5}{4} \right] = -\frac{5}{4} \text{ and}$$

$$\int_1^{33} (x-1)^{-1/5} dx = \lim_{t \rightarrow 1^+} \int_t^{33} (x-1)^{-1/5} dx = \lim_{t \rightarrow 1^+} \left[\frac{5}{4}(x-1)^{4/5} \right]_t^{33} = \lim_{t \rightarrow 1^+} \left[\frac{5}{4} \cdot 16 - \frac{5}{4}(t-1)^{4/5} \right] = 20.$$

Thus, $\int_0^{33} (x-1)^{-1/5} dx = -\frac{5}{4} + 20 = \frac{75}{4}. \quad \text{Convergent}$

34. $f(y) = 1/(4y-1)$ has an infinite discontinuity at $y = \frac{1}{4}$.

$$\int_{1/4}^1 \frac{1}{4y-1} dy = \lim_{t \rightarrow (1/4)^+} \int_t^1 \frac{1}{4y-1} dy = \lim_{t \rightarrow (1/4)^+} \left[\frac{1}{4} \ln |4y-1| \right]_t^1 = \lim_{t \rightarrow (1/4)^+} \left[\frac{1}{4} \ln 3 - \frac{1}{4} \ln(4t-1) \right] = \infty,$$

so $\int_{1/4}^1 \frac{1}{4y-1} dy$ diverges, and hence, $\int_0^1 \frac{1}{4y-1} dy$ diverges. Divergent

35. $I = \int_0^3 \frac{dx}{x^2 - 6x + 5} = \int_0^3 \frac{dx}{(x-1)(x-5)} = I_1 + I_2 = \int_0^1 \frac{dx}{(x-1)(x-5)} + \int_1^3 \frac{dx}{(x-1)(x-5)}.$

$$\text{Now } \frac{1}{(x-1)(x-5)} = \frac{A}{x-1} + \frac{B}{x-5} \Rightarrow 1 = A(x-5) + B(x-1).$$

Set $x = 5$ to get $1 = 4B$, so $B = \frac{1}{4}$. Set $x = 1$ to get $1 = -4A$, so $A = -\frac{1}{4}$. Thus

$$\begin{aligned} I_1 &= \lim_{t \rightarrow 1^-} \int_0^t \left(\frac{-\frac{1}{4}}{x-1} + \frac{\frac{1}{4}}{x-5} \right) dx = \lim_{t \rightarrow 1^-} \left[-\frac{1}{4} \ln|x-1| + \frac{1}{4} \ln|x-5| \right]_0^t \\ &= \lim_{t \rightarrow 1^-} [(-\frac{1}{4} \ln|t-1| + \frac{1}{4} \ln|t-5|) - (-\frac{1}{4} \ln|-1| + \frac{1}{4} \ln|-5|)] \\ &= \infty, \quad \text{since } \lim_{t \rightarrow 1^-} (-\frac{1}{4} \ln|t-1|) = \infty. \end{aligned}$$

Since I_1 is divergent, I is divergent.

36. $\int_{\pi/2}^{\pi} \csc x dx = \lim_{t \rightarrow \pi^-} \int_{\pi/2}^t \csc x dx = \lim_{t \rightarrow \pi^-} [\ln|\csc x - \cot x|]_{\pi/2}^t = \lim_{t \rightarrow \pi^-} [\ln(\csc t - \cot t) - \ln(1-0)] \\ = \lim_{t \rightarrow \pi^-} \ln \left(\frac{1-\cos t}{\sin t} \right) = \infty. \quad \text{Divergent}$

$$\begin{aligned}
 37. \int_{-1}^0 \frac{e^{1/x}}{x^3} dx &= \lim_{t \rightarrow 0^-} \int_{-1}^t \frac{1}{x} e^{1/x} \cdot \frac{1}{x^2} dx = \lim_{t \rightarrow 0^-} \int_{-1}^{1/t} ue^u (-du) \quad \left[\begin{array}{l} u = 1/x, \\ du = -dx/x^2 \end{array} \right] \\
 &= \lim_{t \rightarrow 0^-} [(u-1)e^u]_{1/t}^{-1} \quad \left[\begin{array}{l} \text{use parts} \\ \text{or Formula 96} \end{array} \right] = \lim_{t \rightarrow 0^-} \left[-2e^{-1} - \left(\frac{1}{t} - 1 \right) e^{1/t} \right] \\
 &= -\frac{2}{e} - \lim_{s \rightarrow -\infty} (s-1)e^s \quad [s = 1/t] = -\frac{2}{e} - \lim_{s \rightarrow -\infty} \frac{s-1}{e^{-s}} \stackrel{\text{H}}{=} -\frac{2}{e} - \lim_{s \rightarrow -\infty} \frac{1}{-e^{-s}} \\
 &= -\frac{2}{e} - 0 = -\frac{2}{e}. \quad \text{Convergent}
 \end{aligned}$$

$$\begin{aligned}
 38. \int_0^1 \frac{e^{1/x}}{x^3} dx &= \lim_{t \rightarrow 0^+} \int_t^1 \frac{1}{x} e^{1/x} \cdot \frac{1}{x^2} dx = \lim_{t \rightarrow 0^+} \int_{1/t}^1 ue^u (-du) \quad \left[\begin{array}{l} u = 1/x, \\ du = -dx/x^2 \end{array} \right] \\
 &= \lim_{t \rightarrow 0^+} [(u-1)e^u]_1^{1/t} \quad \left[\begin{array}{l} \text{use parts} \\ \text{or Formula 96} \end{array} \right] = \lim_{t \rightarrow 0^+} \left[\left(\frac{1}{t} - 1 \right) e^{1/t} - 0 \right] \\
 &= \lim_{s \rightarrow \infty} (s-1)e^s \quad [s = 1/t] = \infty. \quad \text{Divergent}
 \end{aligned}$$

$$\begin{aligned}
 39. I &= \int_0^2 z^2 \ln z dz = \lim_{t \rightarrow 0^+} \int_t^2 z^2 \ln z dz = \lim_{t \rightarrow 0^+} \left[\frac{z^3}{3^2} (3 \ln z - 1) \right]_t^2 \quad \left[\begin{array}{l} \text{integrate by parts} \\ \text{or use Formula 101} \end{array} \right] \\
 &= \lim_{t \rightarrow 0^+} \left[\frac{8}{9} (3 \ln 2 - 1) - \frac{1}{9} t^3 (3 \ln t - 1) \right] = \frac{8}{3} \ln 2 - \frac{8}{9} - \frac{1}{9} \lim_{t \rightarrow 0^+} [t^3 (3 \ln t - 1)] = \frac{8}{3} \ln 2 - \frac{8}{9} - \frac{1}{9} L.
 \end{aligned}$$



$$\text{Now } L = \lim_{t \rightarrow 0^+} [t^3 (3 \ln t - 1)] = \lim_{t \rightarrow 0^+} \frac{3 \ln t - 1}{t^{-3}} \stackrel{\text{H}}{=} \lim_{t \rightarrow 0^+} \frac{3/t}{-3/t^4} = \lim_{t \rightarrow 0^+} (-t^3) = 0.$$

Thus, $L = 0$ and $I = \frac{8}{3} \ln 2 - \frac{8}{9}$. Convergent


40. Integrate by parts with $u = \ln x$, $dv = dx/\sqrt{x}$ \Rightarrow $du = dx/x$, $v = 2\sqrt{x}$.

$$\begin{aligned}
 \int_0^1 \frac{\ln x}{\sqrt{x}} dx &= \lim_{t \rightarrow 0^+} \int_t^1 \frac{\ln x}{\sqrt{x}} dx = \lim_{t \rightarrow 0^+} \left(\left[2\sqrt{x} \ln x \right]_t^1 - 2 \int_t^1 \frac{dx}{\sqrt{x}} \right) = \lim_{t \rightarrow 0^+} \left(-2\sqrt{t} \ln t - 4 \left[\sqrt{x} \right]_t^1 \right) \\
 &= \lim_{t \rightarrow 0^+} (-2\sqrt{t} \ln t - 4 + 4\sqrt{t}) = -4
 \end{aligned}$$

$$\text{since } \lim_{t \rightarrow 0^+} \sqrt{t} \ln t = \lim_{t \rightarrow 0^+} \frac{\ln t}{t^{-1/2}} \stackrel{\text{H}}{=} \lim_{t \rightarrow 0^+} \frac{1/t}{-t^{-3/2}/2} = \lim_{t \rightarrow 0^+} (-2\sqrt{t}) = 0. \quad \text{Convergent}$$




43.



$$\begin{aligned} \text{Area} &= \int_{-\infty}^{\infty} \frac{2}{x^2 + 9} dx = 2 \cdot 2 \int_0^{\infty} \frac{1}{x^2 + 9} dx = 4 \lim_{t \rightarrow \infty} \int_0^t \frac{1}{x^2 + 9} dx \\ &= 4 \lim_{t \rightarrow \infty} \left[\frac{1}{3} \tan^{-1} \frac{x}{3} \right]_0^t = \frac{4}{3} \lim_{t \rightarrow \infty} \left[\tan^{-1} \frac{t}{3} - 0 \right] = \frac{4}{3} \cdot \frac{\pi}{2} = \frac{2\pi}{3} \end{aligned}$$


44.



$$\begin{aligned} \text{Area} &= \int_0^{\infty} \frac{x}{x^2 + 9} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{x}{x^2 + 9} dx = \lim_{t \rightarrow \infty} \left[\frac{1}{2} \ln(x^2 + 9) \right]_0^t \\ &= \frac{1}{2} \lim_{t \rightarrow \infty} [\ln(t^2 + 9) - \ln 9] = \infty \end{aligned}$$

Infinite area


45.



$$\begin{aligned} \text{Area} &= \int_0^{\pi/2} \sec^2 x dx = \lim_{t \rightarrow (\pi/2)^-} \int_0^t \sec^2 x dx = \lim_{t \rightarrow (\pi/2)^-} [\tan x]_0^t \\ &= \lim_{t \rightarrow (\pi/2)^-} (\tan t - 0) = \infty \end{aligned}$$

Infinite area

46.



$$\begin{aligned} \text{Area} &= \int_{-2}^0 \frac{1}{\sqrt{x+2}} dx = \lim_{t \rightarrow -2^+} \int_t^0 \frac{1}{\sqrt{x+2}} dx = \lim_{t \rightarrow -2^+} [2\sqrt{x+2}]_t^0 \\ &= \lim_{t \rightarrow -2^+} (2\sqrt{2} - 2\sqrt{t+2}) = 2\sqrt{2} - 0 = 2\sqrt{2} \end{aligned}$$

47. (a)


t	$\int_1^t g(x) dx$
2	0.447453
5	0.577101
10	0.621306
100	0.668479
1000	0.672957
10,000	0.673407

$$g(x) = \frac{\sin^2 x}{x^2}.$$

It appears that the integral is convergent.

(b) $-1 \leq \sin x \leq 1 \Rightarrow 0 \leq \sin^2 x \leq 1 \Rightarrow 0 \leq \frac{\sin^2 x}{x^2} \leq \frac{1}{x^2}$. Since $\int_1^{\infty} \frac{1}{x^2} dx$ is convergent

[Equation 2 with $p = 2 > 1$], $\int_1^{\infty} \frac{\sin^2 x}{x^2} dx$ is convergent by the Comparison Theorem.



Since $\int_1^\infty f(x) dx$ is finite and the area under $g(x)$ is less than the area under $f(x)$ on any interval $[1, t]$, $\int_1^\infty g(x) dx$ must be finite; that is, the integral is convergent.

48. (a)

t	$\int_2^t g(x) dx$
5	3.830327
10	6.801200
100	23.328769
1000	69.023361
10,000	208.124560


$$g(x) = \frac{1}{\sqrt{x} - 1}.$$

It appears that the integral is divergent.

(b) For $x \geq 2$, $\sqrt{x} > \sqrt{x} - 1 \Rightarrow \frac{1}{\sqrt{x}} < \frac{1}{\sqrt{x} - 1}$. Since $\int_2^\infty \frac{1}{\sqrt{x}} dx$ is divergent [Equation 2 with $p = \frac{1}{2} \leq 1$],

$\int_2^\infty \frac{1}{\sqrt{x} - 1} dx$ is divergent by the Comparison Theorem.

(c)



Since $\int_2^\infty f(x) dx$ is infinite and the area under $g(x)$ is greater than the area under $f(x)$ on any interval $[2, t]$, $\int_2^\infty g(x) dx$ must be infinite; that is, the integral is divergent.

49. For $x > 0$, $\frac{x}{x^3 + 1} < \frac{x}{x^3} = \frac{1}{x^2}$. $\int_1^\infty \frac{1}{x^2} dx$ is convergent by Equation 2 with $p = 2 > 1$, so $\int_1^\infty \frac{x}{x^3 + 1} dx$ is convergent

by the Comparison Theorem. $\int_0^1 \frac{x}{x^3 + 1} dx$ is a constant, so $\int_0^\infty \frac{x}{x^3 + 1} dx = \int_0^1 \frac{x}{x^3 + 1} dx + \int_1^\infty \frac{x}{x^3 + 1} dx$ is also convergent.

50. For $x \geq 1$, $\frac{2 + e^{-x}}{x} > \frac{2}{x}$ [since $e^{-x} > 0$] $> \frac{1}{x}$. $\int_1^\infty \frac{1}{x} dx$ is divergent by Equation 2 with $p = 1 \leq 1$, so

$\int_1^\infty \frac{2 + e^{-x}}{x} dx$ is divergent by the Comparison Theorem.

51. For $x > 1$, $f(x) = \frac{x+1}{\sqrt{x^4-x}} > \frac{x+1}{\sqrt{x^4}} > \frac{x}{\sqrt{x^4}} = \frac{1}{x^2}$, so $\int_2^\infty f(x) dx$ diverges by comparison with $\int_2^\infty \frac{1}{x^2} dx$, which diverges

by Equation 2 with $p = 1 \leq 1$. Thus, $\int_1^\infty f(x) dx = \int_1^2 f(x) dx + \int_2^\infty f(x) dx$ also diverges.

52. For $x \geq 0$, $\arctan x < \frac{\pi}{2} < 2$, so $\frac{\arctan x}{2+e^x} < \frac{2}{2+e^x} < \frac{2}{e^x} = 2e^{-x}$. Now

$$I = \int_0^\infty 2e^{-x} dx = \lim_{t \rightarrow \infty} \int_0^t 2e^{-x} dx = \lim_{t \rightarrow \infty} [-2e^{-x}]_0^t = \lim_{t \rightarrow \infty} \left(-\frac{2}{e^t} + 2 \right) = 2, \text{ so } I \text{ is convergent, and by comparison,}$$

$\int_0^\infty \frac{\arctan x}{2+e^x} dx$ is convergent.

53. For $0 < x \leq 1$, $\frac{\sec^2 x}{x\sqrt{x}} > \frac{1}{x^{3/2}}$. Now

$$I = \int_0^1 x^{-3/2} dx = \lim_{t \rightarrow 0^+} \int_t^1 x^{-3/2} dx = \lim_{t \rightarrow 0^+} \left[-2x^{-1/2} \right]_t^1 = \lim_{t \rightarrow 0^+} \left(-2 + \frac{2}{\sqrt{t}} \right) = \infty, \text{ so } I \text{ is divergent, and by}$$

comparison, $\int_0^1 \frac{\sec^2 x}{x\sqrt{x}} dx$ is divergent.

54. For $0 < x \leq 1$, $\frac{\sin^2 x}{\sqrt{x}} \leq \frac{1}{\sqrt{x}}$. Now

$$I = \int_0^\pi \frac{1}{\sqrt{x}} dx = \lim_{t \rightarrow 0^+} \int_t^\pi x^{-1/2} dx = \lim_{t \rightarrow 0^+} \left[2x^{1/2} \right]_t^\pi = \lim_{t \rightarrow 0^+} (2\pi - 2\sqrt{t}) = 2\pi - 0 = 2\pi, \text{ so } I \text{ is convergent, and by}$$

comparison, $\int_0^\pi \frac{\sin^2 x}{\sqrt{x}} dx$ is convergent.

55. $\int_0^\infty \frac{dx}{\sqrt{x}(1+x)} = \int_0^1 \frac{dx}{\sqrt{x}(1+x)} + \int_1^\infty \frac{dx}{\sqrt{x}(1+x)} = \lim_{t \rightarrow 0^+} \int_t^1 \frac{dx}{\sqrt{x}(1+x)} + \lim_{t \rightarrow \infty} \int_1^t \frac{dx}{\sqrt{x}(1+x)}$. Now

$$\int \frac{dx}{\sqrt{x}(1+x)} = \int \frac{2u du}{u(1+u^2)} \quad \left[u = \sqrt{x}, x = u^2, \frac{du}{dx} = \frac{1}{2u} \right] = 2 \int \frac{du}{1+u^2} = 2 \tan^{-1} u + C = 2 \tan^{-1} \sqrt{x} + C, \text{ so}$$

$$\begin{aligned} \int_0^\infty \frac{dx}{\sqrt{x}(1+x)} &= \lim_{t \rightarrow 0^+} [2 \tan^{-1} \sqrt{x}]_t^1 + \lim_{t \rightarrow \infty} [2 \tan^{-1} \sqrt{x}]_1^t \\ &= \lim_{t \rightarrow 0^+} [2(\frac{\pi}{4}) - 2 \tan^{-1} \sqrt{t}] + \lim_{t \rightarrow \infty} [2 \tan^{-1} \sqrt{t} - 2(\frac{\pi}{4})] = \frac{\pi}{2} - 0 + 2(\frac{\pi}{2}) - \frac{\pi}{2} = \pi. \end{aligned}$$

56. $\int_2^\infty \frac{dx}{x\sqrt{x^2-4}} = \int_2^3 \frac{dx}{x\sqrt{x^2-4}} + \int_3^\infty \frac{dx}{x\sqrt{x^2-4}} = \lim_{t \rightarrow 2^+} \int_t^3 \frac{dx}{x\sqrt{x^2-4}} + \lim_{t \rightarrow \infty} \int_3^t \frac{dx}{x\sqrt{x^2-4}}$. Now

$$\int \frac{dx}{x\sqrt{x^2-4}} = \int \frac{2 \sec \theta \tan \theta d\theta}{2 \sec \theta 2 \tan \theta} \quad \left[\begin{array}{l} x = 2 \sec \theta, \text{ where} \\ 0 \leq \theta < \pi/2 \text{ or } \pi \leq \theta < 3\pi/2 \end{array} \right] = \frac{1}{2}\theta + C = \frac{1}{2}\sec^{-1}(\frac{1}{2}x) + C, \text{ so}$$

$$\int_2^\infty \frac{dx}{x\sqrt{x^2-4}} = \lim_{t \rightarrow 2^+} [\frac{1}{2}\sec^{-1}(\frac{1}{2}x)]_t^3 + \lim_{t \rightarrow \infty} [\frac{1}{2}\sec^{-1}(\frac{1}{2}x)]_3^t = \frac{1}{2}\sec^{-1}(\frac{3}{2}) - 0 + \frac{1}{2}(\frac{\pi}{2}) - \frac{1}{2}\sec^{-1}(\frac{3}{2}) = \frac{\pi}{4}.$$

57. If $p = 1$, then $\int_0^1 \frac{dx}{x^p} = \lim_{t \rightarrow 0^+} \int_t^1 \frac{dx}{x} = \lim_{t \rightarrow 0^+} [\ln x]_t^1 = \infty$. Divergent.

If $p \neq 1$, then $\int_0^1 \frac{dx}{x^p} = \lim_{t \rightarrow 0^+} \int_t^1 \frac{dx}{x^p}$ [note that the integral is not improper if $p < 0$]

$$= \lim_{t \rightarrow 0^+} \left[\frac{x^{-p+1}}{-p+1} \right]_t^1 = \lim_{t \rightarrow 0^+} \frac{1}{1-p} \left[1 - \frac{1}{t^{p-1}} \right]$$

If $p > 1$, then $p - 1 > 0$, so $\frac{1}{t^{p-1}} \rightarrow \infty$ as $t \rightarrow 0^+$, and the integral diverges.

If $p < 1$, then $p - 1 < 0$, so $\frac{1}{t^{p-1}} \rightarrow 0$ as $t \rightarrow 0^+$ and $\int_0^1 \frac{dx}{x^p} = \frac{1}{1-p} \left[\lim_{t \rightarrow 0^+} (1 - t^{1-p}) \right] = \frac{1}{1-p}$.

Thus, the integral converges if and only if $p < 1$, and in that case its value is $\frac{1}{1-p}$.

58. Let $u = \ln x$. Then $du = dx/x \Rightarrow \int_e^\infty \frac{dx}{x(\ln x)^p} = \int_1^\infty \frac{du}{u^p}$. By Example 4, this converges to $\frac{1}{p-1}$ if $p > 1$ and diverges otherwise.

59. First suppose $p = -1$. Then

$\int_0^1 x^p \ln x \, dx = \int_0^1 \frac{\ln x}{x} \, dx = \lim_{t \rightarrow 0^+} \int_t^1 \frac{\ln x}{x} \, dx = \lim_{t \rightarrow 0^+} \left[\frac{1}{2} (\ln x)^2 \right]_t^1 = -\frac{1}{2} \lim_{t \rightarrow 0^+} (\ln t)^2 = -\infty$, so the

integral diverges. Now suppose $p \neq -1$. Then integration by parts gives

$\int x^p \ln x \, dx = \frac{x^{p+1}}{p+1} \ln x - \int \frac{x^p}{p+1} \, dx = \frac{x^{p+1}}{p+1} \ln x - \frac{x^{p+1}}{(p+1)^2} + C$. If $p < -1$, then $p+1 < 0$, so

$\int_0^1 x^p \ln x \, dx = \lim_{t \rightarrow 0^+} \left[\frac{x^{p+1}}{p+1} \ln x - \frac{x^{p+1}}{(p+1)^2} \right]_t^1 = \frac{-1}{(p+1)^2} - \left(\frac{1}{p+1} \right) \lim_{t \rightarrow 0^+} \left[t^{p+1} \left(\ln t - \frac{1}{p+1} \right) \right] = \infty$.

If $p > -1$, then $p+1 > 0$ and

$$\begin{aligned} \int_0^1 x^p \ln x \, dx &= \frac{-1}{(p+1)^2} - \left(\frac{1}{p+1} \right) \lim_{t \rightarrow 0^+} \frac{\ln t - 1/(p+1)}{t^{-(p+1)}} \stackrel{H}{=} \frac{-1}{(p+1)^2} - \left(\frac{1}{p+1} \right) \lim_{t \rightarrow 0^+} \frac{1/t}{-(p+1)t^{-(p+2)}} \\ &= \frac{-1}{(p+1)^2} + \frac{1}{(p+1)^2} \lim_{t \rightarrow 0^+} t^{p+1} = \frac{-1}{(p+1)^2} \end{aligned}$$

Thus, the integral converges to $-\frac{1}{(p+1)^2}$ if $p > -1$ and diverges otherwise.

60. (a) $n = 0$: $\int_0^\infty x^n e^{-x} \, dx = \lim_{t \rightarrow \infty} \int_0^t e^{-x} \, dx = \lim_{t \rightarrow \infty} [-e^{-x}]_0^t = \lim_{t \rightarrow \infty} [-e^{-t} + 1] = 0 + 1 = 1$

$n = 1$: $\int_0^\infty x^n e^{-x} \, dx = \lim_{t \rightarrow \infty} \int_0^t x e^{-x} \, dx$. To evaluate $\int x e^{-x} \, dx$, we'll use integration by parts with $u = x$, $dv = e^{-x} \, dx \Rightarrow du = dx$, $v = -e^{-x}$.

So $\int x e^{-x} \, dx = -x e^{-x} - \int -e^{-x} \, dx = -x e^{-x} - e^{-x} + C = (-x - 1)e^{-x} + C$ and

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_0^t x e^{-x} \, dx &= \lim_{t \rightarrow \infty} [(-x - 1)e^{-x}]_0^t = \lim_{t \rightarrow \infty} [(-t - 1)e^{-t} + 1] = \lim_{t \rightarrow \infty} [-te^{-t} - e^{-t} + 1] \\ &= 0 - 0 + 1 \quad [\text{use l'Hospital's Rule}] = 1 \end{aligned}$$

n = 2: $\int_0^\infty x^n e^{-x} dx = \lim_{t \rightarrow \infty} \int_0^t x^2 e^{-x} dx$. To evaluate $\int x^2 e^{-x} dx$, we could use integration by parts

again or Formula 97. Thus,

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_0^t x^2 e^{-x} dx &= \lim_{t \rightarrow \infty} [-x^2 e^{-x}]_0^t + 2 \lim_{t \rightarrow \infty} \int_0^t x e^{-x} dx \\ &= 0 + 0 + 2(1) \quad [\text{use l'Hospital's Rule and the result for } n = 1] = 2 \end{aligned}$$

$$\begin{aligned} \mathbf{n = 3:} \quad \int_0^\infty x^n e^{-x} dx &= \lim_{t \rightarrow \infty} \int_0^t x^3 e^{-x} dx \stackrel{?}{=} \lim_{t \rightarrow \infty} [-x^3 e^{-x}]_0^t + 3 \lim_{t \rightarrow \infty} \int_0^t x^2 e^{-x} dx \\ &= 0 + 0 + 3(2) \quad [\text{use l'Hospital's Rule and the result for } n = 2] = 6 \end{aligned}$$

(b) For $n = 1, 2$, and 3 , we have $\int_0^\infty x^n e^{-x} dx = 1, 2$, and 6 . The values for the integral are equal to the factorials for n , so

we guess $\int_0^\infty x^n e^{-x} dx = n!$.

(c) Suppose that $\int_0^\infty x^k e^{-x} dx = k!$ for some positive integer k . Then $\int_0^\infty x^{k+1} e^{-x} dx = \lim_{t \rightarrow \infty} \int_0^t x^{k+1} e^{-x} dx$.

To evaluate $\int x^{k+1} e^{-x} dx$, we use parts with $u = x^{k+1}$, $dv = e^{-x} dx \Rightarrow du = (k+1)x^k dx$, $v = -e^{-x}$.

So $\int x^{k+1} e^{-x} dx = -x^{k+1} e^{-x} - \int -(k+1)x^k e^{-x} dx = -x^{k+1} e^{-x} + (k+1) \int x^k e^{-x} dx$ and

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_0^t x^{k+1} e^{-x} dx &= \lim_{t \rightarrow \infty} [-x^{k+1} e^{-x}]_0^t + (k+1) \lim_{t \rightarrow \infty} \int_0^t x^k e^{-x} dx \\ &= \lim_{t \rightarrow \infty} [-t^{k+1} e^{-t} + 0] + (k+1)k! = 0 + 0 + (k+1)! = (k+1)! \end{aligned}$$

so the formula holds for $k+1$. By induction, the formula holds for all positive integers. (Since $0! = 1$, the formula holds for $n = 0$, too.)

61. (a) $I = \int_{-\infty}^\infty x dx = \int_{-\infty}^0 x dx + \int_0^\infty x dx$, and $\int_0^\infty x dx = \lim_{t \rightarrow \infty} \int_0^t x dx = \lim_{t \rightarrow \infty} [\frac{1}{2}x^2]_0^t = \lim_{t \rightarrow \infty} [\frac{1}{2}t^2 - 0] = \infty$,

so I is divergent.

(b) $\int_{-t}^t x dx = [\frac{1}{2}x^2]_{-t}^t = \frac{1}{2}t^2 - \frac{1}{2}(-t)^2 = 0$, so $\lim_{t \rightarrow \infty} \int_{-t}^t x dx = 0$. Therefore, $\int_{-\infty}^\infty x dx \neq \lim_{t \rightarrow \infty} \int_{-t}^t x dx$.

62. Let $k = \frac{M}{2RT}$ so that $\bar{v} = \frac{4}{\sqrt{\pi}} k^{3/2} \int_0^\infty v^3 e^{-kv^2} dv$. Let I denote the integral and use parts to integrate I . Let $\alpha = v^2$,

$$d\beta = ve^{-kv^2} dv \Rightarrow d\alpha = 2v dv, \beta = -\frac{1}{2k} e^{-kv^2}$$

$$I = \lim_{t \rightarrow \infty} \left[-\frac{1}{2k} v^2 e^{-kv^2} \right]_0^t + \frac{1}{k} \int_0^\infty ve^{-kv^2} dv \Big|_0^t = -\frac{1}{2k} \lim_{t \rightarrow \infty} (t^2 e^{-kt^2}) + \frac{1}{k} \lim_{t \rightarrow \infty} \left[-\frac{1}{2k} e^{-kv^2} \right]$$

$$\stackrel{H}{=} -\frac{1}{2k} \cdot 0 - \frac{1}{2k^2} (0 - 1) = \frac{1}{2k^2}$$

$$\text{Thus, } \bar{v} = \frac{4}{\sqrt{\pi}} k^{3/2} \cdot \frac{1}{2k^2} = \frac{2}{(k\pi)^{1/2}} = \frac{2}{[\pi M / (2RT)]^{1/2}} = \frac{2\sqrt{2}\sqrt{RT}}{\sqrt{\pi M}} = \sqrt{\frac{8RT}{\pi M}}$$

$$63. \text{ Volume} = \int_1^\infty \pi \left(\frac{1}{x}\right)^2 dx = \pi \lim_{t \rightarrow \infty} \int_1^t \frac{dx}{x^2} = \pi \lim_{t \rightarrow \infty} \left[-\frac{1}{x}\right]_1^t = \pi \lim_{t \rightarrow \infty} \left(1 - \frac{1}{t}\right) = \pi < \infty$$

64. Work = $\int_R^\infty \frac{GMm}{r^2} dr = \lim_{t \rightarrow \infty} \int_R^t \frac{GMm}{r^2} dr = \lim_{t \rightarrow \infty} GMm \left[\frac{-1}{r} \right]_R^t = GMm \lim_{t \rightarrow \infty} \left(\frac{-1}{t} + \frac{1}{R} \right) = \frac{GMm}{R}$, where
 M = mass of the earth = 5.98×10^{24} kg, m = mass of satellite = 10^3 kg, R = radius of the earth = 6.37×10^6 m, and
 G = gravitational constant = 6.67×10^{-11} N·m²/kg.

Therefore, Work = $\frac{6.67 \times 10^{-11} \cdot 5.98 \times 10^{24} \cdot 10^3}{6.37 \times 10^6} \approx 6.26 \times 10^{10}$ J.

65. Work = $\int_R^\infty F dr = \lim_{t \rightarrow \infty} \int_R^t \frac{GmM}{r^2} dr = \lim_{t \rightarrow \infty} GmM \left(\frac{1}{R} - \frac{1}{t} \right) = \frac{GmM}{R}$. The initial kinetic energy provides the work,
so $\frac{1}{2}mv_0^2 = \frac{GmM}{R} \Rightarrow v_0 = \sqrt{\frac{2GM}{R}}$.

66. $y(s) = \int_s^R \frac{2r}{\sqrt{r^2 - s^2}} x(r) dr$ and $x(r) = \frac{1}{2}(R - r)^2 \Rightarrow$

$$\begin{aligned} y(s) &= \lim_{t \rightarrow s^+} \int_t^R \frac{r(R - r)^2}{\sqrt{r^2 - s^2}} dr = \lim_{t \rightarrow s^+} \int_t^R \frac{r^3 - 2Rr^2 + R^2r}{\sqrt{r^2 - s^2}} dr \\ &= \lim_{t \rightarrow s^+} \left[\int_t^R \frac{r^3 dr}{\sqrt{r^2 - s^2}} - 2R \int_t^R \frac{r^2 dr}{\sqrt{r^2 - s^2}} + R^2 \int_t^R \frac{r dr}{\sqrt{r^2 - s^2}} \right] = \lim_{t \rightarrow s^+} (I_1 - 2RI_2 + R^2I_3) = L \end{aligned}$$

For I_1 : Let $u = \sqrt{r^2 - s^2} \Rightarrow u^2 = r^2 - s^2$, $r^2 = u^2 + s^2$, $2r dr = 2u du$, so, omitting limits and constant of integration,

$$\begin{aligned} I_1 &= \int \frac{(u^2 + s^2)u}{u} du = \int (u^2 + s^2) du = \frac{1}{3}u^3 + s^2u = \frac{1}{3}u(u^2 + 3s^2) \\ &= \frac{1}{3}\sqrt{r^2 - s^2}(r^2 - s^2 + 3s^2) = \frac{1}{3}\sqrt{r^2 - s^2}(r^2 + 2s^2) \end{aligned}$$


For I_2 : Using Formula 44, $I_2 = \frac{r}{2}\sqrt{r^2 - s^2} + \frac{s^2}{2} \ln|r + \sqrt{r^2 - s^2}|$.

For I_3 : Let $u = r^2 - s^2 \Rightarrow du = 2r dr$. Then $I_3 = \frac{1}{2} \int \frac{du}{\sqrt{u}} = \frac{1}{2} \cdot 2\sqrt{u} = \sqrt{r^2 - s^2}$.

Thus,

$$\begin{aligned} L &= \lim_{t \rightarrow s^+} \left[\frac{1}{3}\sqrt{r^2 - s^2}(r^2 + 2s^2) - 2R \left(\frac{r}{2}\sqrt{r^2 - s^2} + \frac{s^2}{2} \ln|r + \sqrt{r^2 - s^2}| \right) + R^2\sqrt{r^2 - s^2} \right]_t^R \\ &= \lim_{t \rightarrow s^+} \left[\frac{1}{3}\sqrt{R^2 - s^2}(R^2 + 2s^2) - 2R \left(\frac{R}{2}\sqrt{R^2 - s^2} + \frac{s^2}{2} \ln|R + \sqrt{R^2 - s^2}| \right) + R^2\sqrt{R^2 - s^2} \right] \\ &\quad - \lim_{t \rightarrow s^+} \left[\frac{1}{3}\sqrt{t^2 - s^2}(t^2 + 2s^2) - 2R \left(\frac{t}{2}\sqrt{t^2 - s^2} + \frac{s^2}{2} \ln|t + \sqrt{t^2 - s^2}| \right) + R^2\sqrt{t^2 - s^2} \right] \\ &= \left[\frac{1}{3}\sqrt{R^2 - s^2}(R^2 + 2s^2) - Rs^2 \ln|R + \sqrt{R^2 - s^2}| \right] - \left[-Rs^2 \ln|s| \right] \\ &= \frac{1}{3}\sqrt{R^2 - s^2}(R^2 + 2s^2) - Rs^2 \ln\left(\frac{R + \sqrt{R^2 - s^2}}{s}\right) \end{aligned}$$

67. We would expect a small percentage of bulbs to burn out in the first few hundred hours, most of the bulbs to burn out after close to 700 hours, and a few overachievers to burn on and on.



(b) $r(t) = F'(t)$ is the rate at which the fraction $F(t)$ of burnt-out bulbs increases as t increases. This could be interpreted as a fractional burnout rate.

(c) $\int_0^\infty r(t) dt = \lim_{x \rightarrow \infty} F(x) = 1$, since all of the bulbs will eventually burn out.

$$68. I = \int_0^\infty te^{kt} dt = \lim_{s \rightarrow \infty} \left[\frac{1}{k^2} (kt - 1) e^{kt} \right]_0^s \quad [\text{Formula 96, or parts}] = \lim_{s \rightarrow \infty} \left[\left(\frac{1}{k} se^{ks} - \frac{1}{k^2} e^{ks} \right) - \left(-\frac{1}{k^2} \right) \right].$$

Since $k < 0$ the first two terms approach 0 (you can verify that the first term does so with l'Hospital's Rule), so the limit is equal to $1/k^2$. Thus, $M = -kI = -k(1/k^2) = -1/k = -1/(-0.000121) \approx 8264.5$ years.

$$69. I = \int_a^\infty \frac{1}{x^2 + 1} dx = \lim_{t \rightarrow \infty} \int_a^t \frac{1}{x^2 + 1} dx = \lim_{t \rightarrow \infty} [\tan^{-1} x]_a^t = \lim_{t \rightarrow \infty} (\tan^{-1} t - \tan^{-1} a) = \frac{\pi}{2} - \tan^{-1} a.$$

$I < 0.001 \Rightarrow \frac{\pi}{2} - \tan^{-1} a < 0.001 \Rightarrow \tan^{-1} a > \frac{\pi}{2} - 0.001 \Rightarrow a > \tan(\frac{\pi}{2} - 0.001) \approx 1000.$

70. $f(x) = e^{-x^2}$ and $\Delta x = \frac{4-0}{8} = \frac{1}{2}$.

$$\int_0^4 f(x) dx \approx S_8 = \frac{1}{2 \cdot 3} [f(0) + 4f(0.5) + 2f(1) + \dots + 2f(3) + 4f(3.5) + f(4)] \approx \frac{1}{6}(5.31717808) \approx 0.8862$$

$$\text{Now } x > 4 \Rightarrow -x \cdot x < -x \cdot 4 \Rightarrow e^{-x^2} < e^{-4x} \Rightarrow \int_4^\infty e^{-x^2} dx < \int_4^\infty e^{-4x} dx.$$

$$\int_4^\infty e^{-4x} dx = \lim_{t \rightarrow \infty} [-\frac{1}{4} e^{-4x}]_4^t = -\frac{1}{4}(0 - e^{-16}) = 1/(4e^{16}) \approx 0.0000000281 < 0.0000001, \text{ as desired.}$$

71. (a) $F(s) = \int_0^\infty f(t)e^{-st} dt = \int_0^\infty e^{-st} dt = \lim_{n \rightarrow \infty} \left[-\frac{e^{-st}}{s} \right]_0^n = \lim_{n \rightarrow \infty} \left(\frac{e^{-sn}}{-s} + \frac{1}{s} \right).$ This converges to $\frac{1}{s}$ only if $s > 0$.

Therefore $F(s) = \frac{1}{s}$ with domain $\{s \mid s > 0\}$.

$$(b) F(s) = \int_0^\infty f(t)e^{-st} dt = \int_0^\infty e^t e^{-st} dt = \lim_{n \rightarrow \infty} \int_0^n e^{t(1-s)} dt = \lim_{n \rightarrow \infty} \left[\frac{1}{1-s} e^{t(1-s)} \right]_0^n \\ = \lim_{n \rightarrow \infty} \left(\frac{e^{(1-s)n}}{1-s} - \frac{1}{1-s} \right)$$

This converges only if $1-s < 0 \Rightarrow s > 1$, in which case $F(s) = \frac{1}{s-1}$ with domain $\{s \mid s > 1\}$.

(c) $F(s) = \int_0^\infty f(t)e^{-st} dt = \lim_{n \rightarrow \infty} \int_0^n te^{-st} dt.$ Use integration by parts: let $u = t$, $dv = e^{-st} dt \Rightarrow du = dt$,

$$v = -\frac{e^{-st}}{s}. \text{ Then } F(s) = \lim_{n \rightarrow \infty} \left[-\frac{t}{s} e^{-st} - \frac{1}{s^2} e^{-st} \right]_0^n = \lim_{n \rightarrow \infty} \left(\frac{-n}{se^{sn}} - \frac{1}{s^2 e^{sn}} + 0 + \frac{1}{s^2} \right) = \frac{1}{s^2} \text{ only if } s > 0.$$

Therefore, $F(s) = \frac{1}{s^2}$ and the domain of F is $\{s \mid s > 0\}$.

72. $0 \leq f(t) \leq M e^{at} \Rightarrow 0 \leq f(t)e^{-st} \leq M e^{at} e^{-st}$ for $t \geq 0$. Now use the Comparison Theorem:

$$\int_0^\infty M e^{at} e^{-st} dt = \lim_{n \rightarrow \infty} M \int_0^n e^{t(a-s)} dt = M \cdot \lim_{n \rightarrow \infty} \left[\frac{1}{a-s} e^{t(a-s)} \right]_0^n = M \cdot \lim_{n \rightarrow \infty} \frac{1}{a-s} [e^{n(a-s)} - 1]$$

This is convergent only when $a - s < 0 \Rightarrow s > a$. Therefore, by the Comparison Theorem, $F(s) = \int_0^\infty f(t) e^{-st} dt$ is

also convergent for $s > a$.

73. $G(s) = \int_0^\infty f'(t) e^{-st} dt$. Integrate by parts with $u = e^{-st}$, $dv = f'(t) dt \Rightarrow du = -s e^{-st}$, $v = f(t)$:

$$G(s) = \lim_{n \rightarrow \infty} [f(t)e^{-st}]_0^n + s \int_0^\infty f(t)e^{-st} dt = \lim_{n \rightarrow \infty} f(n)e^{-sn} - f(0) + sF(s)$$

But $0 \leq f(t) \leq M e^{at} \Rightarrow 0 \leq f(t)e^{-st} \leq M e^{at} e^{-st}$ and $\lim_{t \rightarrow \infty} M e^{t(a-s)} = 0$ for $s > a$. So by the Squeeze Theorem,

$\lim_{t \rightarrow \infty} f(t)e^{-st} = 0$ for $s > a \Rightarrow G(s) = 0 - f(0) + sF(s) = sF(s) - f(0)$ for $s > a$.

74. Assume without loss of generality that $a < b$. Then

$$\begin{aligned} \int_{-\infty}^a f(x) dx + \int_a^\infty f(x) dx &= \lim_{t \rightarrow -\infty} \int_t^a f(x) dx + \lim_{u \rightarrow \infty} \int_a^u f(x) dx \\ &= \lim_{t \rightarrow -\infty} \int_t^a f(x) dx + \lim_{u \rightarrow \infty} \left[\int_a^b f(x) dx + \int_b^u f(x) dx \right] \\ &= \lim_{t \rightarrow -\infty} \int_t^a f(x) dx + \int_a^b f(x) dx + \lim_{u \rightarrow \infty} \int_b^u f(x) dx \\ &= \lim_{t \rightarrow -\infty} \left[\int_t^a f(x) dx + \int_a^b f(x) dx \right] + \int_b^\infty f(x) dx \\ &= \lim_{t \rightarrow -\infty} \int_t^b f(x) dx + \int_b^\infty f(x) dx = \int_{-\infty}^b f(x) dx + \int_b^\infty f(x) dx \end{aligned}$$

75. We use integration by parts: let $u = x$, $dv = x e^{-x^2} dx \Rightarrow du = dx$, $v = -\frac{1}{2} e^{-x^2}$. So

$$\int_0^\infty x^2 e^{-x^2} dx = \lim_{t \rightarrow \infty} \left[-\frac{1}{2} x e^{-x^2} \right]_0^t + \frac{1}{2} \int_0^\infty e^{-x^2} dx = \lim_{t \rightarrow \infty} \left[-\frac{t}{2e^{t^2}} \right] + \frac{1}{2} \int_0^\infty e^{-x^2} dx = \frac{1}{2} \int_0^\infty e^{-x^2} dx$$

(The limit is 0 by l'Hospital's Rule.)


76. $\int_0^\infty e^{-x^2} dx$ is the area under the curve $y = e^{-x^2}$ for $0 \leq x < \infty$ and $0 < y \leq 1$. Solving $y = e^{-x^2}$ for x , we get

$y = e^{-x^2} \Rightarrow \ln y = -x^2 \Rightarrow -\ln y = x^2 \Rightarrow x = \pm\sqrt{-\ln y}$. Since x is positive, choose $x = \sqrt{-\ln y}$, and the area is represented by $\int_0^1 \sqrt{-\ln y} dy$. Therefore, each integral represents the same area, so the integrals are equal.

77. For the first part of the integral, let $x = 2 \tan \theta \Rightarrow dx = 2 \sec^2 \theta d\theta$.

$$\int \frac{1}{\sqrt{x^2 + 4}} dx = \int \frac{2 \sec^2 \theta}{2 \sec \theta} d\theta = \int \sec \theta d\theta = \ln |\sec \theta + \tan \theta|.$$

From the figure, $\tan \theta = \frac{x}{2}$, and $\sec \theta = \frac{\sqrt{x^2 + 4}}{2}$. So



$$\begin{aligned}
 I &= \int_0^\infty \left(\frac{1}{\sqrt{x^2 + 4}} - \frac{C}{x+2} \right) dx = \lim_{t \rightarrow \infty} \left[\ln \left| \frac{\sqrt{x^2 + 4}}{2} + \frac{x}{2} \right| - C \ln|x+2| \right]_0^t \\
 &= \lim_{t \rightarrow \infty} \left[\ln \frac{\sqrt{t^2 + 4} + t}{2} - C \ln(t+2) - (\ln 1 - C \ln 2) \right] \\
 &= \lim_{t \rightarrow \infty} \left[\ln \left(\frac{\sqrt{t^2 + 4} + t}{2(t+2)^C} \right) + \ln 2^C \right] = \ln \left(\lim_{t \rightarrow \infty} \frac{t + \sqrt{t^2 + 4}}{(t+2)^C} \right) + \ln 2^{C-1}
 \end{aligned}$$

$$\text{Now } L = \lim_{t \rightarrow \infty} \frac{t + \sqrt{t^2 + 4}}{(t+2)^C} \stackrel{H}{=} \lim_{t \rightarrow \infty} \frac{1 + t/\sqrt{t^2 + 4}}{C(t+2)^{C-1}} = \frac{2}{C \lim_{t \rightarrow \infty} (t+2)^{C-1}}.$$

If $C < 1$, $L = \infty$ and I diverges.

If $C = 1$, $L = 2$ and I converges to $\ln 2 + \ln 2^0 = \ln 2$.

If $C > 1$, $L = 0$ and I diverges to $-\infty$.

$$\begin{aligned}
 78. I &= \int_0^\infty \left(\frac{x}{x^2 + 1} - \frac{C}{3x + 1} \right) dx = \lim_{t \rightarrow \infty} \left[\frac{1}{2} \ln(x^2 + 1) - \frac{1}{3} C \ln(3x + 1) \right]_0^t = \lim_{t \rightarrow \infty} \left[\ln(t^2 + 1)^{1/2} - \ln(3t + 1)^{C/3} \right] \\
 &= \lim_{t \rightarrow \infty} \left(\ln \frac{(t^2 + 1)^{1/2}}{(3t + 1)^{C/3}} \right) = \ln \left(\lim_{t \rightarrow \infty} \frac{\sqrt{t^2 + 1}}{(3t + 1)^{C/3}} \right)
 \end{aligned}$$

For $C \leq 0$, the integral diverges. For $C > 0$, we have

$$L = \lim_{t \rightarrow \infty} \frac{\sqrt{t^2 + 1}}{(3t + 1)^{C/3}} \stackrel{H}{=} \lim_{t \rightarrow \infty} \frac{t/\sqrt{t^2 + 1}}{C(3t + 1)^{(C/3)-1}} = \frac{1}{C} \lim_{t \rightarrow \infty} \frac{1}{(3t + 1)^{(C/3)-1}}$$

For $C/3 < 1 \Leftrightarrow C < 3$, $L = \infty$ and I diverges.

For $C = 3$, $L = \frac{1}{3}$ and $I = \ln \frac{1}{3}$.

For $C > 3$, $L = 0$ and I diverges to $-\infty$.

79. No, $I = \int_0^\infty f(x) dx$ must be divergent. Since $\lim_{x \rightarrow \infty} f(x) = 1$, there must exist an N such that if $x \geq N$, then $f(x) \geq \frac{1}{2}$.

Thus, $I = I_1 + I_2 = \int_0^N f(x) dx + \int_N^\infty f(x) dx$, where I_1 is an ordinary definite integral that has a finite value, and I_2 is improper and diverges by comparison with the divergent integral $\int_N^\infty \frac{1}{2} dx$.

80. As in Exercise 55, we let $I = \int_0^\infty \frac{x^a}{1+x^b} dx = I_1 + I_2$, where $I_1 = \int_0^1 \frac{x^a}{1+x^b} dx$ and $I_2 = \int_1^\infty \frac{x^a}{1+x^b} dx$. We will

show that I_1 converges for $a > -1$ and I_2 converges for $b > a + 1$, so that I converges when $a > -1$ and $b > a + 1$.

I_1 is improper only when $a < 0$. When $0 \leq x \leq 1$, we have $\frac{1}{1+x^b} \leq 1 \Rightarrow \frac{1}{x^{-a}(1+x^b)} \leq \frac{1}{x^{-a}}$. The integral

$\int_0^1 \frac{1}{x^{-a}} dx$ converges for $-a < 1$ [or $a > -1$] by Exercise 57, so by the Comparison Theorem, $\int_0^1 \frac{1}{x^{-a}(1+x^b)} dx$

converges for $-1 < a < 0$. I_1 is not improper when $a \geq 0$, so it has a finite real value in that case. Therefore, I_1 has a finite real value (converges) when $a > -1$.

I_2 is always improper. When $x \geq 1$, $\frac{x^a}{1+x^b} = \frac{1}{x^{-a}(1+x^b)} = \frac{1}{x^{-a}+x^{b-a}} < \frac{1}{x^{b-a}}$. By (2), $\int_1^\infty \frac{1}{x^{b-a}} dx$ converges

for $b - a > 1$ (or $b > a + 1$), so by the Comparison Theorem, $\int_1^\infty \frac{x^a}{1+x^b} dx$ converges for $b > a + 1$.

Thus, I converges if $a > -1$ and $b > a + 1$.