

Read the whole handout [§5.0 Set Theory Transition](#). Then do the below problems.

- ▷. As in class, let $\mathbb{R}^{>0}$ be the positive real numbers (so $\mathbb{R}^{>0} = \{x \in \mathbb{R} : x > 0\}$).
- ▶. Let the universal set U be \mathbb{R} . An index set I will be $\mathbb{R}^{>0}$ or a subset of $\mathbb{R}^{>0}$. For each $r \in \mathbb{R}^{>0}$, define T_r to be the closed interval $[-r^2, r^2]$ of \mathbb{R} , i.e.,

$$T_r = \{x \in \mathbb{R} : -r^2 \leq x \leq r^2\}.$$

Find the sets listed below the dotted line. No justification needed. Your answer may be in: set builder notation, roster method, interval (closed, open, or clopen) of the real line notation, \mathbb{R} , \emptyset .
Hint. In *Thinking Land*, draw a pictorial representation for T_1, T_2, T_3, T_{17} and for some arbitrary T_r 's.

- ☺. Here is a sample. The index set is $I = \{m \in \mathbb{N} : 1 \leq m \leq 6\}$. (Note $I = \{1, 2, 3, 4, 5, 6\}$ so $\bigcap_{i \in I} T_i = \bigcap_{i=1}^6 T_i$.)

$$\bigcap_{i \in I} T_i = [-1, 1]$$

Hint. $\emptyset \neq \{0\}$

-
1. The index set is $I = \{m \in \mathbb{N} : 1 \leq m \leq 6\}$.

$$\bigcup_{i \in I} T_i = \textit{cut this out and put your solution here}$$

2. The index set is \mathbb{N} .

$$\bigcap_{i \in \mathbb{N}} T_i =$$

3. The index set is \mathbb{N} .

$$\bigcup_{i \in \mathbb{N}} T_i =$$

4. The index set is $\mathbb{R}^{>0}$.

$$\bigcap_{i \in \mathbb{R}^{>0}} T_i =$$

5. The index set is $\mathbb{R}^{>0}$.

$$\bigcup_{i \in \mathbb{R}^{>0}} T_i =$$