## Homework assigned Wednesday, February 29.

The following problems are to prepare for results that we are about to prove.

**Problem 1.** Let f(z) be analytic in a domain D and let  $a \in K$ . Let r > 0 be so small that the disk  $|z-a| \leq r$  is contained in D.

(a) Parametrize |z - a| = r by  $z = a + re^{it}$  with  $0 \le t \le 2\pi$ . Use this reparameterizing to show that

$$\int_{|z-a|=r} \frac{f(z)}{z-a} \, dz = i \int_0^{2\pi} f(a+re^{it}) \, dt$$

(b) Use part (a) to show



FIGURE 1

In the next problem we will be using Cauchy's theorem, which we now recall.

**Theorem 1** (Cauchy's Theorem). Let D be a bounded domain with nice boundary and f(z)a function that is analytic on the closure of D. Then

$$\int_{\partial D} f(z) \, dz = 0$$

where, as usual, we orient  $\partial D$  so as we move with the inside on our left.

**Problem** 2. In Figure 1 we have a bounded domain with nice boundary and a point a inside. Let f(z) be a function that is analytic on the closure of D. Let r be a small positive number and  $D_r$  the domain D with the inside of the circle |z-a| = r removed. That is  $D_r$ is the region inside of D and outside of |z - a| = r.

(a) Explain why

$$\int_{\partial D_r} \frac{f(z)}{z-a} \, dz = 0.$$

*Hint:* The function  $g(z) = \frac{f(z)}{z-a}$  is analytic in  $D_r$ . (b) The boundary of  $\partial D_r$  has two pieces. First there is the boundary,  $\partial D$ , of the original domain and second there is the circle |z - a| = r. Thus

$$\int_{\partial D_r} \frac{f(z)}{z-a} dz = \int_{\partial D} \frac{f(z)}{z-a} dz - \int_{|z-a|=r} \frac{f(z)}{z-a} dz.$$

Explain why the sign on the second integral is negative. *Hint:* We always move along the boundary with the inside on our left.

(c) Combine parts (a) and (b) to conclude

$$\int_{\partial D} \frac{f(z)}{z-a} \, dz = \int_{|z-a|=r} \frac{f(z)}{z-a} \, dz.$$

(d) In the last equation take the limit at r goes to 0 and part (b) of Problem 1 to conclude

$$\int_{\partial D} \frac{f(z)}{z-a} \, dz = 2\pi i f(a)$$

We have thus proven the following, which is maybe the most important result in complex analysis.

**Theorem 2** (Cauchy Integral Formula). Let D be a bounded domain with nice boundary and f(z) be analytic on the closure of D. Then for any point  $a \in D$ 

$$f(a) = \frac{1}{2\pi i} \int_{\partial D} \frac{f(z) \, dz}{z - a}.$$

*Example.* Let  $\gamma$  be the path in Figure 2.



FIGURE 2

We now use the Cauchy Integral formula to evaluate

$$\int_{\gamma} \frac{e^z}{z^2 + \pi^2} \, dz.$$

This function is analytic except where the denominator becomes zero. That is where  $z^2 + z^2$  $\pi^2 = 0$ . Note that  $z^2 + \pi^2 = (z - \pi i)(z + \pi i)$ . So that the bad points are  $z = \pi i$  and  $z = -\pi i$ . Thus our integral becomes

$$\int_{\gamma} \frac{e^z}{(z-\pi i)(z+\pi i)} \, dz.$$

We only need to work about the point  $\pi i$  as it is the only non-analytic point inside of  $\gamma$ . Rewrite the integral as

$$\int_{\gamma} \frac{e^z/(z+\pi i)}{(z-\pi i)} dz = \int_{\gamma} \frac{f(z)}{(z-\pi i)} dz$$
$$f(z) = -\frac{e^z}{z}.$$

where

 $\int (z)^{-} z + \pi i$ 

The function f(z) is analytic inside of  $\gamma$ . So by the Cauchy integral formula

$$\int_{\gamma} \frac{e^z}{z^2 + \pi^2} \, dz = \int_{\gamma} \frac{f(z)}{(z - \pi i)} \, dz = 2\pi i f(\pi i) = 2\pi i \frac{e^{\pi i}}{\pi i + \pi i} = e^{\pi i} = -1.$$

**Problem 3.** Let  $z_1$  be a complex number and  $\gamma$  a simple closed curve that does not pass through  $z_1$ . Show

$$\int_{\gamma} \frac{dz}{z - z_1} = \begin{cases} 2\pi i, & \text{if } z_1 \text{ is inside of } \gamma, \\ 0, & \text{if } z_1 \text{ is outside of } \gamma. \end{cases}$$

*Hint:* Use part (d) of Problem 2, or the Cauchy Integral Formula, with f(z) = 1, D the region inside of  $\gamma$ , and  $z = z_1$ .



FIGURE 3

**Problem** 4. Figure 3 shows the points i, -i, 0, and 4 along with three paths  $\alpha, \beta$ , and  $\gamma$ . Use either part (d) or Problem 2 or the Cauchy integral formula to

(a) Evaluate  $\int_{\alpha} \frac{2z+1}{z(z-4)(z^2+1)} dz,$ 

(b) Evaluate 
$$\int_{\beta} \frac{2z+1}{z(z-4)(z^2+1)} dz$$
, and

(c) Evaluate 
$$\int_{\gamma} \frac{2z+1}{z(z-4)(z^2+1)} dz$$
.