

## Homework assigned Friday, February 3.

Here we will mostly be looking at consequences of the Cauchy-Riemann equations. That is if  $f = u + iv$  in an open set, then

$$u_x = v_y \quad \text{and} \quad u_y = -v_x.$$

**Definition 1.** Let  $U$  be an open set in the complex plane  $\mathbf{C}$ . Then a function  $h: U \rightarrow \mathbf{R}$  is *harmonic* iff

$$h_{xx} + h_{yy} = 0.$$

(It is being assumed that the first and second partial derivatives of  $h$  exist and are continuous.)  $\square$

A very important result is that the real and imaginary parts of an analytic function are harmonic. To be precise

**Theorem 2.** Let  $f = u + iv$  be analytic in the open set  $U$ . Assume that  $u$  and  $v$  have continuous first and second partial derivatives. Then both  $u$  and  $v$  are harmonic.

While this is important it is not hard:

**Problem 1.** Prove the last theorem. *Hint:* It is a more or less direct consequence of the Cauchy-Riemann equations. As a start note

$$u_{xx} = (u_x)_x = (v_y)_x = v_{xy}$$

with a similar formula for  $u_{yy}$  in terms of  $v_{xy}$ . If you want more or a hint see page 88 of the text.  $\square$

Here is a variant of some of the problems we did in class today.

**Problem 2.** Let  $f = u + iv$  be analytic in a connected open set  $U$ . Assume that  $u^2 - v^2 = c$  where  $c \neq 0$  is a constant. Show  $f$  is constant. *Hint:* It is enough to show that  $u$  and  $v$  are constant. And for that it is enough to show  $u_x = u_y = 0$  and  $v_x = v_y = 0$ . Take the first partial derivatives of the equation  $u^2 - v^2 = c$  with respect to  $x$  and  $y$  and use the Cauchy-Riemann equations.  $\square$

A consequence of our proof of the Cauchy-Riemann equations is

**Proposition 3.** Let  $f = u + iv$  be analytic in an open set  $U$ . Then the derivative of  $f$  is given by either of the formulas

$$f' = u_x + iv_x \quad \text{and} \quad f' = v_y - iu_y$$

(In practice we usually just use  $f' = u_x + iv_x$ .)  $\square$

Here is an example similar to an example we did in class, if  $f(z) = e^{2z}$ , then

$$f(z) = e^{2x} \cos(2y) + ie^{2x} \sin(2y) = u + iv.$$

Thus

$$f'(z) = u_x + iv_x = 2e^{2x} \cos(2y) + i2e^{2x} \sin(2y) = 2e^{2z}$$

just as we expected.

**Problem 3.** Use Proposition 3 to show the following (which we are all familiar with for real values, but which we still need to verify for complex values.)

- (a) If  $f(z) = \cos(z)$ , then  $f'(z) = \sin(z)$ .
- (b) If  $f(z) = \sin(z)$ , then  $f'(z) = \cos(z)$ .
- (c) If  $f(z) = \log(z)$ , then  $f'(z) = \frac{1}{z}$ .

While at this point is not clear there is much relationship between analytic function and functions that can be expressed as a convergent power series, it will turn out that the two are closely related. Here is a start on that

**Proposition 4.** Consider the power series

$$f(z) = \sum_{n=0}^{\infty} a_n z^n.$$

Assume this converges for  $z = z_1$ . Then the series converges for all  $z$  with  $|z| < |z_1|$ .

**Problem 4.** Prove this. Before starting we make a few observations. As the series for  $f(z_1)$  converges, the terms go to zero. That is  $\lim_{n \rightarrow \infty} a_n z_1^n = 0$ . This implies the terms are bounded, that is there is a constant,  $C$ , so that

$$|a_n z_1^n| \leq C.$$

Define

$$r = \frac{|z|}{|z_1|} = \left| \frac{z}{z_1} \right|.$$

By hypothesis  $|z| < |z_1|$ , so

$$r < 1.$$

Thus by our basic results about geometric series

$$(1) \quad \sum_{n=0}^{\infty} C r^n < \infty$$

Now proceed with the proof as follows.

- (a) Show  $|a_n z^n| \leq C r^n$ . *Hint:*  $|a_n z^n| = |a_n z_1^n| |z/z_1|^n$ .
- (b) Finish the proof by use of the comparison theorem (look this up if you have forgotten it) part (a) and (1).

**Problem 5** (Not to be handed in). Review the definition of the gradient and the chain rule for functions of two variables. In particular that the gradient of a function is orthogonal to the level curves of the function.