

Mathematics 700 Homework due Monday October 4

The rank plus nullity theorem is one of the great (and underrated) existence theorems in elementary mathematics. Here are some examples for you to work out.

1. (From algebra.) It is a basic fact that if $p(x)$ is a polynomial of degree $\leq n$ over the field \mathbb{F} that has $(n + 1)$ roots then $p(x)$ is the zero polynomial. Use this fact to show that for any elements $z_0, \dots, z_n, w_0, \dots, w_n \in \mathbb{F}$ with z_0, \dots, z_n distinct there is a unique polynomial $p(x)$ of degree $\leq n$ so that $p(z_i) = w_i$ for $i = 0, \dots, n$. This is often stated loosely and a little imprecisely as: It is possible to assign (or interpolate) the values of a polynomial of degree $\leq n$ at $n + 1$ points arbitrarily. (Note that we have seen one proof of this, by use of Lagrange interpolation polynomials, already.) HINT: Let \mathcal{P}_n be the polynomials of degree $\leq n$ and show that the map $V: \mathcal{P}_n \rightarrow \mathbb{F}^{n+1}$ given by $V(p) := (p(z_0), \dots, p(z_n))$ is linear. Then apply the rank plus nullity theorem.
2. (From ordinary differential equations.) Let $C^2(\mathbb{R})$ be the vector space of all real valued functions on \mathbb{R} that are twice continuously differentiable. Let a and b be real numbers and let V be the two dimensional subspace of $C^2(\mathbb{R})$ spanned by the two functions $e^{ax} \cos(bx)$ and $e^{ax} \sin(bx)$. Let a_0, a_1 , and a_2 be any real numbers and define $L: C^2(\mathbb{R}) \rightarrow C^2(\mathbb{R})$ by $Ly := a_2 y'' + a_1 y' + a_0 y$. Assume that the only solution to $Ly = 0$ in V is $y = 0$. Then show that for any $f \in V$ there is a unique $y_p \in V$ so that $Ly_p = f$. COMMENT: This is basically a justification for a special case of the method of undetermined coefficients you have seen in your differential equations class. It is not much harder to justify the entire method along these same lines.
3. (From partial differential equations.) Let \mathcal{HP}_n^2 be the homogeneous polynomials of degree n in the two variables x and y with real coefficients. That is elements of \mathcal{HP}_n^2 are of the form $a_0 x^n + a_1 x^{n-1} y + \dots + a_k x^{n-k} y^k + \dots + a_n y^n$ where a_0, \dots, a_n are real numbers. A **harmonic polynomial of degree n** is an element h of \mathcal{HP}_n^2 that satisfies the partial differential equation

$$\frac{\partial^2 h}{\partial x^2} + \frac{\partial^2 h}{\partial y^2} = 0.$$

Show that there are lots of harmonic polynomials. HINT: Let \mathcal{H}_n^2 be the space of all harmonic polynomials of degree n . Define a

linear map $\Delta: \mathcal{H}\mathcal{P}_n^2 \rightarrow \mathcal{H}\mathcal{P}_{n-2}^2$ by

$$\Delta f := \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}.$$

Now what are the dimensions of $\mathcal{H}\mathcal{P}_n^2$ and $\mathcal{H}\mathcal{P}_{n-2}^2$ and what is the null space of Δ ? With just a little work you should be able to give the exact dimension of \mathcal{H}_n^2 .

4. (From algebraic geometry) We say that a curve in the plane \mathbb{R}^2 has a **polynomial parameterization** iff it is of the form

$$\begin{aligned} c(t) &= (A(t), B(t)) \\ &= (a_0 + a_1t + a_2t^2 + \cdots + a_mt^m, b_0 + b_1t + b_2t^2 + \cdots + b_nt^n) \end{aligned}$$

where $A(t)$ and $B(t)$ are polynomials as indicated. Then show that every curve with an polynomial parameterization is algebraic in the sense that there is a nonzero polynomial $p(x, y)$ in two variables so that

$$p(A(t), B(t)) \equiv 0.$$

HINT: Let \mathcal{P}_k^2 be the vector space of polynomials of degree total degree $\leq k$ in the two variables x and y . Let \mathcal{P}_d^1 be the polynomials of degree $\leq d$ in the variable t . Given the two polynomials $A(t)$ and $B(t)$ assume that $m \leq n$ and define a map $E: \mathcal{P}_k^2 \rightarrow \mathcal{P}_{nk}^1$ by

$$E(f)(t) := f(A(t), B(t))$$

and show that E is linear. Now compute dimensions of \mathcal{P}_{nk}^1 and \mathcal{P}_k^2 and let k get large. Are you able to give a bound on the degree of p in terms of the dimensions of $A(t)$ and $B(t)$?