

Mathematics 700 Homework due Wednesday, October 6

The following are problems on finding the matrices of linear maps. There are examples in Chapter 10 of the text that are relevant to these problems.

- Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be given by

$$T(x, y) := (x - 2y, -4x + y, 7x - 11y).$$

Then find the matrix of T with respect to the standard bases of each of \mathbb{R}^2 and \mathbb{R}^3 . (Recall that the standard basis of \mathbb{R}^n is the basis $e_1 = (1, 0, 0, \dots, 0)$, $e_2 = (0, 1, 0, \dots, 0)$, $e_3 = (0, 0, 1, \dots, 0)$ )

- With T as in the last problem find the matrix of T with respect to the bases $\mathcal{V} = \{(1, 2), (3, 2)\}$ of \mathbb{R}^2 and $\mathcal{W} := \{(1, 1, 1), (0, 2, 1), (0, 0, 3)\}$ of \mathbb{R}^3 .
- Letting \mathcal{P}_3 be the real polynomials of degree ≤ 3 and using the standard basis $\mathcal{V} := \{1, x, x^2, x^3\}$ of \mathcal{P}_3 find the matrices, the rank and the nullity of the following linear maps
 - $(Tp)(x) = p(x - 2)$,
 - $(Cp)(x) = (x + 1)^3 p\left(\frac{x - 1}{x + 1}\right)$,
 - $Ap = p + p' + p'' + p''' + p''''$,
 - $(Pp)(x) = e^{-x} \int_{-\infty}^x p(t)e^t dt$.
 - $(Bp)(x) = \int_0^x p'(t) dt$
 - $(Vp)(x) = \frac{d}{dx} \int_0^x p(t) dt$
- Let $M_{2 \times 2}$ be the vector space of 2×2 matrices over the real numbers. Let

$$A := \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}.$$

Use for $M_{2 \times 2}$ the the ordered basis basis $\mathcal{V} := \{E_{11}, E_{12}, E_{21}, E_{22}$ where

$$E_{11} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, E_{12} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, E_{21} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, E_{22} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

Then find the matrices of the following linear maps from $M_{2 \times 2}$ to itself.

- $LX = AX$,
- $RX = XA$,
- $CX = AX - XA$,

- (d) $TX = X^t$ (the transpose of X),
 - (e) $SX = \frac{1}{2}(X + X^t)$ and find the rank and nullity of this map,
and
 - (f) $GX = \frac{1}{2}(X - X^t)$ and find the rank and nullity of this map.
5. The set of complex numbers $\mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$ is a two dimensional vector space over the real numbers \mathbb{R} . Using the basis $\mathcal{B} = \{1, i\}$ for this real vector space find the matrices of the following linear maps
- (a) $Jz = iz$,
 - (b) $Cz = \bar{z}$ (where \bar{z} is the complex conjugate of z),
 - (c) $Tz = (2 + 3i)z$,
 - (d) $Mz = (a + bi)z$, and
 - (e) $Rz = \frac{1}{2}(z + \bar{z})$.