

---

 Math 141 HANDOUT
 

---

Fundamental Theorem of Calculus (FTC)

Let  $f: [a, b] \rightarrow \mathbb{R}$  be a continuous function.

Let  $F: [a, b] \rightarrow \mathbb{R}$  be a function.

- If  $F$  is an antiderivative of  $f$  on  $[a, b]$  (i.e.  $F'(x) = f(x)$  for each  $x \in [a, b]$ ), then

$$\int_a^b f(x) dx \equiv \int_a^b F'(x) dx = F(b) - F(a).$$

- If  $F(x) = \int_a^x f(t) dt$  for each  $x \in [a, b]$ , then  $F$  is an antiderivative of  $f$  on  $[a, b]$ , i.e.

$$F'(x) \equiv D_x \left[ \int_a^x f(t) dt \right] = f(x).$$

Basic Differentiation Rules

If the functions  $y = f(x)$  and  $y = g(x)$  are differentiable at  $x$  and  $a$  and  $b$  are constants, then:

1.  $D_x [af(x) + bg(x)] = af'(x) + bg'(x)$
2.  $D_x [f(x) \cdot g(x)] = f'(x) \cdot g(x) + f(x) \cdot g'(x)$
3.  $D_x \left[ \frac{f(x)}{g(x)} \right] = \frac{f'(x) \cdot g(x) - f(x) \cdot g'(x)}{[g(x)]^2} \quad \text{provided } g(x) \neq 0$
4.  $D_x [f(x)]^r = r [f(x)]^{r-1} f'(x) \quad \text{provided } r \in \mathbb{Q}$

If  $f$  is differentiable at  $x$  and  $g$  is differentiable at  $f(x)$ , then:

5.  $D_x [g(f(x))] = g'(f(x)) f'(x)$

Generalized Exponential and Logarithmic Functions with base  $a$  where  $a > 0$  but  $a \neq 1$

DERIVATIVES	FTC	INTEGRALS
$D_x \ln u  \stackrel{u \neq 0}{=} \frac{1}{u} \frac{du}{dx}$	$\rightleftharpoons$	$\int \frac{du}{u} \stackrel{u \neq 0}{=} \ln u  + C$
$D_x e^u = e^u \frac{du}{dx}$		$\int e^u du = e^u + C$
$D_x \log_a  u  \stackrel{u \neq 0}{=} \frac{1}{u} \frac{1}{\ln a} \frac{du}{dx}$		
$D_x a^u = a^u \ln a \frac{du}{dx}$		$\int a^u du = \frac{a^u}{\ln a} + C$

TRIG and CALCULUS

DERIVATIVES

$\xrightarrow{\text{FTC}}$

INTEGRALS

$$\begin{aligned}
 D_x \sin u &= \cos u \frac{du}{dx} & \int \cos u du &= \sin u + C \\
 D_x \tan u &= \sec^2 u \frac{du}{dx} & \int \sec^2 u du &= \tan u + C \\
 D_x \sec u &= \sec u \tan u \frac{du}{dx} & \int \sec u \tan u du &= \sec u + C \\
 D_x \cos u &= -\sin u \frac{du}{dx} & \int \sin u du &= -\cos u + C \\
 D_x \cot u &= -\csc^2 u \frac{du}{dx} & \int \csc^2 u du &= -\cot u + C \\
 D_x \csc u &= -\csc u \cot u \frac{du}{dx} & \int \csc u \cot u du &= -\csc u + C
 \end{aligned}$$

MORE INTEGRALS

$$\begin{aligned}
 \int \tan u du &= -\ln |\cos u| + C & = \ln |\sec u| + C \\
 \int \cot u du &= \ln |\sin u| + C & = -\ln |\csc u| + C \\
 \int \sec u du &= \ln |\sec u + \tan u| + C & = -\ln |\sec u - \tan u| + C \\
 \int \csc u du &= -\ln |\csc u + \cot u| + C & = \ln |\csc u - \cot u| + C
 \end{aligned}$$

DERIVATIVES

$\xrightarrow{\text{FTC}}$

INTEGRALS ( $a > 0$ )

$$\begin{aligned}
 D_x \sin^{-1} u &= \frac{1}{\sqrt{1-u^2}} \frac{du}{dx} & \int \frac{1}{\sqrt{a^2-u^2}} du &= \sin^{-1} \frac{u}{a} + C \\
 D_x \tan^{-1} u &= \frac{1}{1+u^2} \frac{du}{dx} & \int \frac{1}{a^2+u^2} du &= \frac{1}{a} \tan^{-1} \frac{u}{a} + C \\
 D_x \sec^{-1} u &= \frac{1}{|u|\sqrt{u^2-1}} \frac{du}{dx} & \int \frac{1}{u\sqrt{u^2-a^2}} du &= \frac{1}{a} \sec^{-1} \frac{|u|}{a} + C \\
 D_x \cos^{-1} u &= \frac{-1}{\sqrt{1-u^2}} \frac{du}{dx} \\
 D_x \cot^{-1} u &= \frac{-1}{1+u^2} \frac{du}{dx} \\
 D_x \csc^{-1} u &= \frac{-1}{|u|\sqrt{u^2-1}} \frac{du}{dx}
 \end{aligned}$$

Natural Logarithm Fn. $y = \ln x$	AND	Natural Exponential Fn. $y = \exp x$
-----------------------------------	-----	--------------------------------------

$$\begin{array}{lll} \ln x \stackrel{x>0}{\equiv} \int_1^x \frac{dt}{t} & & \text{nat. exp. fn. } \equiv \text{inverse of the nat. log. fn.} \\ \ln: (0, \infty) \rightarrow (-\infty, \infty) & & \exp: (-\infty, \infty) \rightarrow (0, \infty) \\ y = \ln x & \iff & x = \exp y \\ \exists \text{ a unique } e \in \mathbb{R} \text{ so that } \ln e = 1 & & e^x \stackrel{x \in \mathbb{R}}{\equiv} \exp(x) \end{array}$$

$$\begin{array}{ll} \begin{array}{c} x, y > 0 \text{ \& } r \in \mathbb{Q} \\ e^{\ln x} = x \\ \ln 1 = 0 \\ \ln(xy) = \ln x + \ln y \\ \ln\left(\frac{x}{y}\right) = \ln x - \ln y \\ \ln(x^r) = r(\ln x) \end{array} & \begin{array}{c} x, y \in \mathbb{R} \text{ \& } r \in \mathbb{Q} \\ \ln(e^x) = x \\ e^0 = 1 \\ e^x e^y = e^{x+y} \\ \frac{e^x}{e^y} = e^{x-y} \\ (e^x)^r = e^{xr} \end{array} \end{array}$$

Generalized Exponential $y = a^x$ and Logarithmic $y = \log_a x$ Functions
--

with base $a$ where $a > 0$ but $a \neq 1$ and $b > 0$ but $b \neq 1$
---

$$\log_e \equiv \ln$$

$$\begin{array}{lll} f(x) = a^x \equiv e^{x \ln a} & & : (-\infty, \infty) \rightarrow (0, \infty) \\ g(x) = \log_a x \equiv \text{the inverse of the fn. } f(x) = a^x & & : (0, \infty) \rightarrow (-\infty, \infty) \\ y = \log_a x & \iff & x = a^y \\ (\log_a b)(\log_b c) = \log_a c & \Rightarrow & \log_a x = \frac{\ln x}{\ln a} \end{array}$$

$$\begin{array}{ll} \begin{array}{c} x, y > 0 \text{ \& } r \in \mathbb{R} \\ a^{\log_a x} = x \\ \log_a 1 = 0 \\ \log_a(xy) = \log_a x + \log_a y \\ \log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y \\ \log_a(x^r) = r(\log_a x) \end{array} & \begin{array}{c} x, y \in \mathbb{R} \text{ \& } r \in \mathbb{R} \text{ \& } 0 < b \neq 1 \\ \log_a(a^x) = x \\ a^0 = 1 \\ a^x a^y = a^{x+y} \\ \frac{a^x}{a^y} = a^{x-y} \\ (a^x)^r = a^{xr} \\ (ab)^x = a^x b^x \text{ and } \left(\frac{a}{b}\right)^x = \frac{a^x}{b^x} \end{array} \end{array}$$

### Basic Trig

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} \quad \sin \theta = \frac{\text{opp}}{\text{hyp}} \quad \tan \theta = \frac{\text{opp}}{\text{adj}}$$

$$\tan x = \frac{\sin x}{\cos x} \quad \cot x = \frac{\cos x}{\sin x} \quad \sec x = \frac{1}{\cos x} \quad \csc x = \frac{1}{\sin x}$$

### Basic Inverse Trig Functions

$y = \sin \theta \Leftrightarrow \theta = \sin^{-1} y$	where	$-1 \leq y \leq 1$	and	$\frac{-\pi}{2} \leq \theta \leq \frac{\pi}{2}$
$y = \cos \theta \Leftrightarrow \theta = \cos^{-1} y$	where	$-1 \leq y \leq 1$	and	$0 \leq \theta \leq \pi$
$y = \tan \theta \Leftrightarrow \theta = \tan^{-1} y$	where	$y \in \mathbb{R}$	and	$\frac{-\pi}{2} < \theta < \frac{\pi}{2}$
$y = \cot \theta \Leftrightarrow \theta = \cot^{-1} y$	where	$y \in \mathbb{R}$	and	$0 < \theta < \pi$
$y = \sec \theta \Leftrightarrow \theta = \sec^{-1} y$	where	$ y  \geq 1$	and	$0 \leq \theta \leq \pi, \theta \neq \frac{\pi}{2}$
$y = \csc \theta \Leftrightarrow \theta = \csc^{-1} y$	where	$ y  \geq 1$	and	$\frac{-\pi}{2} \leq \theta \leq \frac{\pi}{2}, \theta \neq 0$

## Math 142

Integration by Parts:	$\int u \, dv = uv - \int v \, du$
-----------------------	------------------------------------

### Trig Identities

$$\begin{aligned}\cos(2x) &= \cos^2 x - \sin^2 x \\ \cos^2 x &= \frac{1 + \cos(2x)}{2} \\ \cos(s+t) &= \cos s \cos t - \sin s \sin t \\ \cos(s-t) &= \cos s \cos t + \sin s \sin t\end{aligned}$$

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \sin^2 x &= \frac{1 - \cos(2x)}{2} \\ \sin(s+t) &= \sin s \cos t + \cos s \sin t \\ \sin(s-t) &= \sin s \cos t - \cos s \sin t\end{aligned}$$

### Trig Substitution

IF INTEGRAND INVOLVES	THEN MAKE THE SUBSTITUTION	RESTRICTION ON $\theta$
$a^2 - u^2$	$u = a \sin \theta \rightsquigarrow \theta = \sin^{-1} \frac{u}{a}$	$\frac{-\pi}{2} \leq \theta \leq \frac{\pi}{2}$
$a^2 + u^2$	$u = a \tan \theta \rightsquigarrow \theta = \tan^{-1} \frac{u}{a}$	$\frac{-\pi}{2} < \theta < \frac{\pi}{2}$
$u^2 - a^2$	$u = a \sec \theta \rightsquigarrow \theta = \sec^{-1} \frac{u}{a}$	$0 \leq \theta \leq \pi, \theta \neq \frac{\pi}{2}$