

## MATH 122: DEFINITE INTEGRAL DOMINOES - SOLUTIONS

ANN CLIFTON  
UNIVERSITY OF SOUTH CAROLINA

Goal: To match definite integrals with their numerical or geometric representations.

$$(a) \int_0^1 (x+1)(x-1)dx = \int_0^1 (x^2-1)dx = \frac{x^3}{3} - x \Big|_0^1 = \left(\frac{1^3}{3} - 1\right) - \left(\frac{0^3}{3} - 0\right) = -\frac{2}{3}$$

$$(b) \int_0^1 (x-8x^3)dx = \frac{x^2}{2} - 2x^4 \Big|_0^1 = \left(\frac{1^2}{2} - 2(1)^4\right) - \left(\frac{0^2}{2} - 2(0)^4\right) = -\frac{3}{2}$$

$$(c) \int_0^1 \frac{x^3-x}{x+1} dx = \int_0^1 \frac{x(x^2-1)}{x+1} dx = \int_0^1 \frac{x(x+1)(x-1)}{x+1} dx = \int_0^1 x(x-1)dx = \int_0^1 (x^2-x)dx = \frac{x^3}{3} - \frac{x^2}{2} \Big|_0^1 = \left(\frac{1}{3} - \frac{1}{2}\right) - (0-0) = -\frac{1}{6}$$

$$(d) \int_0^1 e^{x+1} dx = e^{x+1} \Big|_0^1 = e^2 - e^1 = e^2 - e$$

$$(e) \int_0^1 2x^3 dx = \frac{x^4}{2} \Big|_0^1 = \frac{1}{2} - 0 = \frac{1}{2}$$

$$(f) \int_0^1 (x^2+1)dx = \frac{x^3}{3} + x \Big|_0^1 = \left(\frac{1}{3} + 1\right) - (0+0) = \frac{4}{3}$$

$$(g) \int_0^1 x^2 dx = \frac{x^3}{3} \Big|_0^1 = \frac{1}{3} - 0 = \frac{1}{3}$$

$$(h) \int_0^1 \sqrt[3]{x} dx = \int_0^1 x^{1/3} dx = \frac{x^{1/3+1}}{1/3+1} \Big|_0^1 = \frac{x^{4/3}}{4/3} \Big|_0^1 = \frac{3}{4} x^{4/3} \Big|_0^1 = \frac{3}{4}(1)^{4/3} - \frac{3}{4}(0)^{4/3} = \frac{3}{4}$$

$$(i) \int_0^1 \sqrt{x} dx = \int_0^1 x^{1/2} dx = \frac{x^{1/2+1}}{1/2+1} \Big|_0^1 = \frac{2}{3} x^{3/2} \Big|_0^1 = \frac{2}{3} (1)^{3/2} - 0 = \frac{2}{3}$$

$$(j) \int_0^1 \pi dx = \pi x \Big|_0^1 = \pi(1) - \pi(0) = \pi$$

$$(k) \int_0^1 (x+1)^2 dx = \int_0^1 (x^2 + 2x + 1) dx = \frac{x^3}{3} + x^2 + x \Big|_0^1 = \left(\frac{1^3}{3} + (1)^2 + 1\right) - (0) = \frac{7}{3}$$

$$(l) \int_0^1 (x+1)^3 dx = \int_0^1 (x+1)(x^2 + 2x + 1) dx = \int_0^1 (x^3 + 3x^2 + 3x + 1) dx = \frac{x^4}{4} + x^3 + \frac{3x^2}{2} + x \Big|_0^1 = \left(\frac{1^4}{4} + 1 + \frac{3}{2} + 1\right) - (0) = \frac{15}{4}$$