Math 580/780I Review Sheet for Test 2, Fall 2019

Stuff to Know:

- The Chinese Remainder Theorem (Theorem 14)!!
- The Euclidean algorithm for polynomials.
- How to find u(x) and v(x) such that $f(x)u(x) + g(x)v(x) = \gcd(f(x), g(x))$.
- The elementary symmetric functions.
- The Remainder Theorem.
- Arithmetic modulo polynomials.
- The proof of Theorem 16.
- Know Lagrange's Theorem (Theorem 18).
- How to compute orders of integers modulo some integer n > 0.
- How to determine if a number is a primitive root modulo a prime (or integer).
- How to do the quiz problems and the review problems below.
- How to do the old quiz and old test problems.
- How to do homework problems and examples from class. This includes items not mentioned above. See class presentations for examples.

Practice Problems:

- 1. What is the remainder when 1234567891011...4647 is divided by 30?
- 2. Solve the system of congruences below. (Hint: $2^3 \cdot 3 \cdot 5 \cdot 19 = 2280$.)

$$x \equiv 10 \pmod{24}$$

 $x \equiv 18 \pmod{20}$
 $x \equiv 13 \pmod{15}$
 $x \equiv 3 \pmod{95}$

- 3. What are the first two positive integers n that satisfy 8|n, 9|(n + 1), 10|(n + 2), 11|(n + 3) and 12|(n + 4)?
- 4. What is the remainder when 3^{100} is divided by 45?
- 5. What is the remainder when 2^{200} is divided by 216? Note that 2 and 216 are not relatively prime.

6. If α_1 , α_2 and α_3 are the roots of $x^3 + 2x^2 - 3x + 5$, then what are the values of

$$\alpha_1+\alpha_2+\alpha_3, \qquad \alpha_1^2+\alpha_2^2+\alpha_3^2, \qquad \text{and} \qquad \alpha_1^3+\alpha_2^3+\alpha_3^3\,?$$

7. What is the remainder when we divide

$$x^{2010} + 2x^{2005} + 4x^{1020} - 16x^{1010} + x^{24} - 2x^{12} + x^7 - x^3 + 1$$
 by $x^5 + 2$?

- 8. Let p be a prime $\equiv 5 \pmod 8$, so p = 8k + 5 for some integer k. Explain why $x^4 + 1 \equiv 0 \pmod p$ has no solutions.
- 9. (a) State Lagrange's Theorem.
 - (b) If p is a prime $\equiv 1 \mod 0$, then it is known that each cube $x \mod p$ satisfies the congruence $x^{(p-1)/3} \equiv 1 \pmod p$. What does Lagrange's Theorem imply about the number of cubes modulo such a prime?
- 10. (a) Which of the following can be the order of an integer a modulo 50? There may be more than one correct answer. Find all of them.

- (b) The number 3 is a primitive root modulo 50. Find an integer $a \in \{1, 2, ..., 49\}$ that has order 5 modulo 50. If you didn't use that 3 is a primitive root modulo 50, figure out how to.
- 11. Is 2 a primitive root modulo 23? How about modulo 29?
- 12. The following are consecutive powers of 5 modulo 37 that your professor has kindly provided for you:

$$5^{24} \equiv 26 \pmod{37}, \quad 5^{25} \equiv 19 \pmod{37}, \quad 5^{26} \equiv 21 \pmod{37},$$

 $5^{27} \equiv 31 \pmod{37}, \quad 5^{28} \equiv 7 \pmod{37}, \quad 5^{29} \equiv 35 \pmod{37},$
 $5^{30} \equiv 27 \pmod{37}, \quad 5^{31} \equiv 24 \pmod{37}, \quad 5^{32} \equiv 9 \pmod{37}.$

In particular, note that $5^{31} \equiv 24 \pmod{37}$. There is an integer $k \in \{1, 2, 3, \dots, 36\}$ satisfying $k^7 \equiv 24 \pmod{37}$. Find that value of k. (Hint: By Fermat's Little Theorem, $5^{36} \equiv 1 \pmod{37}$ so $5^{67} \equiv 5^{31} \equiv 24 \pmod{37}$. Find the right power of 5 that is congruent to 24 modulo 37.)