

Level 1:

Level 1: If $\vec{u} = \langle 2, -1, -2 \rangle$, then what is the magnitude (or length) of \vec{u} ?

Level 1: If $\vec{u} = \langle 2, -1, -2 \rangle$, then what is the magnitude (or length) of \vec{u} ?

(a) 2

(b) 3

(c) 6

(d) 9

Level 1: If $\vec{u} = \langle 2, -1, -2 \rangle$, then what is the magnitude (or length) of \vec{u} ?

(a) 2

(b) 3

(c) 6

(d) 9

IS THAT YOUR FINAL ANSWER?

Level 1: If $\vec{u} = \langle 2, -1, -2 \rangle$, then what is the magnitude (or length) of \vec{u} ?

(a) 2

(b) 3

(c) 6

(d) 9

Level 2:

Level 2: What is the value of $\int_0^\pi \int_0^{\pi-x} dy dx$?

Level 2: What is the value of $\int_0^\pi \int_0^{\pi-x} dy dx$?

(a) $\pi^2/2$

(b) $\pi^2/3$

(c) 1

(d) 0

Level 2: What is the value of $\int_0^\pi \int_0^{\pi-x} dy dx$?

(a) $\pi^2/2$

(b) $\pi^2/3$

(c) 1

(d) 0

IS THAT YOUR FINAL ANSWER?

Level 2: What is the value of $\int_0^\pi \int_0^{\pi-x} dy dx$?

(a) $\pi^2/2$

(b) $\pi^2/3$

(c) 1

(d) 0

Level 3:

Level 3: Let $\vec{u} = \langle 1, -2, 2 \rangle$ and $\vec{v} = \langle -1, 0, -1 \rangle$.
What is the smallest angle θ between these two vectors?

Level 3: Let $\vec{u} = \langle 1, -2, 2 \rangle$ and $\vec{v} = \langle -1, 0, -1 \rangle$.
What is the smallest angle θ between these two vectors?

- (a) $3\pi/4$ (b) $2\pi/3$ (c) $5\pi/3$ (d) $\pi/2$

Level 3: Let $\vec{u} = \langle 1, -2, 2 \rangle$ and $\vec{v} = \langle -1, 0, -1 \rangle$.
What is the smallest angle θ between these two vectors?

- (a) $3\pi/4$ (b) $2\pi/3$ (c) $5\pi/3$ (d) $\pi/2$

IS THAT YOUR FINAL ANSWER?

Level 3: Let $\vec{u} = \langle 1, -2, 2 \rangle$ and $\vec{v} = \langle -1, 0, -1 \rangle$.
What is the smallest angle θ between these two vectors?

- (a) $3\pi/4$ (b) $2\pi/3$ (c) $5\pi/3$ (d) $\pi/2$

Level 4:

Level 4: Which of the following is an equation for the tangent plane to the surface $z = x^2 - 2y^2$ at the point $(3, 2, 1)$?

Level 4: Which of the following is an equation for the tangent plane to the surface $z = x^2 - 2y^2$ at the point $(3, 2, 1)$?

(a) $2x - 4y = 1$

(b) $2x - 4y - z = -3$

(c) $6x - 8y - z = 3$

(d) $6x - 8y - z = 1$

Level 4: Which of the following is an equation for the tangent plane to the surface $z = x^2 - 2y^2$ at the point $(3, 2, 1)$?

- (a) $2x - 4y = 1$ (b) $2x - 4y - z = -3$
(c) $6x - 8y - z = 3$ (d) $6x - 8y - z = 1$

IS THAT YOUR FINAL ANSWER?

Level 4: Which of the following is an equation for the tangent plane to the surface $z = x^2 - 2y^2$ at the point $(3, 2, 1)$?

(a) $2x - 4y = 1$

(b) $2x - 4y - z = -3$

(c) $6x - 8y - z = 3$

(d) $6x - 8y - z = 1$

Level 5:

Level 5: If the Cartesian coordinates (rectangular coordinates) of a point are $(0, \sqrt{3}, -1)$, then what is the corresponding angle ϕ in spherical coordinates?

Level 5: If the Cartesian coordinates (rectangular coordinates) of a point are $(0, \sqrt{3}, -1)$, then what is the corresponding angle ϕ in spherical coordinates?

(a) $\pi/6$

(b) $\pi/3$

(c) $2\pi/3$

(d) $5\pi/6$

Level 5: If the Cartesian coordinates (rectangular coordinates) of a point are $(0, \sqrt{3}, -1)$, then what is the corresponding angle ϕ in spherical coordinates?

(a) $\pi/6$

(b) $\pi/3$

(c) $2\pi/3$

(d) $5\pi/6$

IS THAT YOUR FINAL ANSWER?

Level 5: If the Cartesian coordinates (rectangular coordinates) of a point are $(0, \sqrt{3}, -1)$, then what is the corresponding angle ϕ in spherical coordinates?

(a) $\pi/6$

(b) $\pi/3$

(c) $2\pi/3$

(d) $5\pi/6$

Level 6:

Level 6: For the function $f(x, y) = 3x^2y + 3x^2 - y^3$,
the point $(1, -1)$ determines . . .

Level 6: For the function $f(x, y) = 3x^2y + 3x^2 - y^3$, the point $(1, -1)$ determines . . .

(a) a local maximum

(b) a local minimum

(c) a saddle point

(d) not a critical point

Level 6: For the function $f(x, y) = 3x^2y + 3x^2 - y^3$, the point $(1, -1)$ determines . . .

- (a) a local maximum
- (b) a local minimum
- (c) a saddle point
- (d) not a critical point

IS THAT YOUR FINAL ANSWER?

Level 6: For the function $f(x, y) = 3x^2y + 3x^2 - y^3$, the point $(1, -1)$ determines . . .

(a) a local maximum

(b) a local minimum

(c) a saddle point

(d) not a critical point

Level 7:

Level 7: Which of the following are parametric equations for the line passing through $(11, 3, 8)$ and $(2, 9, 6)$?

Level 7: Which of the following are parametric equations for the line passing through $(11, 3, 8)$ and $(2, 9, 6)$?

(a) $x = 2 - 9t$
 $y = 9 + 6t$
 $z = 8 - 2t$

(b) $x = 24 + 18t$
 $y = 12 - 12t$
 $z = 8 + 4t$

(c) $x = 20 + 18t$
 $y = -3 - 12t$
 $z = 10 + 4t$

(d) $x = 2 + 9t$
 $y = 9 - 6t$
 $z = 10 + 2t$

Level 7: Which of the following are parametric equations for the line passing through $(11, 3, 8)$ and $(2, 9, 6)$?

(a) $x = 2 - 9t$
 $y = 9 + 6t$
 $z = 8 - 2t$

(b) $x = 24 + 18t$
 $y = 12 - 12t$
 $z = 8 + 4t$

(c) $x = 20 + 18t$
 $y = -3 - 12t$
 $z = 10 + 4t$

(d) $x = 2 + 9t$
 $y = 9 - 6t$
 $z = 10 + 2t$

IS THAT YOUR FINAL ANSWER?

Level 7: Which of the following are parametric equations for the line passing through $(11, 3, 8)$ and $(2, 9, 6)$?

(a) $x = 2 - 9t$
 $y = 9 + 6t$
 $z = 8 - 2t$

(b) $x = 24 + 18t$
 $y = 12 - 12t$
 $z = 8 + 4t$

(c) $x = 20 + 18t$
 $y = -3 - 12t$
 $z = 10 + 4t$

(d) $x = 2 + 9t$
 $y = 9 - 6t$
 $z = 10 + 2t$

Level 8:

Level 8: What is the value of $\int_0^\pi \int_0^{\pi-x} \frac{\sin y}{\pi-y} dy dx$?

Level 8: What is the value of $\int_0^\pi \int_0^{\pi-x} \frac{\sin y}{\pi-y} dy dx$?

(a) **3**

(b) **2**

(c) **1**

(d) **0**

Level 8: What is the value of $\int_0^\pi \int_0^{\pi-x} \frac{\sin y}{\pi-y} dy dx$?

(a) **3**

(b) **2**

(c) **1**

(d) **0**

IS THAT YOUR FINAL ANSWER?

Level 8: What is the value of $\int_0^\pi \int_0^{\pi-x} \frac{\sin y}{\pi-y} dy dx$?

(a) 3

(b) 2

(c) 1

(d) 0

Level 9:

Level 9: Which of the following is a point P on the graph of $x^2 - y^2 = z^2 - 1$ with the tangent plane at P parallel to the xz -plane?

Level 9: Which of the following is a point P on the graph of $x^2 - y^2 = z^2 - 1$ with the tangent plane at P parallel to the xz -plane?

(a) $(0, -1, 0)$

(b) $(2, 1, 2)$

(c) $(1, 1, -1)$

(d) $(0, 0, 1)$

Level 9: Which of the following is a point P on the graph of $x^2 - y^2 = z^2 - 1$ with the tangent plane at P parallel to the xz -plane?

(a) $(0, -1, 0)$

(b) $(2, 1, 2)$

(c) $(1, 1, -1)$

(d) $(0, 0, 1)$

IS THAT YOUR FINAL ANSWER?

Level 9: Which of the following is a point P on the graph of $x^2 - y^2 = z^2 - 1$ with the tangent plane at P parallel to the xz -plane?

(a) $(0, -1, 0)$

(b) $(2, 1, 2)$

(c) $(1, 1, -1)$

(d) $(0, 0, 1)$

Level 10:

Level 10: Calculate $\iiint_S (x^2 + y^2)^{1/2} dV$ where S is the solid between the paraboloids $z = x^2 + y^2 - 1$ and $z = -2x^2 - 2y^2 + 8$.

Level 10: Calculate $\iiint_S (x^2 + y^2)^{1/2} dV$ where S is the solid between the paraboloids $z = x^2 + y^2 - 1$ and $z = -2x^2 - 2y^2 + 8$.

- (a) $81\pi/2$ (b) $36\pi\sqrt{3}/5$ (c) $162\pi/5$ (d) $81\pi/5$

Level 10: Calculate $\iiint_S (x^2 + y^2)^{1/2} dV$ where S is the solid between the paraboloids $z = x^2 + y^2 - 1$ and $z = -2x^2 - 2y^2 + 8$.

- (a) $81\pi/2$ (b) $36\pi\sqrt{3}/5$ (c) $162\pi/5$ (d) $81\pi/5$

IS THAT YOUR FINAL ANSWER?

Level 10: Calculate $\iiint_S (x^2 + y^2)^{1/2} dV$ where S is the solid between the paraboloids $z = x^2 + y^2 - 1$ and $z = -2x^2 - 2y^2 + 8$.

- (a) $81\pi/2$ (b) $36\pi\sqrt{3}/5$ (c) $162\pi/5$ (d) $81\pi/5$

Level 11:

Level 11: A drop of water is placed gently onto the surface of $z = 3y^2 - 4x^2$ at the point $(1, 1, -1)$. In what direction does the drop begin to move? Express your answer as a unit vector $\langle \mathbf{a}, \mathbf{b} \rangle$ (so the raindrop will go in this direction along the surface).

Level 11: A drop of water is placed gently onto the surface of $z = 3y^2 - 4x^2$ at the point $(1, 1, -1)$. In what direction does the drop begin to move? Express your answer as a unit vector $\langle \mathbf{a}, \mathbf{b} \rangle$ (so the raindrop will go in this direction along the surface).

(a) $\langle 4/5, -3/5 \rangle$

(b) $\langle -4/5, -3/5 \rangle$

(c) $\langle -4/5, 3/5 \rangle$

(d) $\langle 4/5, 3/5 \rangle$

Level 11: A drop of water is placed gently onto the surface of $z = 3y^2 - 4x^2$ at the point $(1, 1, -1)$. In what direction does the drop begin to move? Express your answer as a unit vector $\langle \mathbf{a}, \mathbf{b} \rangle$ (so the raindrop will go in this direction along the surface).

(a) $\langle 4/5, -3/5 \rangle$

(b) $\langle -4/5, -3/5 \rangle$

(c) $\langle -4/5, 3/5 \rangle$

(d) $\langle 4/5, 3/5 \rangle$

IS THAT YOUR FINAL ANSWER?

Level 11: A drop of water is placed gently onto the surface of $z = 3y^2 - 4x^2$ at the point $(1, 1, -1)$. In what direction does the drop begin to move? Express your answer as a unit vector $\langle \mathbf{a}, \mathbf{b} \rangle$ (so the raindrop will go in this direction along the surface).

(a) $\langle 4/5, -3/5 \rangle$

(b) $\langle -4/5, -3/5 \rangle$

(c) $\langle -4/5, 3/5 \rangle$

(d) $\langle 4/5, 3/5 \rangle$

Level 12:

Level 12: What is the value of

$$\lim_{h \rightarrow 0} \frac{e^{x(y+h)} - e^{xy}}{h} ?$$

Level 12: What is the value of

$$\lim_{h \rightarrow 0} \frac{e^{x(y+h)} - e^{xy}}{h} ?$$

- (a) e^{xy} (b) xe^{xy} (c) ye^{xy} (d) does not exist

Level 12: What is the value of

$$\lim_{h \rightarrow 0} \frac{e^{x(y+h)} - e^{xy}}{h} ?$$

- (a) e^{xy} (b) xe^{xy} (c) ye^{xy} (d) does not exist

IS THAT YOUR FINAL ANSWER?

Level 12: What is the value of

$$\lim_{h \rightarrow 0} \frac{e^{x(y+h)} - e^{xy}}{h} ?$$

- (a) e^{xy} (b) xe^{xy} (c) ye^{xy} (d) does not exist

Level 13:

Level 13: What is the value of $\iiint_S (x^2 + y^2)^{3/2} dV$

where S is the solid inside the cylinder $x^2 + y^2 = 1$, below the cone $z = 4 - \sqrt{x^2 + y^2}$ and above the plane $z = 0$?

Level 13: What is the value of $\iiint_S (x^2 + y^2)^{3/2} dV$

where S is the solid inside the cylinder $x^2 + y^2 = 1$, below the cone $z = 4 - \sqrt{x^2 + y^2}$ and above the plane $z = 0$?

- (a) $8\pi/5$ (b) $9\pi/10$ (c) $10\pi/3$ (d) $19\pi/15$

Level 13: What is the value of $\iiint_S (x^2 + y^2)^{3/2} dV$

where S is the solid inside the cylinder $x^2 + y^2 = 1$, below the cone $z = 4 - \sqrt{x^2 + y^2}$ and above the plane $z = 0$?

- (a) $8\pi/5$ (b) $9\pi/10$ (c) $10\pi/3$ (d) $19\pi/15$

IS THAT YOUR FINAL ANSWER?

Level 13: What is the value of $\iiint_S (x^2 + y^2)^{3/2} dV$

where S is the solid inside the cylinder $x^2 + y^2 = 1$, below the cone $z = 4 - \sqrt{x^2 + y^2}$ and above the plane $z = 0$?

- (a) $8\pi/5$ (b) $9\pi/10$ (c) $10\pi/3$ (d) $19\pi/15$

Level 14:

Level 14: What is the shortest distance from a point on the graph of $z = x^2 - 3y^2 - 1$ to the origin $(0, 0, 0)$?

Level 14: What is the shortest distance from a point on the graph of $z = x^2 - 3y^2 - 1$ to the origin $(0, 0, 0)$?

(a) $1/2$

(b) $\sqrt{2}/2$

(c) $\sqrt{3}/2$

(d) 1

Level 14: What is the shortest distance from a point on the graph of $z = x^2 - 3y^2 - 1$ to the origin $(0, 0, 0)$?

(a) $1/2$

(b) $\sqrt{2}/2$

(c) $\sqrt{3}/2$

(d) 1

IS THAT YOUR FINAL ANSWER?

Level 14: What is the shortest distance from a point on the graph of $z = x^2 - 3y^2 - 1$ to the origin $(0, 0, 0)$?

(a) $1/2$

(b) $\sqrt{2}/2$

(c) $\sqrt{3}/2$

(d) 1

Level 15:

Level 15: What is the value of

$$\int_0^\pi \int_0^2 \int_0^1 zx^2 \sin(xyz) \, dx \, dy \, dz ?$$

Level 15: What is the value of

$$\int_0^\pi \int_0^2 \int_0^1 zx^2 \sin(xyz) \, dx \, dy \, dz ?$$

- (a) $\pi/4$ (b) $\pi/3$ (c) $\pi/2$
(d) none of the above

Level 15: What is the value of

$$\int_0^{\pi} \int_0^2 \int_0^1 zx^2 \sin(xyz) \, dx \, dy \, dz ?$$

- (a) $\pi/4$ (b) $\pi/3$ (c) $\pi/2$
(d) none of the above

IS THAT YOUR FINAL ANSWER?

Level 15: What is the value of

$$\int_0^\pi \int_0^2 \int_0^1 zx^2 \sin(xyz) \, dx \, dy \, dz ?$$

- (a) $\pi/4$ (b) $\pi/3$ (c) $\pi/2$
(d) none of the above