

# A Distribution Problem for Powerfree Values of Irreducible Polynomials

by B. Beasley and M. Filaseta

**Theorem 2.** *Let  $k \geq (\sqrt{2} - 1/2)g$ . Given any  $\gamma > 0$ , there exists a  $\delta = \delta(\gamma) > 0$  such that*

$$\sum_{\substack{s_{n+1} \leq X \\ s_{n+1} - s_n \leq X^\delta}} (s_{n+1} - s_n)^\gamma \sim B(\gamma, f, k)X,$$

for some constant  $B(\gamma, f, k)$  depending only on  $\gamma$ ,  $f(x)$ , and  $k$ .

**Definition 1.** Given  $f(x) \in \mathbb{Z}[x]$ , let  $s_n = s_n(f)$  be the  $n$ th positive integer  $m$  such that  $f(m)$  is  $k$ -free. Let

$$L(h) = L(h, X) = L(h, X, f) = |\{n \in \mathbb{Z}^+ : h < s_{n+1} - s_n \leq 2h, X/2 < s_{n+1} \leq X\}|.$$

$$(2) \quad L(2h) \ll \frac{X}{h^{\gamma+\varepsilon}}$$

**Comment:** (2) & Theorem 3  $\implies$  Theorems 1 & 2 (sort-of).

**Notation:**  $D = R(f, f')$ , the resultant of  $f$  and  $f'$

$$z = 4g$$

$$\mathcal{A} = \{p : p|D \text{ or } p \leq z\}$$

$$Q = \prod_{p \in \mathcal{A}} p^k$$

$$H = \frac{h \log h}{8gQ}$$

$$\rho(q) = \rho(f, q) = |\{a \in \mathbb{Z}_q : f(a) \equiv 0 \pmod{q}\}|$$

$$T \in \mathcal{T} = \{2^j H : j = 0, 1, \dots, J\}$$

**Lemma 4.** *Let*

$$F(n) = F(n, h, f) = \sum_{n < m \leq n+h} \sum_{\substack{p > H \\ p^k | f(m)}} 1.$$

*If there are no integers  $m$  in  $(n, n + h]$  such that  $f(m)$  is  $k$ -free, then*

$$F(n) \geq \frac{h}{4Q}.$$

**Lemma 5.** For  $r$  a positive integer, define

$$M_r = M_r(h, X, f) = \sum_{X/2 < n \leq X} F^r(n).$$

Then

$$L(2h) \ll_r \frac{M_r}{h^{r+1}} + 1.$$

**Definition 2.** Let  $S_r(T)$  be the number of  $2r$ -tuples  $(p_1, \dots, p_r, m_1, \dots, m_r)$  with  $T < p_1 < p_2 < \dots < p_r \leq 2T$  such that, for each  $j \in \{1, 2, \dots, r\}$ ,

$$|m_1 - m_j| < h, \quad \frac{X}{2} < m_j \leq X + h \leq 2X, \quad \text{and} \quad p_j^k | f(m_j).$$

**Notation:**  $G(n, T) = G(n, T, f) = \sum_{n < m \leq n+h} \sum_{\substack{p^k | f(m) \\ T < p \leq 2T}} 1$

**Lemma 6.** Let  $r$  be a positive integer. If  $r = 1$ , then

$$\sum_{X/2 < n \leq X} \binom{G(n, T)}{r} \leq (2g)^r h S_r(T).$$

If  $r > 1$ , then

$$\sum_{\substack{X/2 < n \leq X \\ G(n, T) \geq 2gr}} \binom{G(n, T)}{r} \leq (2g)^r h S_r(T).$$

**Lemma 7.** If  $T \geq H$  and  $r \in \mathbb{Z}^+$ , then

$$S_r(T) \ll \frac{h^{r-1} X}{T^{(k-1)r} \log^r T} + \frac{h^{r-1} T^r}{\log^r T}.$$

**Lemma 8.** For  $\mathcal{T}$  as previously defined,

$$\sum_{\substack{T \in \mathcal{T} \\ U < T \leq V}} T^a \ll \begin{cases} U^a & \text{if } a < 0 \\ V^a & \text{if } a > 0. \end{cases}$$

**Lemma 9.** Let  $k \geq (\sqrt{2} - 1/2)g$ . Let  $\varepsilon \in (0, (k-1)/k]$  such that

$$\varepsilon < 1 - \frac{8g(g-1)}{(2k+g)^2 - 4}.$$

If  $X^{1-\varepsilon} < T \ll X^{g/k}$ , then there is a  $\xi = \xi(\varepsilon) > 0$  such that

$$S_1(T) \ll X^{1-\xi}.$$