

MARK BOX		
PROBLEM	POINTS	
0A	11	
0B	8	
0C	8	
0D	3	
<hr/>		
Total for 0	30	
1	10	
2	10	
3	10	
4-13	40=10x4	
%	100	

NAME: _____

PIN: _____

INSTRUCTIONS

- Instructions for the different types of problems on this exam.
 - (1) Problem 0.
 - Fill-in the blank/box or circle the correct answer (can circle at most one choice).
 - No need to show work.
 - (2) Multiple Choice.
 - Circle the correct answer. Circle at most one choice.
 - No need to show work (but can do scratch work if needed). No partial credit.
 - (3) Work Out.
 - Partial credit possible.
 - Justify your answer below the provided box/line.
 - * Show all your work. Work in a logical fashion. Use proper notation.
 - * Explain your thoughts (using English words).
 - * You will be graded on the quality and correctness of your justification.
 - Then put your answer in/on the provided box/line.
 - No points for a correct answer that does not have a justification.
- The MARK BOX above indicates the problems (check you have them all) along with their points.
- Write your PIN on the top center on **each page**.
- **Write on only the front side of the paper.** Only the front side will graded.
- If you run out of space for a solution at the place the solution should go, then leave me a note at the where you solution should be telling me where to go look for the rest of your solution.
- During the exam, the use of unauthorized materials is prohibited. Unauthorized materials include: books, notes, electronic devices (e.g., cell phone, smart watch, earbuds). Unauthorized materials (including cell phones) must be in a secured (e.g. zipped up, snapped closed) bag placed completely under your desk or, if you did not bring such a bag, then place on the front desk during the exam (and pick it after the exam). **So no electronic devices allowed in your pockets.**
- At a student's request, a clock will be projected to the screen.
- During this exam, do not leave your seat without permission.
- If you have a question, then raise your hand.
- When you finish: turn your exam over, put your pencil down and raise your hand.
- No portion, nor the statements of the problems, may leave the classroom.
- Not following instructions can result in a lose of points. Cheating is grounds for a F in the course.
- This exam covers (from *Calculus* by Thomas, 13th ed., ET): §10.7–10.10, 11.1–11.4.

Honor Code Statement

I understand that it is the responsibility of every member of the Carolina community to uphold and maintain the University of South Carolina's Honor Code.

As a Carolinian, I certify that I have neither given nor received unauthorized aid on this exam.

I understand that if it is determined that I used any unauthorized assistance or otherwise violated the University's Honor Code then I will receive a failing grade for this course and be referred to the academic Dean and the Office of Academic Integrity for additional disciplinary actions.

Furthermore, I have not only read but will also follow the instructions on the exam.

Signature : _____

<p>The rest of this page is left intentionally blank. You may use the space as scratch paper. Nothing written below will be looked at nor graded.</p>

0B. Taylor/Maclaurin Polynomials and Series.

For this part, fill-in the 9 boxes.

Let $y = f(x)$ be a function with derivatives of all orders in an interval I containing x_0 .

Let $y = P_N(x)$ be the N^{th} -order Taylor polynomial of $y = f(x)$ about x_0 .

Let $y = R_N(x)$ be the N^{th} -order Taylor remainder of $y = f(x)$ about x_0 .

Let $y = P_\infty(x)$ be the Taylor series of $y = f(x)$ about x_0 .

Let c_n be the n^{th} Taylor coefficient of $y = f(x)$ about x_0 .

- a. The formula for c_n is

$c_n =$

- b. In open form (i.e., with ... and without a \sum -sign)

$P_N(x) =$

- c. In closed form (i.e., with a \sum -sign and without ...)

$P_N(x) =$

- d. In open form (i.e., with ... and without a \sum -sign)

$P_\infty(x) =$

- e. In closed form (i.e., with a \sum -sign and without ...)

$P_\infty(x) =$

- f. We know that $f(x) = P_N(x) + R_N(x)$. Taylor's BIG Theorem tells us that, for each $x \in I$,

$R_N(x) =$

for some c

- g. A Maclaurin series is a Taylor series with the center specifically specified as $x_0 =$

0C. Commonly Used Taylor Series

Fill in the blank boxes with the choices a – ℓ, which are provided below.

You may use a choice more than once or not at all.

Sample questions, which are needed later in the exam, are already done for you.

a. $\sum_{n=0}^{\infty} x^n$

d. $\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$

g. $x \in \mathbb{R}$

j. $(-1, 1]$

b. $\sum_{n=0}^{\infty} \frac{x^n}{n!}$

e. $\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$

h. $(-1, 1)$

k. $[-1, 1)$

c. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n}$

f. $\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$

i. $[-1, 1]$

ℓ. none of the others

sample. A power series expansion for $y = \frac{1}{1-x}$ is and is valid precisely when .

sample. A power series expansion for $y = e^x$ is and is valid precisely when .

o.1. A power series expansion for $y = \cos x$ is and is valid precisely when .

o.2. A power series expansion for $y = \sin x$ is and is valid precisely when .

o.3. A power series expansion for $y = \ln(1+x)$ is and is valid precisely when .

o.4. A power series expansion for $y = \tan^{-1} x$ is and is valid precisely when .

0D. Parametric Curves In this part, fill in the 3 boxes. Consider the curve \mathcal{C} parameterized by

$$x = x(t)$$

$$y = y(t)$$

for $a \leq t \leq b$.

1) Express $\frac{dy}{dx}$ in terms of derivatives with respect to t . Answer: $\frac{dy}{dx} =$

2) The tangent line to \mathcal{C} when $t = t_0$ is $y = mx + b$ where m is evaluated at $t = t_0$.

3) The arc length of \mathcal{C} , expressed as an integral with respect to t , is


Arc Length =

Work Out: problems 1–3.

1. Consider the formal power series

$$\sum_{n=1}^{\infty} \frac{(2x+8)^n}{n}.$$

As we did in class, in the box below draw a diagram indicating for which x 's this series is: absolutely convergent, conditionally convergent, and divergent. Of course, indicate your reasoning. Don't forget to check the endpoints, if there are any.



2. Hint: $e^{x^2} = e^{(x^2)}$.
- 2.1. Using a *Commonly Used Taylor Series* (see problem **0C**), find a power series representation, centered about $x_0 = 0$, for the function

$$f(x) = e^{x^2} \quad (2.1)$$

and say when it is valid.

Express your series in CLOSED form (i.e., with a \sum -sign and without ...).

Soln: $e^{x^2} =$

, valid when

- 2.2. Using your solution for first part of this problem, find a power series representation, centered about $x_0 = 0$, for

$$\int e^{x^2} dx. \quad (2.2)$$

Express your series in CLOSED form (i.e., with a \sum -sign and without ...).

Note: we cannot express $\int e^{x^2} dx$ as an elementary function (loosely speaking, you cannot integrate e^{x^2}).

Soln: $\int e^{x^2} dx =$

 + C

3. Hint: $\sqrt{e} = e^{\frac{1}{2}}$.
3.1. Using a *Commonly Used Taylor Series* (see problem 0C), express the number

$$\sqrt{e} \tag{3.1}$$

as a numerical series. Express your series in CLOSED form (i.e., with a \sum -sign and without ...).

Soln: $\sqrt{e} =$

- 3.2. In the first part of this problem, you found a_n 's so that $\sqrt{e} = \sum_{n=0}^{\infty} a_n$. Now estimate the error in approximating \sqrt{e} by the partial sum $\sum_{n=0}^2 a_n$ of your infinite series $\sum_{n=0}^{\infty} a_n$ in Part 3.1.

answer: $\left| \sqrt{e} - \sum_{n=0}^2 a_n \right| \leq$

Multiple Choice: problems 4-13.

4. Let the function $y = f(x)$ have a power series power series representation $\sum_{n=0}^{\infty} c_n x^n$, which is valid in some interval $(-R, R)$ where $R > 0$.
- Then $f(0)$ must be 0.
 - Then $f(0)$ must be c_0 .
 - Then $f(0)$ must be c_1 .
 - Then we know that $f(0)$ exists but we do not know what the value of $f(0)$ is.
 - None of the others.
5. Let the function $y = f(x)$ have a power series power series representation $\sum_{n=0}^{\infty} a_n x^n$, which is valid in some interval J containing 0 and the radius of J strictly positive. Consider the two statements:
- (A) If $y = f(x)$ is an even function (i.e., $f(-x) = f(x)$), then $a_1 = a_3 = a_5 = \dots = 0$.
- (B) If $y = f(x)$ is an odd function (i.e., $f(-x) = -f(x)$), then $a_0 = a_2 = a_4 = \dots = 0$.
- Both (A) and (B) are true.
 - Both (A) and (B) are false.
 - (A) is true but (B) is false.
 - (A) is false but (B) is true.
 - None of the others.

►. **Problems 4 and 5 were meant to help you with Problem 0C. ☺☺☺**

6. Find the interval of convergence of the power series

$$\sum_{n=1}^{\infty} \frac{(x-2)^n}{10^n}$$

Recall that the interval of convergence is the set of x 's for which the power series converges, either absolutely or conditionally.

- $(-10, 10)$
 - $[-10, 10]$
 - $(-8, 12)$
 - $[-8, 12]$
 - None of the others.
7. Find the 3rd order Taylor polynomial, about the center $x_0 = 1$, for the function $f(x) = x^5 - x^2 + 5$.
- $p_3(x) = 5 + 3(x-1) + 9(x-1)^2 + 10(x-1)^3$
 - $p_3(x) = 5 + 3(x-1) + 18(x-1)^2 + 60(x-1)^3$
 - $p_3(x) = 5 + 3x + 9x^2 + 10x^3$
 - $p_3(x) = 5 + 3x + 18x^2 + 60x^3$
 - None of the others.
8. Using the geometric series, find a power series representation about (i.e., centered at) $x_0 = 5$ for the function

$$g(x) = \frac{3}{x-2}$$

and indicate when the representation is valid.

- $\sum_{n=0}^{\infty} \left(\frac{-1}{3}\right)^n (x-5)^n$, valid on $(2, 8)$.
- $\sum_{n=0}^{\infty} \left(\frac{1}{3}\right)^n (x-5)^n$, valid on $(2, 8)$.
- $\sum_{n=0}^{\infty} (-1)^n (x-5)^n$, valid on $(4, 6)$.
- $\sum_{n=0}^{\infty} (x-5)^n$, valid on $(4, 6)$.
- None of the others.

9. Using a known (commonly used) Taylor series, find the Taylor series for

$$f(x) = \frac{1}{(1-x)^4}$$

about the center $x_0 = 0$ which is valid for $|x| < 1$. Hint. Start with the Taylor series expansion

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k \quad \text{valid for } |x| < 1$$

and differentiate (as many times as needed). Be careful and don't forget the chain rule:

$$D_x(1-x)^{-1} = (-1)(1-x)^{-2} D_x(1-x) = (-1)(1-x)^{-2}(-1) = (1-x)^{-2}.$$

- $\sum_{n=0}^{\infty} \frac{(n)(n-1)(n-2)}{6} x^{n-3}$
- $\sum_{n=0}^{\infty} (n)(n-1)(n-2) x^n$
- $\sum_{n=0}^{\infty} \frac{(n+3)(n+2)(n+1)}{6} x^n$
- $\sum_{n=0}^{\infty} (-1)^n \frac{(n+3)(n+2)(n+1)}{6} x^n$
- None of the others.

10. Find a parameterization for the line segment from $(-1, 2)$ to $(10, -6)$ for $0 \leq t \leq 1$.

- $x = 10 - 8t$ and $y = -1 + t$
- $x = -1 + 11t$ and $y = 2 - 8t$
- $x = -1 + 11t$ and $y = -6 - 8t$
- $x = -1 - 11t$ and $y = -8t$
- None of the others.

11. A parametrization of a circle with center at $(0,0)$ and radius 1, which is traced out twice in the clockwise direction is
- $x(t) = \cos t$ and $y(t) = \sin t$ for $0 \leq t \leq 2\pi$
 - $x(t) = \cos t$ and $y(t) = \sin t$ for $0 \leq t \leq 4\pi$
 - $x(t) = \cos t$ and $y(t) = -\sin t$ for $0 \leq t \leq 2\pi$
 - $x(t) = \cos t$ and $y(t) = -\sin t$ for $0 \leq t \leq 4\pi$
 - None of the others.

12. Find an equation for the line tangent to the curve parameterized by

$$x = 2t^2 + 3$$

$$y = t^4$$

at the point defined by the value $t = -1$.

- $y = x - 6$
 - $y = x - 4$
 - $y = -x - 6$
 - $y = -x - 4$
 - None of the others.
13. Find the Cartesian coordinates of the point with polar coordinates

$$\left(-3, \frac{5\pi}{6}\right)$$

a. $\left(\frac{3}{2}, -\frac{3\sqrt{3}}{2}\right)$

b. $\left(-\frac{3}{2}, \frac{3\sqrt{3}}{2}\right)$

c. $\left(\frac{3\sqrt{3}}{2}, -\frac{3}{2}\right)$

d. $\left(-\frac{3\sqrt{3}}{2}, \frac{3}{2}\right)$

- e. None of the others.