

CONSTRUCTING COMPLETE PROJECTIVELY FLAT CONNECTIONS

RALPH HOWARD

ABSTRACT. On any open subset U of the Euclidean space \mathbf{R}^n there is complete torsion free connection whose geodesics are reparameterizations of the intersections of the straight lines of \mathbf{R}^n with U . For any positive integer m there is a complete projectively flat torsion free connection on the two dimensional torus such that for any point p there is another point q so that any broken geodesic from p to q has at least m breaks. This example is also homogeneous with respect to a transitive Lie group action.

1. INTRODUCTION.

The propose of this note is to tie up a couple of loose ends in the classical theory of linear connections. First, in [6, p. 395], Spivak rises the question of if, on a compact manifold with complete connection, any two points can be joined by a geodesic. The answer is “no” even when the connection is projectively flat and homogeneous:

Theorem 1. *Let T^2 be the two dimensional torus. Then for any positive integer m there is a complete torsion free projectively flat connection, ∇ , on T^2 such that for any point $p \in T^2$ there is a point $q \in T^2$ with the property that any broken ∇ -geodesic between p and q has at least m breaks. Moreover if T^2 is viewed as a Lie group in the usual manner, this connection is invariant under translations by elements of T^2 .*

Another natural question is: For a connected open subset, U , of the Euclidean space, \mathbf{R}^n , is the usual flat connection restricted to U projectively equivalent to complete torsion free connection on U ? This is true and is a special case of a more general result about connections on incomplete Riemannian manifolds.

Theorem 2. *Let (M, g) be a not necessarily complete Riemannian manifold. Then there is a complete torsion free connection on M that is projective with the metric connection on M . In particular any connected open subset M of the Euclidean space, \mathbf{R}^n , has a complete torsion free connection ∇ such that the geodesics of ∇ are reparameterizations of straight line segments of $M \subseteq \mathbf{R}^n$.*

Date: March 5, 2000.

Supported in part by DoD Grant No. N00014-97-1-0806.

The main tool is Proposition 2.2 which gives an elementary method of constructing complete torsion free connections that are projective with a given torsion free connection.

1.1. Definitions, notation and preliminaries. All of our manifolds are smooth (*i.e.* C^∞), Hausdorff, paracompact, and connected. The tangent bundle of M is denoted by $T(M)$. If $f: M \rightarrow N$ is a smooth map between manifolds, then the derivative map is $f_{*x}: T(M)_x \rightarrow T(M)_{f(x)}$.

We will use the term *connection* to stand for a linear connection on the tangent bundle (also called a Koszul connection) as defined in [4, Prop. 2.8 p. 123 and Prop. 7.5 p. 143] or [6, p. 241]. Let $c: (a, b) \rightarrow M$ be a smooth immersed curve. Then c is a ∇ -geodesic iff $\nabla_{c'(t)}c'(t) = 0$. The curve is a ∇ -pregeodesic iff there is a reparameterization of c that is a geodesic. This is equivalent to $\nabla_{c'(t)}c'(t) = \alpha(t)c'(t)$ for some smooth function $\alpha: (a, b) \rightarrow \mathbf{R}$. Given a pregeodesic $c: (a, b) \rightarrow M$ then an *affine parameterization* of c is a reparameterization $\sigma: (a_1, b_1) \rightarrow (a, b)$ so that $c \circ \sigma$ is a geodesic.

If $f: M \rightarrow N$ is a local diffeomorphism and ∇ is a connection on N then the *pull back connection* is the connection $f^*\nabla$ defined on M by $f_*((f^*\nabla)_X Y) = \nabla_{f_*X} f_*Y$. The connection ∇ on M is *homogeneous* on M iff there is a transitive action on M by a Lie group, G , so that $\varphi^*\nabla = \nabla$ for all $\varphi \in G$.

Two connections $\bar{\nabla}$ and ∇ on M are *projective* iff all geodesics of $\bar{\nabla}$ are pregeodesics of ∇ . This is an equivalence relation on the set of connections on M . If ∇_i is a connection on M_i for $i = 1, 2$ then a map $f: M_1 \rightarrow M_2$ is a *projective map* iff it is a local diffeomorphism and maps ∇_1 -geodesics to ∇_2 -pregeodesics. This is equivalent to the connections ∇_1 and $f^*\nabla_2$ on M_1 being projective. The connection ∇ is *projectively flat* iff every point $p \in M$ has an open neighborhood U and projective map $f: U \rightarrow \mathbf{R}^n$ where \mathbf{R}^n has its standard flat connection. Or what, is the same thing for every geodesic c of M the image $f \circ c$ is a reparameterization of interval in a line of \mathbf{R}^n . There is a well known criterion, due to Hermann Weyl, for two connections to be projective. A proof can be found in [6, Cor 19 p. 277].

1.1. Proposition (H. Weyl). *Two connections $\bar{\nabla}$ and ∇ on a manifold are projective and have the same torsion tensor if and only if there is a smooth one form ω so that the connections are related by*

$$(1.1) \quad \nabla_X Y = \bar{\nabla}_X Y + \omega(X)Y + \omega(Y)X.$$

Therefore if this relation holds and $\bar{\nabla}$ is torsion free, then so is ∇ . \square

Only the easy direction of this result will be used. That is if $\bar{\nabla}$ is torsion free and ∇ is given by (1.1) then ∇ is torsion free and projective with $\bar{\nabla}$. Note in this case if $c: (a, b) \rightarrow M$ is a $\bar{\nabla}$ -geodesic then (1.1) implies $\nabla_{c'(t)}c'(t) = 2\omega(c'(t))c'(t)$ and therefore c is a ∇ -pregeodesic. That ∇ is torsion free is equally as elementary.

The connection ∇ is *complete* iff every ∇ -geodesic defined on a subinterval of \mathbf{R} extends to a ∇ -geodesic defined on all of \mathbf{R} . Letting \exp^∇ be the

exponential of ∇ (cf. [4, p. 140]), then ∇ is easily seen to be complete if and only if the domain of \exp^∇ is all of $T(M)$. A curve $c: [0, b) \rightarrow M$ is an *inextendible $\bar{\nabla}$ -geodesic ray* iff c is a $\bar{\nabla}$ -geodesic and has no extension to $[0, b + \varepsilon)$ as a $\bar{\nabla}$ -geodesic for any $\varepsilon > 0$. Therefore when $b = \infty$, so that $[0, \infty)$ is the domain of c , c is always inextendible.

1.2. Proposition. *Let $\bar{\nabla}$ be a torsion free connection on the manifold M and let ∇ be torsion free and projective with $\bar{\nabla}$. Then ∇ is complete if and only if every inextendible $\bar{\nabla}$ -geodesic ray $c: [0, b) \rightarrow M$ has an orientation preserving reparameterization $\sigma: [0, \infty) \rightarrow [0, b)$ such that $c \circ \sigma$ is a ∇ -geodesic.*

Proof. First assume that the reparameterization condition holds and we will show that ∇ is complete by showing the domain of the exponential map of ∇ is all of $T(M)$. Let $v \in T(M)$. As 0 is in the domain of \exp^∇ , assume $v \neq 0$. Let $c: [0, b) \rightarrow M$ be the inextendible $\bar{\nabla}$ -geodesic ray with $c'(0) = v$. By assumption there is an orientation preserving reparameterization $\sigma: [0, \infty) \rightarrow [0, b)$ such that $\tilde{c} := c \circ \sigma$ is a ∇ -geodesic. As the reparameterization is orientation preserving $\tilde{c}'(0) = \lambda c'(0) = v$ for some positive constant λ . Then $\hat{c}: [0, \infty) \rightarrow M$ given by $\hat{c}(t) := \tilde{c}(t/\lambda)$ is also a ∇ -geodesic and $\hat{c}'(0) = v$. From the definition of \exp^∇ we have for all $t \geq 0$ that tv is in the domain of \exp^∇ and $\exp^\nabla(tv) = \hat{c}(t)$. In particular letting $t = 1$ shows that v is in the domain of \exp^∇ and completes the proof that ∇ is complete.

Conversely assume ∇ is complete and let $c: [0, b) \rightarrow M$ be an inextendible $\bar{\nabla}$ -geodesic ray. Assume, toward a contradiction, there is an orientation preserving reparameterization $\sigma: [0, b_1) \rightarrow [0, b)$ with $b_1 < \infty$ and so that $\tilde{c} = c \circ \sigma$ is a ∇ geodesic. Then, as ∇ is complete, the curve \tilde{c} extends to a ∇ -geodesic $\hat{c}: [0, \infty) \rightarrow M$ and therefore is a proper extension of \tilde{c} . But then \hat{c} can be reparameterized as a $\bar{\nabla}$ -geodesic that extends c , contradicting that c was an inextendible $\bar{\nabla}$ -geodesic ray and completing the proof. \square

2. CONSTRUCTING COMPLETE PROJECTIVELY EQUIVALENT CONNECTIONS ON INCOMPLETE RIEMANNIAN MANIFOLDS.

We first observe that for some choices of the one form ω in Weyl's result 1.1 there is an explicit formula for reparameterizing a $\bar{\nabla}$ -geodesic as a ∇ -geodesic.

2.1. Lemma. *Let $\bar{\nabla}$ be a smooth manifold and let $\bar{\nabla}$ be a connection on M and let $v: M \rightarrow (0, \infty)$ be a smooth positive function. Define a new connection by*

$$(2.1) \quad \nabla_X Y = \bar{\nabla}_X Y + \frac{1}{2v} dv(X)Y + \frac{1}{2v} dv(Y)X$$

Let $c: (a, b) \rightarrow M$ be a $\bar{\nabla}$ -geodesic and $\sigma: (\alpha, \beta) \rightarrow (a, b)$ an orientation preserving reparameterization of c so that $\tilde{c} = c \circ \sigma$ is a ∇ -geodesic. Then

the inverse of σ , $\sigma^{-1}: (a, b) \rightarrow (\alpha, \beta)$, is given by

$$(2.2) \quad \sigma^{-1}(t) = C_0 + C_1 \int_{t_0}^t v(c(\tau)) d\tau$$

where $t_0 \in (a, b)$, $C_0, C_1 \in \mathbf{R}$ and $C_1 > 0$.

Proof. Let t be the natural coordinate on (a, b) and s the coordinate on (α, β) related to t by $t = \sigma(s)$. Our goal is to find $s = s(t) = \sigma^{-1}(t)$. Note $dt = \sigma'(s) ds$ so that $\sigma'(s) = \frac{dt}{ds}$. Therefore

$$\tilde{c}'(s) = (c \circ \sigma)'(s) = \sigma'(s)c'(\sigma(s)) = \left. \frac{dt}{ds} \frac{dc}{dt} \right|_{t=\sigma(s)}.$$

Because of this, and because it makes applications of the chain rule easier to follow, we will denote $\tilde{c}'(s)$ as $\frac{dc}{ds}$ and think of s as “the affine parameter for ∇ along c ”. We will abuse notation a bit and write $v(t) = v(c(t))$. As $\bar{\nabla} \frac{dc}{dt} \frac{dc}{dt} = \bar{\nabla}_{c'(t)} c'(t) = 0$, we have using (2.1) that $\bar{\nabla} \frac{dc}{ds} \frac{dc}{dt} = \frac{dt}{ds} \bar{\nabla} \frac{dc}{dt} \frac{dc}{dt} = 0$, and $dv \left(\frac{dc}{ds} \right) = \frac{dv}{ds}$

$$\begin{aligned} 0 &= \nabla \frac{dc}{ds} \frac{dc}{ds} = \bar{\nabla} \frac{dc}{ds} \frac{dc}{ds} + \frac{1}{v} \left(\frac{dv}{ds} \right) \frac{dc}{ds} = \bar{\nabla} \frac{dc}{ds} \left(\frac{dt}{ds} \frac{dc}{dt} \right) + \frac{d(\ln v)}{ds} \frac{dc}{ds} \\ &= \frac{d^2 t}{ds^2} \frac{dc}{dt} + \frac{dt}{ds} \bar{\nabla} \frac{dc}{ds} \frac{dc}{dt} + \frac{d(\ln v)}{ds} \frac{dc}{ds} = \frac{d^2 t}{ds^2} \frac{dc}{dt} + \frac{d(\ln v)}{ds} \frac{dt}{ds} \frac{dc}{dt} \\ &= \left(\frac{dt}{ds} \right) \left(\left(\frac{dt}{ds} \right)^{-1} \frac{d^2 t}{ds^2} + \frac{d(\ln v)}{ds} \right) \frac{dc}{dt} = \left(\frac{dt}{ds} \right) \left(\frac{d}{ds} \ln \left(v \frac{dt}{ds} \right) \right) \frac{dc}{dt}. \end{aligned}$$

This shows that $\ln \left(v \frac{dt}{ds} \right)$, and therefore also $v \frac{dt}{ds}$, is constant. As $v, \frac{dt}{ds} > 0$ (the reparameterization is orientation preserving implies $\frac{dt}{ds} = \sigma'(s) > 0$) there is a constant $C_1 > 0$ such that

$$v(t) \frac{dt}{ds} = \frac{1}{C_1}.$$

This differential equation can be integrated to give $s(t) = \sigma^{-1}(t)$ as a function of t and the result is the required formula (2.2). \square

2.2. Proposition. *Let M be a smooth manifold with smooth torsion free connection $\bar{\nabla}$ and let $v: M \rightarrow (0, \infty)$ be a smooth positive function. Then the connection ∇ defined by (2.1) is a torsion free connection projective with $\bar{\nabla}$ and ∇ is complete if and only if for each inextendible $\bar{\nabla}$ -geodesic ray $c: [0, b) \rightarrow M$ the growth condition*

$$(2.3) \quad \int_0^b v(c(t)) dt = \infty.$$

holds.

Proof. That ∇ is projective to $\bar{\nabla}$ and torsion free follows from Proposition 1.1 using $\omega = (2v)^{-1}dv$. So all that is left to check is that ∇ is complete if and only if (2.3) holds along inextendible $\bar{\nabla}$ -geodesic rays.

First assume that the growth condition (2.3) holds along inextendible $\bar{\nabla}$ -geodesic rays. Let $c: [0, b) \rightarrow M$ be such a ray and let $\sigma: [0, \beta) \rightarrow [0, b)$ be an orientation preserving reparameterization of c so that $\tilde{c} = c \circ \sigma$ is a ∇ -geodesic. We claim that $\beta = \infty$. By Lemma 2.1 $\sigma^{-1}(t)$ is given by

$$(2.4) \quad \sigma^{-1}(t) = C_1 \int_0^t v(c(\tau)) d\tau$$

with $C_1 > 0$. But then the growth condition (2.3) implies $\beta = C_1 \int_0^b v(c(\tau)) d\tau = \infty$. As c was any inextendible $\bar{\nabla}$ -geodesic ray, the completeness of ∇ follows from Proposition 1.2.

Conversely assume ∇ is complete and let $c: [0, b) \rightarrow M$ be an inextendible $\bar{\nabla}$ -geodesic ray. Then by Proposition 1.2 there is an orientation preserving reparameterization $\sigma: [0, \infty) \rightarrow [0, b)$ so that $\tilde{c} = c \circ \sigma$ is a ∇ -geodesic. Again Lemma 2.1 implies that σ^{-1} is given by (2.4). Therefore $C_1 \int_0^b v(c(\tau)) d\tau = \lim_{t \uparrow b} \sigma^{-1}(t) = \infty$ which shows that the condition (2.3) holds along all inextendible $\bar{\nabla}$ -geodesic rays. \square

For a general connection, $\bar{\nabla}$, it is not clear how to choose a positive smooth function v so that the growth condition (2.3) holds along all inextendible $\bar{\nabla}$ -geodesics rays. However when ∇ is the metric connection of a Riemannian metric the behavior of geodesics is closely related to the properties of the distance function of the metric and this can be exploited to find an appropriate v .

Proof of Theorem 2. If (M, g) is complete as a metric space, then the metric connection $\bar{\nabla}$ is complete (cf. [7, p. 462]) and taking $\nabla = \bar{\nabla}$ completes the proof. Therefore assume that M is incomplete. Let \bar{M} be the completion of M as a metric space and let $\partial M = \bar{M} \setminus M$ be the boundary of M in \bar{M} . For $x \in M$ let $\delta(x)$ be the distance of x from ∂M . A standard partition of unity argument shows that there is a smooth function v on M so that

$$v(x) \geq \max\{1, 1/\delta(x)\}$$

for all $x \in M$. Let $c: [0, b) \rightarrow M$ be an inextendible $\bar{\nabla}$ -geodesic ray. There are two cases: $b = \infty$ and $b < \infty$. In the case $b = \infty$, then from the definition of v we have $v(c(t)) \geq 1$ and so $\int_0^b v(c(t)) dt \geq \int_0^\infty 1 dt = \infty$ and the condition (2.3) holds in this case.

In the second case, where $b < \infty$, the length of the velocity vector $c'(t)$ is constant and thus there is a constant $C > 0$ so that for all $t_1, t_2 \in [0, b)$ the distance $d(c(t_1), c(t_2))$ between $c(t_1)$ and $c(t_2)$ satisfies

$$d(c(t_1), c(t_2)) \leq C|t_2 - t_1|.$$

Therefore in the completion \bar{M} the limit $p = \lim_{t \uparrow b} c(t)$ will exist and from the definition of δ as the distance from the boundary ∂M the estimate

$\delta(c(t)) \leq d(c(t), p) \leq C|b - t|$ holds. This yields

$$\int_0^b v(c(t)) dt \geq \int_0^b \frac{dt}{\delta(c(t))} \geq \int_0^b \frac{dt}{C|b - t|} = \infty.$$

Thus (2.3) holds in all cases and therefore ∇ is complete by Proposition 2.2 \square

2.3. *Remark.* In a complete Riemannian manifold any two points can be joined by a geodesic. For complete connections this is no longer true and Hicks [3] has constructed an example of a complete connection on a manifold, M , so that for any positive integer m there are two points of M that not only can not be connected by a geodesic, but any broken geodesic between the points must have at least m breaks. For open sets U in \mathbf{R}^2 the behavior of geodesics is easy to visualize and, using Theorem 2, it is trivial to generate such examples that are also projectively flat. For example, set




FIGURE 1. Let $U \subset \mathbf{R}^2$ be the compliment of the pictured rays. Then there a complete torsion free connection on U whose geodesics are the restriction of the line segments of \mathbf{R}^2 to U .

$K := \bigcup_{k=-\infty}^{\infty} \{2k\} \times [-1, \infty) \cup \bigcup_{k=-\infty}^{\infty} \{2k + 1\} \times (-\infty, 1]$, which is a union of rays parallel to the y -axis, and let $U = \mathbf{R}^2 \setminus K$ (See Figure 1). Use Theorem 1 to put a complete projectively flat connection on U that has line segments as its geodesics and polygonal paths as its broken geodesics. With this connection U has the property that any broken geodesic between the points $(1/2, 0)$ and $(m + 1/2, 0)$ must have at least $m + 1$ corners. \square

3. HOMOGENEOUS EXAMPLES

Before specializing to two dimensions for the proof of Theorem 1 we do the preliminary calculations in arbitrary dimensions. This leads to higher dimensional examples.

Let $\bar{\nabla}$ be the standard flat connection on \mathbf{R}^n and let $U := \mathbf{R}^n \setminus \{0\}$ be \mathbf{R}^n with the origin deleted. Then any nonsingular linear map $A: \mathbf{R}^n \rightarrow \mathbf{R}^n$ preserves the connection $\bar{\nabla}$ and therefore the general linear group $\mathbf{GL}(n, \mathbf{R})$ has a transitive action on U that preserves $\bar{\nabla}$. Let $\mathbf{O}(n)$ be the orthogonal group of the standard inner product, $\langle \cdot, \cdot \rangle$, on \mathbf{R}^n and let \mathbf{R}^+ be the multiplicative group of positive real numbers. Let G be the product group $G = \mathbf{O}(n) \times \mathbf{R}^+$. View G as a subgroup of $\mathbf{GL}(n, \mathbf{R})$ by letting it act on \mathbf{R}^n by $(P, c)x = cPx$. This action of G is transitive on U and preserves the connection $\bar{\nabla}$. Let $v: U \rightarrow (0, \infty)$ be the function $v(x) = 1/\|x\|$. Then, if $g = (P, c) \in G$, the

pull back of v by g is $(g^*v)(x) = v(gx) = \|cPx\|^{-1} = c^{-1}\|x\|^{-1} = c^{-1}v(x)$ as $P \in \mathbf{O}(n)$ so that $\|Px\| = \|x\|$. The pull back of the one form dv/v is

$$g^* \left(\frac{dv}{v} \right) = \frac{g^*dv}{g^*v} = \frac{d(g^*v)}{g^*v} = \frac{d(c^{-1}v)}{c^{-1}v} = \frac{dv}{v}$$

and so dv/v is invariant under the action of G . Therefore if we define a connection ∇ on U by

(3.1)

$$\nabla_X Y = \overline{\nabla}_X Y + \overline{\nabla}_X Y + \frac{1}{2v} (dv(X)Y + dv(Y)X) \quad \text{with} \quad v(x) = \frac{1}{\|x\|}$$

then ∇ will be invariant under the action of the group G . The inextendible $\overline{\nabla}$ -geodesic rays in U are the curves $c: [0, b) \rightarrow U$ given by $c(t) = x_0 + tx_1$ where $x_1 \neq 0$ and either $b = \infty$ or $c(b) := \lim_{t \uparrow b} c(t) = 0$. In either case it is easy to check that $\int_0^b v(c(t)) dt = \infty$ and therefore by Proposition 2.2 the connection ∇ is complete and projectively flat on U .

To get compact examples let $\lambda > 1$ and let Γ be the cyclic subgroup of G given by $\Gamma := \{(I, \lambda^k) : k \in \mathbf{Z}\}$ where \mathbf{Z} is the integers. The action of Γ on U is fixed point free and properly discontinuous and therefore if M is defined to be the quotient space $M := \Gamma \backslash U$ then M is a smooth manifold (cf. [1, Thm 8.3 p. 97]) and it is not hard to see that M is diffeomorphic to $S^{n-1} \times S^1$. Let $\pi: U \rightarrow M$ be the natural projection. Then π is a covering map and Γ is the group of deck transformations. As the connection ∇ is invariant under these transformations it follows there is a unique connection ∇^M on M so that $\pi^* \nabla^M = \nabla$. The ∇^M -geodesics on M are $\pi \circ c$ where c is a ∇ -geodesic on U . As the ∇ -geodesics in U are complete, it follows that the ∇^M geodesics in M are complete. Also this implies that π is a projective map and therefore ∇^M is projectively flat on M .

For any $g = (P, c) \in G$ and $a = (I, \lambda^k) \in \Gamma$ we have $ag = ga$. As for $x \in U$ the image $\pi(x)$ is the orbit $\pi(x) = \Gamma x$ we see for $g \in \Gamma$ that $\pi(gx) = \Gamma gx = g\Gamma x = g\pi(x)$. Therefore there is a well defined action of G on M given by $g\pi(x) = \pi(gx)$. This action is transitive on M as G is transitive on U .

We now claim that if $x \in U$ and $y = -\alpha x$ for $\alpha > 0$, then there is no geodesic from $\pi(x)$ to $\pi(y)$ in M . Assume, toward a contradiction, that there is a geodesic $c: [a, b] \rightarrow M$ with $c(a) = \pi(x)$ and $c(b) = \pi(y)$. Then there is a unique geodesic $\hat{c}: [a, b] \rightarrow U$ with $\hat{c}(a) = x$ and $\pi \circ \hat{c} = c$. Therefore $\pi(\hat{c}(b)) = c(b) = \pi(y)$ which implies that $\hat{c}(b) = ay$ for some $a \in \Gamma$. From the definition of Γ this implies that for some $k \in \mathbf{Z}$ that $\hat{c}(b) = \lambda^k y = -\lambda^k \alpha x$. But as ∇ is projective with the flat metric $\overline{\nabla}$ the geodesics segments of ∇ are reparameterizations of straight line segments in U . But then \hat{c} is a reparameterization of a straight line segment of U from $\hat{c}(a) = x$ to $\hat{c}(b) = -\lambda^k \alpha x$, which is impossible as $\lambda^k \alpha > 0$ so that any line segment connecting these points must pass through the origin, which is not in U . This contradiction verifies our claim that there is no geodesic

of M from $\pi(x)$ to $\pi(y)$. Letting α vary over the positive real numbers we get uncountable many points $\pi(y)$ that can not be connected to $\pi(x)$ by a geodesic. As every point $p \in M$ is of the form $p = \pi(x)$ this can be summarized as:

3.1. Proposition. *Let $M = \Gamma \backslash U$ and ∇^M be the manifold and connection just constructed. Then M is diffeomorphic to $S^{n-1} \times S^1$ and the connection ∇^M on M is complete, projectively flat and with homogeneous with respect to the group action of G on M . For any $p \in M$ there are uncountable many points q that can not be connected to p by a ∇^M -geodesic. \square*

3.1. Proof of Theorem 1. In the case that $n = 2$ it is possible to be more explicit. On $U = \mathbf{R}^2 \setminus \{0\}$ there are several sets of coordinates that will be convenient to use. First the standard Euclidean coordinates x and y . With respect to these coordinates the standard flat connection $\bar{\nabla}$ is given by $\bar{\nabla}_{\frac{\partial}{\partial x}} \frac{\partial}{\partial x} = \bar{\nabla}_{\frac{\partial}{\partial x}} \frac{\partial}{\partial y} = \bar{\nabla}_{\frac{\partial}{\partial y}} \frac{\partial}{\partial x} = \bar{\nabla}_{\frac{\partial}{\partial y}} \frac{\partial}{\partial y} = 0$.

The simply connected covering space, \widehat{U} , of U is diffeomorphic to \mathbf{R}^2 . Using polar coordinates r, θ on \widehat{U} (with $(r, \theta) \in (0, \infty) \times \mathbf{R}$) we have the usual formula for the covering map: $x = r \cos \theta$ and $y = r \sin \theta$. In polar coordinates the connection is given by

$$\bar{\nabla}_{\frac{\partial}{\partial r}} \frac{\partial}{\partial r} = 0, \quad \bar{\nabla}_{\frac{\partial}{\partial r}} \frac{\partial}{\partial \theta} = \bar{\nabla}_{\frac{\partial}{\partial \theta}} \frac{\partial}{\partial r} = \frac{1}{r} \frac{\partial}{\partial \theta}, \quad \bar{\nabla}_{\frac{\partial}{\partial \theta}} \frac{\partial}{\partial \theta} = -r \frac{\partial}{\partial r}.$$

(More precisely this is the pull back of the connection ∇ to \widehat{U} by the covering map. We will still denote this connection by ∇ .) The function $v = \|(x, y)\|^{-1}$ used in the definition (3.1) of the connection ∇ is given in polar coordinates as $v = r^{-1}$. Then $dv = -r^{-2} dr$. Using this in (3.1) gives

$$\nabla_X Y = \bar{\nabla}_X Y - \frac{1}{r} (dr(X)Y + dr(Y)X)$$

and therefore ∇ is given explicitly in polar coordinates as

$$\nabla_{\frac{\partial}{\partial r}} \frac{\partial}{\partial r} = \frac{-1}{r} \frac{\partial}{\partial r}, \quad \nabla_{\frac{\partial}{\partial r}} \frac{\partial}{\partial \theta} = \nabla_{\frac{\partial}{\partial \theta}} \frac{\partial}{\partial r} = \frac{1}{2r} \frac{\partial}{\partial \theta}, \quad \nabla_{\frac{\partial}{\partial \theta}} \frac{\partial}{\partial \theta} = -r \frac{\partial}{\partial r}.$$

The formulas for ∇ simplify even farther if we replace the coordinate r on \widehat{U} by ρ related to r by $r = e^\rho$. The vector field $\frac{\partial}{\partial \rho}$ is related to the vector field $\frac{\partial}{\partial r}$ by $\frac{\partial}{\partial \rho} = r \frac{\partial}{\partial r}$ and $\frac{\partial}{\partial r} = e^{-\rho} \frac{\partial}{\partial \rho}$. Therefore in the coordinates ρ, θ the connection ∇ is given by

$$\nabla_{\frac{\partial}{\partial \rho}} \frac{\partial}{\partial \rho} = 0, \quad \nabla_{\frac{\partial}{\partial \rho}} \frac{\partial}{\partial \theta} = \nabla_{\frac{\partial}{\partial \theta}} \frac{\partial}{\partial \rho} = \frac{1}{2} \frac{\partial}{\partial \theta}, \quad \nabla_{\frac{\partial}{\partial \theta}} \frac{\partial}{\partial \theta} = -\frac{\partial}{\partial \rho}.$$

This explicit form of the connection ∇ makes it clear that it is invariant under translations $\rho \mapsto \rho + a$ and $\theta \mapsto \theta + b$. From the construction ∇ is complete and projectively flat.

Using the coordinates ρ and θ and letting \mathbf{Z} be the integers, then the original open set U is naturally identified with the quotient group $\mathbf{R}^2 / (\{0\} \times 2\pi\mathbf{Z})$ (that is identify (ρ, θ) with $(\rho, \theta + 2k\pi)$ for $k \in \mathbf{Z}$). As in the original

set U the ∇ -geodesics are reparameterized line segments it is not hard to see that a point $z \in U$ can be connected to a point z_0 on the positive real axis by a ∇ -geodesic if and only if z is not on the negative real axis. That is z can be connected to z_0 by a ∇ -geodesic if and only if $|\theta(z)| < \pi$. (See




FIGURE 2. As the connection ∇ is projective with the usual flat connection, a point z in the set $U = \mathbf{R}^2 \setminus \{0\}$ can be connected to a point z_0 on the positive real axis by a ∇ -geodesic if and only if $|\theta(z)| < \pi$.

Figure 2.) But because of the homogeneity of the connection with respect to translations $\theta \mapsto \theta + b$ this implies:

3.2. Lemma. *Two points $z_1, z_2 \in \widehat{U}$ can be connected by a ∇ -geodesic if and only if $|\theta(z_1) - \theta(z_2)| < \pi$. Therefore if z_1, z_2 satisfy $|\theta(z_1) - \theta(z_2)| \geq m\pi$ for some positive integer m any piecewise broken geodesic from z_1 to z_2 must have at least m breaks. \square*

3.3. Remark. There is a less geometric, but possibly more informative, proof of this lemma. Using the coordinates ρ, θ on \widehat{U} and the coordinates x, y on U , the covering map from \widehat{U} to U is given by $x = e^\rho \cos \theta$ and $y = e^\rho \sin \theta$. In U the ∇ -geodesics are reparameterization of straight lines and thus along a ∇ -geodesic the coordinates x and y are related by $ax + by = 0$ (if geodesic goes through the origin) or $ax + by = 1$ (if it does not pass through the origin). The first case leads to a relation between ρ and θ of the form $e^\rho(a \cos \theta + b \sin \theta) = 0$ along the geodesic which implies $\theta = \theta_0$ on the geodesic, for some constant θ_0 . In the second case we get $e^\rho(a \cos \theta + b \sin \theta) = 1$ along the geodesic. Let $A = \sqrt{a^2 + b^2}$ and let α be so that $A \cos \alpha = a$ and $A \sin \alpha = b$. Then the equation between ρ and θ becomes $e^\rho A \cos(\theta - \alpha) = 1$. From this it follows that given a point in \widehat{U} with coordinates (ρ_0, θ_0) the ∇ -geodesics of \widehat{U} through this point are the line $\theta = \theta_0$ and the curves defined for $|\theta - \alpha| < \pi/2$ by the equation

$$(3.2) \quad e^\rho \cos(\theta - \alpha) = e^{\rho_0} \cos(\theta_0 - \alpha)$$

where α varies over real numbers with $|\alpha - \theta_0| < \pi/2$. This makes it clear a point (ρ_1, θ_1) with $|\theta_1 - \theta_0| \geq \pi$ can not be on a geodesic through (ρ_0, θ_0) . And conversely if $|\theta_1 - \theta_0| < \pi$ then either $\theta_1 = \theta_0$, and the points are both on the geodesic $\theta = \theta_0$, or $\theta_1 \neq \theta_0$ and straightforward calculus argument shows that there is a unique $\alpha \in (\theta_0 - \pi/2, \theta_0 + \pi/2) \cap (\theta_1 - \pi/2, \theta_1 + \pi/2)$ so that $e^{\rho_1} \cos(\theta_1 - \alpha) = e^{\rho_0} \cos(\theta_0 - \alpha)$. For this choice of α both of the points (ρ_0, θ_0) and (ρ_1, θ_1) will be on the ∇ -geodesic defined by (3.2) \square

We now complete the proof of Theorem 1. Given the positive integer m let k be an integer with $k \geq m$. Let T^2 be the torus

$$T^2 = \widehat{U}/(\mathbf{Z} \times 2\pi k\mathbf{Z})$$

(that is identify (ρ, θ) with $(\rho + j, \theta + 2\pi k\ell)$ for $j, \ell \in \mathbf{Z}$). As the connection ∇ is translation invariant it well defined as a connection on T^2 and will be invariant under translations of T^2 when T^2 is viewed as a Lie group. We have already seen that ∇ is complete and projectively flat. Let $\varpi: \widehat{U} \rightarrow T^2$ be the covering map. We now claim that any broken ∇ -geodesic in T^2 from $\varpi(\rho_0, \theta_0)$ to $\varpi(\rho_0, \theta_0 + m\pi)$ must have at least m breaks. For let $c: [a, b] \rightarrow T^2$ be such a broken geodesic. By the Path Lifting Theorem ([2, p. 22] or [5, p. 67]) there is a unique curve $\hat{c}: [a, b] \rightarrow \widehat{U}$ with $\hat{c}(a) = (\rho_0, \theta_0)$ and $\varpi \circ \hat{c} = c$. This curve will also be a broken geodesic. Also $\varpi(\hat{c}(b)) = c(b) = \varpi(\rho_0, \theta_0 + m\pi)$, and therefore $\hat{c}(b) = (\rho_0 + j, \theta_0 + m\pi + 2\pi k\ell)$ for some $j, \ell \in \mathbf{Z}$. The difference in the θ coordinates of the ends of \hat{c} is

$$|\theta_0 + m\pi + 2\pi k\ell - \theta_0| = |m + 2k\ell|\pi \geq m\pi$$

as $k \geq m$. By Lemma 3.2 this implies that \hat{c} has at least m breaks. But then $c = \varpi \circ \hat{c}$ also has at least m breaks. As $\varpi(\rho_0, \theta_0)$ was an arbitrary point of T^2 this completes the proof of Theorem 1. \square

3.4. *Remark.* The connection ∇ has another property worth noting. If $c(t) = (\rho(t), \theta(t))$ is a smooth curve in \widehat{U} then the equations for c to be a ∇ -geodesic are

$$\ddot{\rho} = \dot{\theta}^2, \quad \ddot{\theta} = -\dot{\rho}\dot{\theta}.$$

These imply

$$\frac{1}{2} \frac{d}{dt}(\dot{\rho}^2 + \dot{\theta}^2) = \dot{\rho}\ddot{\rho} + \dot{\theta}\ddot{\theta} = \dot{\rho}\dot{\theta}^2 - \dot{\theta}\dot{\rho}\dot{\theta} = 0.$$

Therefore $\dot{\rho}^2 + \dot{\theta}^2$ is constant along ∇ -geodesics. Thus all ∇ -geodesics have constant speed with respect to the flat Riemannian metric $ds^2 = d\rho^2 + d\theta^2$ on \widehat{U} . As this metric is translation invariant it is also well defined on the torus $T^2 = \widehat{U}/(\mathbf{Z} \times 2\pi k\mathbf{Z})$ and the ∇ -geodesics on T^2 will also have constant speed with respect to this metric. This can be used to give another proof that ∇ is complete \square

REFERENCES

- [1] W. M. Boothby, *An introduction to differentiable manifolds and Riemannian geometry*, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975, Pure and Applied Mathematics, No. 63.
- [2] M. J. Greenberg and J. R. Harper, *Algebraic topology*, Benjamin/Cummings Publishing Co. Inc. Advanced Book Program, Reading, Mass., 1981, A first course.
- [3] N. J. Hicks, *An example concerning affine connexions*, Proc. Amer. Math. Soc. **11** (1960), 952–956.
- [4] S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, vol. 1, John Wiley and Sons, 1963.
- [5] E. H. Spanier, *Algebraic topology*, McGraw-Hill Book Co., New York, 1966.

- [6] M. Spivak, *A comprehensive introduction to differential geometry*, 2 ed., vol. 2, Publish or Perish Inc., Berkeley, 1979.
- [7] ———, *A comprehensive introduction to differential geometry*, 2 ed., vol. 1, Publish or Perish Inc., Berkeley, 1979.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, S.C.
29208, USA

E-mail address: `howard@math.sc.edu`