


# Section 11.3: Polar Coordinates

**Definition:** To define polar coordinates, we first fix an \_\_\_\_\_  $O$  (called the \_\_\_\_\_) and an \_\_\_\_\_ from  $O$  (usually the positive  $x$ -axis). Then each point  $P$  can be located by assigning to it a \_\_\_\_\_  $(r, \theta)$  in which  $r$  gives the directed distance from  $O$  to  $P$  and  $\theta$  gives the directed angle from the initial ray to the ray  $OP$ .



Just like trigonometry,  $\theta$  is positive when measured anticlockwise and negative when measured clockwise. The angle associated with a given point is not unique.

In some cases, we allow  $r$  to be negative. For instance, the point  $P(2, 7\pi/6)$  can be reached by turning  $7\pi/6$  radians anticlockwise from the initial ray and going forward 2 units, or we could turn  $\pi/6$  radians clockwise and go backwards 2 units; corresponding to  $P(-2, \pi/6)$ .

**Example 1:** Find all the polar coordinates of the point  $P(2, \frac{\pi}{6})$ .

**Polar Equations and Graphs:** If we fix  $r$  at a constant value (not equal to zero), the point  $P(r, \theta)$  will lie  $|r|$  unites from the origin  $O$ . As  $\theta$  varies over any interval of length  $2\pi$ ,  $P$  traces a what?

If we fix  $\theta$  at a constant value and let  $r$  vary between  $-\infty$  and  $\infty$ , then the point  $P(r, \theta)$  traces a what?

**Example 2:** A circle or line can have more than one polar equation.

**Example 3:** Equations of the form  $r = a$  and  $\theta = \theta_0$  can be combined to define regions, segments and rays. Graph the sets of points whose polar coordinates satisfy the given conditions:

(a)  $1 \leq r \leq 2$  and  $0 \leq \theta \leq \frac{\pi}{2}$

(b)  $-3 \leq r \leq 2$  and  $\theta = \frac{\pi}{4}$

(c)  $\frac{2\pi}{3} \leq \theta \leq \frac{5\pi}{6}$

**Relating Polar and Cartesian Coordinates:** When we use both polar and Cartesian coordinates in a plane, we place the two origins together and take the initial ray as the positive  $x$ -axis. The ray  $\theta = \pi/2$ ,  $r > 0$  becomes the positive  $y$ -axis. The two coordinate systems are then related by the following:

**Example 4:** Given the polar equation, find the Cartesian equivalent:

(a)  $r \cos(\theta) = 2$

(b)  $r^2 \cos(\theta) \sin(\theta) = 4$

(c)  $r^2 \cos^2(\theta) - r^2 \sin^2(\theta) = 1$

(d)  $r = 1 + 2r \cos(\theta)$

(e)  $r = 1 - \cos(\theta)$

**Example 5:** Find a polar equation for the circle  $x^2 + (y - 3)^2 = 9$ .