Math 142, Exam 3, Fall 2009

Use my paper. **Please turn the problems in order. Please leave 1 square inch in the upper left hand corner for the staple.**

The exam is worth 50 points. There are 9 problems. SHOW your work. $\boxed{\textit{CIRCLE}}$ your answer. **CHECK** your answer whenever possible. **No Calculators.**

1. **Find the limit of the sequence whose n^{th} term is** $a_n = n\sin\left(\frac{3}{n}\right)$.

We see that

$$\lim_{n\to\infty} a_n = \lim_{n\to\infty} n\sin\left(\frac{3}{n}\right) = \lim_{n\to\infty} \frac{\sin\left(\frac{3}{n}\right)}{\frac{1}{n}}.$$

L'hopitals's rule applies since the top and the bottom both go to zero; so,

$$\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{\frac{-3}{n^2}\cos\left(\frac{3}{n}\right)}{\frac{-1}{n^2}} = \lim_{n\to\infty} 3\cos\left(\frac{3}{n}\right) = 3.$$

$$\boxed{\text{The sequence } \{a_n\} \text{ converges to 3.}}$$

2. **Does the series** $\sum_{n=1}^{\infty}\left(1-\frac{2}{n}\right)^n$ **converge? Justify your answer.**

The limit of the individual terms $\left(1-\frac{2}{n}\right)^n$ is equal to

$$\lim_{n\to\infty}\left(1-\frac{2}{n}\right)^n = e^{-2} \neq 0.$$

The Individual Term Test for Divergence tells me that

$$\boxed{\text{the series } \sum_{n=1}^{\infty}\left(1-\frac{2}{n}\right)^n \text{ diverges.}}$$

3. **Consider the following sequence of numbers:** $a_2 = (1-\frac{1}{4})$, $a_3 = (1-\frac{1}{4})(1-\frac{1}{9})$, $a_4 = (1-\frac{1}{4})(1-\frac{1}{9})(1-\frac{1}{16})$, $\dots$, $a_n = (1-\frac{1}{4})(1-\frac{1}{9})(1-\frac{1}{16})\dots(1-\frac{1}{n^2})$, $\dots$. **Does this infinite sequence converge? Justify your answer.**

This is a sequence of positive numbers. Observe that $a_{n+1} = a_n(1 - \frac{1}{(n+1)^2})$. The number $1 - \frac{1}{(n+1)^2}$ is less than 1; so $a_{n+1} < a_n$. We are studying a decreasing bounded sequence of real numbers . The completeness axiom assures us that

$$\boxed{\text{the sequence } \{a_n\} \text{ converges.}}$$

4. **Does the series** $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^{n+1}}{\sqrt{n}}$ **converge? Justify your answer.**

We use the Alternating Series Test. Notice that the series alternates; $\frac{1}{\sqrt{n}} > \frac{1}{\sqrt{n+1}}$ for all n; and $\lim\limits_{n\to\infty} \frac{1}{\sqrt{n}} = 0$. The Alternating Series Test tells us that

$$\boxed{\text{the series } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\sqrt{n}} \text{ converges.}}$$

5. **Does the series** $\sum\limits_{n=1}^{\infty} \dfrac{2\sqrt{n}}{n^2+1}$ **converge? Justify your answer.**

I do a limit comparison with the p-series $\sum\limits_{n=1}^{\infty} \frac{1}{n^{3/2}}$. The p-series $\sum\limits_{n=1}^{\infty} \frac{1}{n^{3/2}}$ has $p = 3/2 > 1$; so $\sum\limits_{n=1}^{\infty} \frac{1}{n^{3/2}}$ converges. Observe that

$$\lim_{n\to\infty} \frac{\frac{2\sqrt{n}}{n^2+1}}{\frac{1}{n^{3/2}}} = \lim_{n\to\infty} \frac{2n^2}{n^2+1} = 2.$$

Two is a number. It is not zero or infinity; so the limit comparison test tells us that $\sum\limits_{n=1}^{\infty} \dfrac{2\sqrt{n}}{n^2+1}$ and $\sum\limits_{n=1}^{\infty} \frac{1}{n^{3/2}}$ both converge or both diverge. We have already seen that $\sum\limits_{n=1}^{\infty} \frac{1}{n^{3/2}}$ converges. We conclude that

$$\boxed{\text{the series } \sum_{n=1}^{\infty} \frac{2\sqrt{n}}{n^2+1} \text{ converges.}}$$

6. **Where does** $f(x)=\displaystyle\sum_{n=1}^{\infty}\frac{(x-2)^n}{n3^n}$ **converge? Justify your answer.**

Apply the ratio test. Let

$$\rho=\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\frac{\frac{(x-2)^{n+1}}{(n+1)3^{n+1}}}{\frac{(x-2)^n}{n3^n}}\right|=\lim_{n\to\infty}\left|\frac{(x-2)^{n+1}n3^n}{(x-2)^n(n+1)3^{n+1}}\right|$$

$$=\lim_{n\to\infty}\frac{|x-2|}{3}\frac{n}{(n+1)}=\frac{|x-2|}{3}$$

Notice that $\rho<1$ when $\frac{|x-2|}{3}<1$; which is the same as $|x-2|<3$; which is the same as $-3<x-2<3$ or $-1<x<5$. If $-1<x<5$, then $f(x)$ converges. If $x<-1$ or $5<x$, then $f(x)$ diverges. At -1,

$$f(-1)=\sum_{n=1}^{\infty}\frac{(-3)^n}{n3^n}=\sum_{n=1}^{\infty}\frac{(-1)^n}{n}.$$

Thus, $f(-1)$ is minus the sum of the Alternating Harmonic series, which we know converges. At 5,

$$f(5)=\sum_{n=1}^{\infty}\frac{3^n}{n3^n}=\sum_{n=1}^{\infty}\frac{1}{n};$$

which is the harmonic series which diverges. We conclude that

the function $f(x)$ converges for $-1\le x<5$ and diverges everywhere else.

7. **Approximate** $e^{\frac{-1}{10}}$ **with an error at most** 10^{-3}**. Explain what you are doing.**

We apply Taylor's Theorem which says that

$$|e^x-P_n(x)|=|R_n(x)|=\frac{e^c x^{n+1}}{(n+1)!}$$

for some c between 0 and x. In particular, when $x=-.1$, we have $-.1\le c\le 0$; hence, $e^{-.1}\le e^c\le e^0=1$ (since $y=e^x$ is an increasing function). So, $|R_n(-1/10)|\le\frac{1}{10^{n+1}(n+1)!}$. We want $|R_n(-1/10)|\le 10^{-3}$. In other words, we want $10^3 \le 10^{n+1}(n+1)!$. This inequality holds for $n=2$. We calculate $P_2(x) = 1 + x + x^2/2$. We conclude that

$$\boxed{1 - \frac{1}{10} + \frac{1}{2\times 10^2} \text{ approximates } e^{\frac{-1}{10}} \text{ with an error of at most } 10^{-3}.}$$

8. **Approximate $\sum_{n=10}^{\infty} \frac{1}{n^2}$. Your approximation should be "close to" but more than the actual value. Explain what you are doing.**

I will use an improper integral to approximate $\sum_{n=10}^{\infty} \frac{1}{n^2}$. Draw the graph of $y = \frac{1}{x^2}$. Approximate the area under the curve starting at $x = 9$ by drawing rectangles with base from one integer to the next and height given by the y-coordinate on the curve above the right hand endpoint. I put a picture on a separate piece of paper. The area inside the boxes underestimates the area under the curve. The area inside the boxes is $\sum_{n=10}^{\infty} \frac{1}{n^2}$. The area under the curve is $\int_9^{\infty} x^{-2}dx$. Thus,

$$\sum_{n=10}^{\infty} \frac{1}{n^2} \le \int_9^{\infty} x^{-2}dx = \lim_{b\to\infty} \left.\frac{-1}{x}\right|_9^b$$

$$= \lim_{b\to\infty} \frac{-1}{b} + \frac{1}{9} = \frac{1}{9}.$$

We conclude that

$$\boxed{\sum_{n=10}^{\infty} \frac{1}{n^2} \le \frac{1}{9}.}$$

9. **A ball is dropped from a height of 100 feet. Each time it bounces, it rebounds to $\frac{4}{5}$ its previous height. Find the total distance it travels before coming to rest.**

The ball goes down 100 feet, up $\left(\frac{4}{5}\right)100$ feet, down $\left(\frac{4}{5}\right)100$ feet, up $\left(\frac{4}{5}\right)^2 100$ feet, down $\left(\frac{4}{5}\right)^2 100$ feet, etc. In total the ball travels

$$100 + \left(\tfrac{4}{5}\right)100 + \left(\tfrac{4}{5}\right)100 + \left(\tfrac{4}{5}\right)^2 100 + \left(\tfrac{4}{5}\right)^2 100 + \left(\tfrac{4}{5}\right)^3 100 + \left(\tfrac{4}{5}\right)^3 100 + \dots$$

$$= 100 + \left[\left(\tfrac{4}{5}\right) 200 + \left(\tfrac{4}{5}\right)^2 200 + \left(\tfrac{4}{5}\right)^3 200 + \ldots \right]$$

feet. The expression inside the brackets is the geometric series with $r = \frac{4}{5}$ and $a = \left(\frac{4}{5}\right) 200$. This geometric series converges to $\frac{a}{1-r} = \frac{\left(\frac{4}{5}\right)200}{\frac{1}{5}} = 800$. The ball travels $100 + 800 = \boxed{900 \text{ feet}}$.