

PRINT Your Name: \_\_\_\_\_

**Quiz for June 14, 2012**

The quiz is worth 5 points. **Remove EVERYTHING from your desk except this quiz and a pen or pencil.** SHOW your work. Express your work in a neat and coherent manner. BOX your answer.

Solve the Initial Value Problem

$$x'' + 10x' + 125x = 0, \quad x(0) = 6, \quad x'(0) = 50.$$

If possible, put your answer in the form  $x(t) = C_1 e^{-pt} \cos(\omega_1 t - \alpha_1)$ .

**Answer.** We try  $x = e^{rt}$ . We immediately are faced with solving the characteristic equation:

$$r^2 + 10r + 125 = 0.$$

So,  $r = \frac{-10 \pm \sqrt{100 - 500}}{2} = \frac{-10 \pm 20i}{2} = -5 \pm 10i$ . So  $x = c_1 e^{-5t} \cos 10t + c_2 e^{-5t} \sin 10t$  is the general solution of the differential equation. We now plug in the Initial Conditions to determine the values of  $c_1$  and  $c_2$ . We calculate

$$x' = e^{-5t}(-10c_1 \sin 10t - 5c_1 \cos 10t + 10c_2 \cos 10t - 5c_2 \sin 10t).$$

We have  $6 = x(0) = c_1$  and  $50 = x'(0) = -5c_1 + 10c_2$ ; so,  $c_1 = 6$  and  $c_2 = 8$ . The solution is

$$x(t) = e^{-5t}(6 \cos 10t + 8 \sin 10t) = 10e^{-5t}\left(\frac{6}{10} \cos 10t + \frac{8}{10} \sin 10t\right).$$

We have arranged the numbers so that  $(\frac{6}{10}, \frac{8}{10})$  is a point on the unit circle. Let  $\alpha = \arccos(\frac{6}{10})$ . It follows that  $\sin \alpha = \frac{8}{10}$ . We have

$$x(t) = 10e^{-5t}(\cos \alpha \cos 10t + \sin \alpha \sin 10t) = 10e^{-5t} \cos(\alpha - 10t) = 10e^{-5t} \cos(10t - \alpha).$$

Our answer is

$x(t) = 10e^{-5t} \cos(10t - \alpha), \text{ where } \alpha = \arccos(\frac{3}{5}).$