

Math 544, Exam 3, Summer 2005, solution

Write your answers as legibly as you can on the blank sheets of paper provided. Use only **one side** of each sheet. Be sure to number your pages. Put your solution to problem 1 first, and then your solution to number 2, etc.; although, by using enough paper, you can do the problems in any order that suits you.

There are 7 problems. Problem 1 is worth 14 points. Each of the other problems is worth 6 points. The exam is worth a total of 50 points. **SHOW** your work. **CIRCLE** your answer. **CHECK** your answer whenever possible. **No Calculators.**

If I know your e-mail address, I will e-mail your grade to you. If I don't already know your e-mail address and you want me to know it, then **send me an e-mail.**

If you would like, I will leave your graded exam outside my office door. You may pick it up any time before the next class. **If you are interested, be sure to tell me.**

I will post the solutions on my website shortly after the class is finished.

1. **Let A be the matrix**

$$A = \begin{bmatrix} 1 & 3 & 4 & 2 & 4 \\ 1 & 3 & 4 & 3 & 6 \\ 2 & 6 & 8 & 5 & 10 \end{bmatrix}$$

- (a) **Find a basis for the null space of A .**
- (b) **Find a basis for the column space of A .**
- (c) **Find a basis for the row space of A .**
- (d) **Write each column of A as a linear combination of your answer to (b).**
- (e) **Write each row of A as a linear combination of your answer to (c).**

Apply elementary row operations $R_2 \mapsto R_2 - R_1$ and $R_3 \mapsto R_3 - 2R_1$ to get

$$\begin{bmatrix} 1 & 3 & 4 & 2 & 4 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix}.$$

Apply elementary row operations $R_1 \mapsto R_1 - 2R_2$ and $R_3 \mapsto R_3 - R_2$ to get

$$\begin{bmatrix} 1 & 3 & 4 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The nullspace of A is the set of all vectors:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = x_2 \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -4 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} 0 \\ 0 \\ 0 \\ -2 \\ 1 \end{bmatrix},$$

where x_2 , x_3 , and x_5 are free to take any value.

(a) It follows that the vectors

$$v_1 = \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -4 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad \text{and} \quad v_3 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ -2 \\ 1 \end{bmatrix}$$

form a basis for the null space of A .

(b) The vectors

$$A_{*,1} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \quad \text{and} \quad A_{*,4} = \begin{bmatrix} 2 \\ 3 \\ 5 \end{bmatrix}$$

form a basis for the column space of A .

(c) The vectors

$$w_1 = [1 \ 3 \ 4 \ 0 \ 0] \quad \text{and} \quad w_2 = [0 \ 0 \ 0 \ 1 \ 2]$$

form a basis for the row space of A .

(d) We see that

$$\begin{aligned} A_{*,1} &= A_{*,1} \\ A_{*,2} &= 3A_{*,1} \\ A_{*,3} &= 4A_{*,1} \\ A_{*,4} &= A_{*,4} \\ A_{*,5} &= 2A_{*,4} \end{aligned}.$$

(e) We see that

$$\begin{aligned} A_{1,*} &= 1w_1 + 2w_2 \\ A_{2,*} &= w_1 + 3w_2 \\ A_{3,*} &= 2w_1 + 5w_2 \end{aligned}.$$

2. **Let $U \subseteq V$ be subspaces of \mathbb{R}^n with $\dim U = \dim V$. Do U and V HAVE to be equal? If yes, prove your answer. If no, give an example.**

YES. Let u_1, \dots, u_r be a basis for U . Thus u_1, \dots, u_r is a linearly independent set in the vector space V . One of the dimension theorems tells us that u_1, \dots, u_r is the beginning of a basis for V ; that is, we may adjoin more vectors to this list, if necessary, to get a basis for V . However every basis for V has $\dim V$ vectors and $\dim V = \dim U = r$. Thus, u_1, \dots, u_r is already a basis for V . Thus U and V are both spanned by u_1, \dots, u_r and $U = V$.

3. **Let A and B be $n \times n$ matrices. Does the null space of AB HAVE to be a subset of the null space of A ? If yes, prove your answer. If no, give an example.**

NO. Let

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \text{ and } B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

So,

$$AB = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

We see that $v = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is in the null space of AB , because $ABv = 0$; but v is not in the null space of A , because $Av = v \neq 0$.

4. **Define “null space”. Use complete sentences. Include everything that is necessary, but nothing more.**

The null space of the matrix A is the set of all column vectors x with $Ax = 0$.

5. **Define “dimension”. Use complete sentences. Include everything that is necessary, but nothing more.**

The dimension of the vector space V is the number of vectors in a basis for V .

6. **Let**

$$V = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \in \mathbb{R}^3 \mid 2x_1 + 3x_3 - 4x_3 = 5 \right\}.$$

Is V a vector space? Explain thoroughly.

NO. The vector $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ is not in V .

7. Let $a = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix}$ and $b = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$ be fixed elements of \mathbb{R}^3 , and let

$$V = \left\{ x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \in \mathbb{R}^3 \mid a^\top x = 0 \text{ and } b^\top x = 0 \right\}.$$

Is V a vector space? Explain thoroughly.

YES. The set V is the null space of

$$\begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{bmatrix}.$$

The null space of any matrix is a vector space.