

## 2.2 Completed Notes

### 2.2: Describing Sets

Definition: A set is any collection of objects with no repetitions. An object in a set is said to be an element of the set. One way to write a set is to list them in  $\{ \}$  with commas in between the elements.

Notation: If  $A$  is a set and  $a$  is an element of  $A$ , we write  $a \in A$ . If  $b$  is not an element of  $A$ , we write  $b \notin A$ .

Example: Write the set of the first five counting numbers and give examples of elements in and not in the set.

$$\{1, 2, 3, 4, 5\} = A$$

$$3 \in A \quad 1 \in A$$

$$6 \notin A \quad 0 \notin A \quad -3 \notin A$$

$$\sqrt{7} \notin A \quad \pi \notin A$$

Example: Describe the elements of the following sets.

(a)  $\{3x \mid x \in \mathbb{Z}\}$  all multiples of 3

$$\{\dots, -9, -6, -3, 0, 3, 6, 9, \dots\}$$

(b)  $\{-x \mid x \in \mathbb{N}\}$

$$\{-1, -2, -3, -4, \dots\}$$

(c)  $\{a/b \mid a, b \in \mathbb{Z}, b \neq 0\}$

all fractions (positive and negative)

(d)  $\{x^2 \mid x \in \mathbb{N}\}$

$$\{1^2, 2^2, 3^2, 4^2, \dots\} = \{1, 4, 9, 16, \dots\}$$

Example: Find the cardinal number of the following sets.

(a)  $S = \{1, 4, 7, 10, 13, \dots, 40\}$

$d = \text{common difference}$

$$n = \frac{\text{last \#} - \text{first \#}}{d} + 1$$

$$= \frac{40 - 1}{3} + 1$$

$$n(S) = \boxed{14}$$

(b)  $T = \{33, 37, 41, 45, 49, \dots, 353\}$

$$\frac{353 - 33}{4} + 1 = \frac{320}{4} + 1 = 81$$

$$n(T) = \boxed{81}$$

Definition: (Set builder notation) Let  $S$  be a set. Then we can write  $S = \{x \mid x \text{ satisfies some conditions}\}$ . This is read "S equals the set of elements  $x$  such that  $x$  satisfies some conditions".

Another way to think of set builder notation is  $\{\text{form of elements} \mid \text{conditions}\}$ . This will show up more in the examples.

Example: Write  $S = \{1, 2, 3, 4, 5\}$  in set builder notation.

$$\{x \mid x \text{ is one of the first 5 counting numbers}\}$$

Definition (Special Sets):

- (1) The Natural Numbers:  $\mathbb{N} = \{1, 2, 3, 4, \dots\}$
- (2) The Integers:  $\mathbb{Z} = \{\dots, -4, -3, -2, -1, 0, 1, 2, 3, 4, \dots\}$
- (3) The Real Numbers:  $\mathbb{R} = \{x \mid x \text{ is any number that can be written as a decimal}\}$

Definition: Two sets are equal if they contain exactly the same elements in any order.

Definition: The cardinal number of a set  $S$ , denoted  $n(S)$  or  $|S|$ , is the number of elements of  $S$ .

Definition: The empty set, denoted  $\emptyset$ , is the set with no elements. The empty set can also be written as  $\{ \}$ .

Definition: A set is a finite set if the cardinal number of the set is 0 or a natural number. A set with infinitely many elements, such as the natural numbers, is called an infinite set.

Example: Find the cardinal number of  $A = \{1, 2, 3, 4\}$ ,  $B = \{0\}$ ,  $C = \{2, 4, 6, 8, \dots\}$ , and  $\emptyset$ .


$$n(A) = 4 \quad n(B) = 1 \quad n(C) = \infty$$

$$n(\emptyset) = 0 \quad C \text{ is infinite}$$

Definition: The universal set, denoted  $U$ , is the set of all elements being considered in a given discussion.

Definition: The complement of a set  $S$ , denoted  $\bar{S}$ , is the set of all elements in  $U$  that are not in  $S$ . That is,  $\bar{S} = \{x \mid x \in U \text{ and } x \notin S\}$ .

A complement can be thought of in the following manner. The shaded region is  $\bar{S}$ .



Example: If  $U = \{1, 2, 3, 4, 5, 6, 7, 8\}$ , find the complements of  $A = \{2, 4, 6, 8\}$  and  $B = \{1, 2, 4, 6, 7\}$ .


$$\bar{A} = \{1, 3, 5\}$$

$$\bar{B} = \{3, 5, 8\}$$

## 2.2 Completed Notes

Definition: If  $A$  and  $B$  are sets, we say that  $A$  is a subset of  $B$ , denoted  $A \subseteq B$ , if every element of  $A$  is an element of  $B$ . If  $A \subseteq B$  and  $A \neq B$ , we say that  $A$  is a proper subset of  $B$ , denoted  $A \subset B$ .

A subset can be thought of in the following manner. In the figure  $A \subseteq B$ :



Example: Fill in the blanks with either  $\subseteq$  or  $\not\subseteq$ .

$\{1, 2, 3, 4\} \subseteq \{1, 2, 3, 4, 5\}$	$\{1, 2, 3, 4\} \subseteq \{1, 2, 3, 4\}$
$\{1, 2, 3, 4, 5\} \not\subseteq \{1, 2, 3, 4\}$	$\{1, 2, 3, 4, 5\} \not\subseteq \{1, 2, 3, 4, 6\}$
$\{0\} \not\subseteq \{1, 2, 3, 4\}$	$\emptyset \subseteq \{1, 2, 3, 4\}$
$\{1, 2, 3, 4\} \not\subseteq \emptyset$	$\emptyset \subseteq \emptyset$

Example: Fill in the blanks with either  $\in$ ,  $\notin$ ,  $\subseteq$ , or  $\not\subseteq$ .

$\{2\} \subseteq \{1, 2, 3\}$	$0 \notin \mathbb{N}$
$2 \in \{1, 2, 3\}$	$\mathbb{Z} \not\subseteq \mathbb{N}$ ← $\mathbb{Z}$ contains 0 and negatives that are not in $\mathbb{N}$
$5 \notin \{1, 2, 3, 4\}$	$5 \notin \{2x \mid x \in \mathbb{Z}\}$ ← all even integers
$\emptyset \subseteq \{1\}$	$\mathbb{R} \subseteq \mathbb{R}$
$0 \notin \emptyset$	$\{a/b \mid a, b \in \mathbb{Z}, b \neq 0\} \subseteq \mathbb{R}$
$\{4\} \not\subseteq \{2\}$	$\{1.5\} \not\subseteq \mathbb{N}$ ← fractions (+ or -)

First item: if it is a set, use  $\subseteq$  or  $\not\subseteq$   
if it is a number, use  $\in$  or  $\notin$