

2.2 Notes

2.2: Describing Sets

Definition: A set is any collection of objects with no repetitions. An object in a set is said to be an element of the set. One way to write a set is to list them in $\{ \}$ with commas in between the elements.

Notation: If A is a set and a is an element of A , we write $a \in A$. If b is not an element of A , we write $b \notin A$.

Example: Write the set of the first five counting numbers and give examples of elements in and not in the set.

Definition: (Set builder notation) Let S be a set. Then we can write $S = \{x \mid x \text{ satisfies some conditions}\}$. This is read 'S equals the set of elements x such that x satisfies some conditions'.

Another way to think of set builder notation is $\{\text{form of elements} \mid \text{conditions}\}$. This will show up more in the examples.

Example: Write $S = \{1, 2, 3, 4, 5\}$ in set builder notation.

Definition (Special Sets):

(1) The Natural Numbers: $\mathbb{N} = \{1, 2, 3, 4, \dots\}$

(2) The Integers: $\mathbb{Z} = \{\dots, -4, -3, -2, -1, 0, 1, 2, 3, 4, \dots\}$

(3) The Real Numbers: $\mathbb{R} = \{x \mid x \text{ is any number that can be written as a decimal}\}$

Example: Use set builder notation to write the set of all real numbers between 0 and 1.

Example: Use set builder notation to write the set of all even integers.

Example: Use set builder notation to write the set of perfect squares 1, 4, 9, 16, etc.

Example: Describe the elements of the following sets.

(a) $\{3x \mid x \in \mathbb{Z}\}$

(b) $\{-x \mid x \in \mathbb{N}\}$

(c) $\{a/b \mid a, b \in \mathbb{Z}, b \neq 0\}$

Definition: Two sets are equal if they contain exactly the same elements in any order.

Definition: The cardinal number of a set S , denoted $n(S)$ or $|S|$, is the number of elements of S .

Definition: The empty set, denoted \emptyset , is the set with no elements. The empty set can also be written as $\{ \}$.

Definition: A set is finite if the cardinal number of the set is 0 or a natural number. A set with infinitely many elements, such as the natural numbers, is called an infinite set.

Example: Find the cardinal number of $A = \{1, 2, 3, 4\}$, $B = \{0\}$, $C = \{2, 4, 6, 8, \dots\}$, and \emptyset .

2.2 Notes

Example: Find the cardinal number of the following sets.


(a) $S = \{1, 4, 7, 10, 13, \dots, 40\}$

(b) $T = \{33, 37, 41, 45, 49, \dots, 353\}$

Definition: The universal set, denoted U , is the set of all elements being considered in a given discussion.

Definition: The complement of a set S , denoted \bar{S} , is the set of all elements in U that are not in S . That is, $\bar{S} = \{x \mid x \in U \text{ and } x \notin S\}$.


A complement can be thought of in the following manner. The shaded region is \bar{S} :



Example: If $U = \{1, 2, 3, 4, 5, 6, 7, 8\}$, find the complements of $A = \{2, 4, 6, 8\}$ and $B = \{1, 2, 4, 6, 7\}$.

Definition: If A and B are sets, we say that A is a subset of B , denoted $A \subseteq B$, if every element of A is an element of B . If $A \subseteq B$ and $A \neq B$, we say that A is a proper subset of B , denoted $A \subset B$.

A subset can be thought of in the following manner. In the figure $A \subseteq B$:



Example: Fill in the blanks with either \subseteq or $\not\subseteq$.

$\{1, 2, 3, 4\} \underline{\hspace{1cm}} \{1, 2, 3, 4, 5\}$	$\{1, 2, 3, 4\} \underline{\hspace{1cm}} \{1, 2, 3, 4\}$
$\{1, 2, 3, 4, 5\} \underline{\hspace{1cm}} \{1, 2, 3, 4\}$	$\{1, 2, 3, 4, 5\} \underline{\hspace{1cm}} \{1, 2, 3, 4, 6\}$
$\{0\} \underline{\hspace{1cm}} \{1, 2, 3, 4\}$	$\emptyset \underline{\hspace{1cm}} \{1, 2, 3, 4\}$
$\{1, 2, 3, 4\} \underline{\hspace{1cm}} \emptyset$	$\emptyset \underline{\hspace{1cm}} \emptyset$

Example: Fill in the blanks with either \in , \notin , \subseteq , or $\not\subseteq$.

$\{2\} \underline{\hspace{1cm}} \{1, 2, 3\}$	$0 \underline{\hspace{1cm}} \mathbb{N}$
$2 \underline{\hspace{1cm}} \{1, 2, 3\}$	$\mathbb{Z} \underline{\hspace{1cm}} \mathbb{N}$
$5 \underline{\hspace{1cm}} \{1, 2, 3, 4\}$	$5 \underline{\hspace{1cm}} \{2x \mid x \in \mathbb{Z}\}$
$\emptyset \underline{\hspace{1cm}} \{1\}$	$\mathbb{R} \underline{\hspace{1cm}} \mathbb{R}$
$0 \underline{\hspace{1cm}} \emptyset$	$\{a/b \mid a, b \in \mathbb{Z}, b \neq 0\} \underline{\hspace{1cm}} \mathbb{R}$
$\{4\} \underline{\hspace{1cm}} \{2\}$	$\{1.5\} \underline{\hspace{1cm}} \mathbb{N}$