

ON THE VARIATIONAL DISTANCE OF TWO TREES*

BY M. A. STEEL AND L. A. SZÉKELY

University of Canterbury and University of South Carolina

A widely-studied model for generating sequences is to ‘evolve’ them on a tree according to a symmetric Markov process. We prove that model trees tend to be maximally “far apart” in terms of variational distance.

1. Introduction. In this paper we investigate sequences that have been generated on the tree by a simple Markov model. Such processes are widely-studied in molecular genetics, and in other areas of applied probability (including broadcasting and statistical physics). More precisely, we study the separation—as measured by variational distance—of the probability distribution on sequence patterns generated by different trees. We find that a large tree generates a probability distribution that is typically at maximal distance from that generated by nearly all other trees.

To describe our results more precisely we first provide some terminology concerning trees and random processes on them. In a tree, vertices of degree 1 are called *leaves*, as opposed to *internal vertices*. A tree is *binary*, if all vertices have degree 1 or 3. Consider a set X of labels. A *phylogenetic X -tree* is a tree, in which leaves are identified with elements of X . (We do not require phylogenetic X -trees be binary by definition for technical reasons, as we will have to deal with subtrees of phylogenetic X -trees.) We will regard two phylogenetic X -trees as being identical, if there is a graph isomorphism between them, which in addition, if restricted to X , is the identity function of X . If $|X| = n$, then the number of different binary phylogenetic X -trees is $(2n - 5)!!$ ($= 1 \times 3 \times 5 \times \cdots \times (2n - 5)$) [16]. For a phylogenetic X -tree

*We thank the NZIMA (Maclaurin Fellowship) for supporting this research. The second author was supported in part by NSF DMS contracts Nr. 0072187 and 0302307.

AMS 2000 subject classifications: Primary 62P10; secondary 05C05, 05C80, 05C90, 92D15

Keywords and phrases: Cavender–Farris–Neyman model, symmetric binary channel, tree-based Markov process, Yule–Harding distribution, phylogeny reconstruction, sequence evolution, q -state Potts model, Jukes–Cantor model, variational distance

\mathcal{T} , let $[\mathcal{T}]$ denote the corresponding unlabelled tree. The *distance* $d_{\mathcal{T}}(u, v)$ between two vertices, u, v in a tree \mathcal{T} is the number of edges on the unique path connecting them.

We now describe a model for the evolution of *binary* sequences on a tree. This model has been described by various authors (and in a range of disciplines including molecular biology, information theory, and physics; for references see [8, 16]). Here we refer to this model as the *CFN model* (short for ‘Cavender-Farris-Neyman model’); it has also been referred to in the literature as the ‘symmetric binary channel’ and the ‘symmetric 2-state Poisson model’. The CFN model provides a simple model for the evolution of purine–pyrimidine sequences. The significance of this simple model is, that phenomena shown for the CFN model often extends to more realistic models of sequence evolution, and we will describe how our main results concerning the CFN model generalise. The term *CFN tree* will refer to a phylogenetic X -tree equipped with a CFN model.

Suppose we have two states, 0 and 1, and a phylogenetic X -tree \mathcal{T} . The CFN model assigns probabilities to the patterns of state of the elements of X as follows. Let us associate a number p_e ($0 < p_e < 1/2$) with the edge e called the *transition probability*. Let ξ_e denote a random indicator variable associated to edge e with $\mathbb{P}[\xi_e = 1] = p_e$, and assume the ξ_e ’s are independent. Fix any vertex v and assign state 0 or 1 to v with equal probability $1/2$. Note that for every vertex u of \mathcal{T} there is a unique path denoted $path(u, v)$ in \mathcal{T} and so we may define

$$(1.1) \quad state(u) = state(v) + \sum_{e \in path(u, v)} \xi_e \pmod{2}.$$

This gives a (joint) probability distribution on the set of all assignment of states (0 or 1) to the vertices of \mathcal{T} , and thereby a marginal distribution on state assignments to the leaves of \mathcal{T} – we call each such assignment $\chi : X \rightarrow \{0, 1\}$ a (state) *pattern*, and we let \mathcal{P}_{χ} denote the probability of generating χ under this model.

The CFN model is thus specified by the pair $(\mathcal{T}, \mathcal{P})$, where \mathcal{P} is the map that associated to each edge e its transition probability $p(e)$. We refer to \mathcal{T} as a *CFN tree* and \mathcal{P} as a *transition mechanism*.

The probability p that the endpoints of a path uw in a CFN tree \mathcal{T} are in different states is nicely related to the transition probabilities of edges of

the uw -path:

$$(1.2) \quad p = \frac{1}{2} \left(1 - \prod_{e \in \text{path}(u,w)} (1 - 2p_e) \right).$$

Formula (1.2) is well-known, and is easy to prove by induction. Formula (1.2) also shows that the transition probability of a path is not less than the largest transition probability on its edges. It is well-known ([18]) that (1) changing the location of v in \mathcal{T} , or (2) substituting a path by a *single edge* in a CFN tree, and assigning to the new edge a transition probability according to (1.2) *does not change* the probability distribution of patterns.

Usually k independent experiments are made to generate random patterns from a binary CFN tree \mathcal{T} , they are called *sites*. The (abstract) *phylogeny reconstruction problem* is the following: from the observed pattern frequencies, determine with a prescribed probability, what was the underlying binary phylogenetic X -tree. We have shown in [6] that if $|X| = n$ and $n \rightarrow \infty$, then $k = \Omega(\log n)$ sites are needed to return the true underlying tree with probability at least $\frac{1}{2} + \epsilon$ with either a deterministic algorithm or with a randomized algorithm whose random bits are independent from the random events on the CFN tree. Sequence length requirements for accurate tree reconstruction is not only of mathematical interest, but also a topical issue in molecular systematics (eg. [3, 15]). We showed in [6] that for fixed $0 < f \leq g < 1/2$, $f \leq p_e \leq g$, and $n \rightarrow \infty$, phylogeny reconstruction is possible for all model trees, when k is a certain polynomial of n ; is possible for some model trees, when k is a logarithmic function of n ; and is possible for almost all model trees, either in the uniform random binary X -tree model or in the Yule–Harding model, when k is a certain polylogarithmic function of n . More recent work by E. Mossel and colleagues [5, 12] has established further instances for which logarithmic dependence of k on n suffices for accurate tree reconstruction and cases for which polynomial dependence is necessary.

In this paper we show *asymptotic results*. The theorems are about n -leaf trees, but their conclusions are $o(1)$ (limit) relations as $n \rightarrow \infty$. The understanding is that for a *sequence* of n -leaf trees satisfying the hypotheses, the limit relation holds. It would be technically more proper to speak about *sequences of trees* in the statements of the theorems, but we follow the tradition of random graph theory ([1, 4]) not speaking explicitly about sequences. With the exception of Section 4 we study problems where the bounds on p_e are *fixed*, and we let $n \rightarrow \infty$. In Section 4 we show that many of the results

generalize if dependence of the bounds on n is allowed but limited.

2. Results. Let us be given two binary phylogenetic X -trees $\mathcal{T}_1, \mathcal{T}_2$ with CFN transition mechanism \mathcal{P}_1 and \mathcal{P}_2 , respectively. The variational distance of their pattern distributions is

$$(2.1) \quad \text{vardist}\left((\mathcal{T}_1, \mathcal{P}_1), (\mathcal{T}_2, \mathcal{P}_2)\right) = \sum_x \left| (\mathcal{P}_1)_x - (\mathcal{P}_2)_x \right|.$$

This distance lies between 0 and 2, and in Theorem 3.1 we show that *almost all* binary trees are maximally distant (in terms of variational distance) from any given binary tree with a given CFN transition mechanism, under mild assumptions on their transition mechanisms. A practitioner may argue that Theorem 3.1 has limited relevance, since the uniform distribution of trees is just one particular prior distribution on trees, and the CFN model is very particular. However, the conclusion of Theorem 3.1 holds not just for the counting measure, but for all permutation invariant measures on phylogenetic X -trees; moreover it holds for more general, and for the applications more realistic classes of transition mechanisms (Theorem 4.1). This result may not be surprising: as we equip randomly selected trees with CFN models, they have many local statistics that are essentially independent and have different marginals in the two trees. Therefore, analogously to the Kakutani Dichotomy, their measures are expected to be (near) orthogonal.

Farach and Kannan [9], [10] designed an algorithm for phylogeny reconstruction based on convergence to the true tree in variational distance and suggested to pay more attention to the variational distance in phylogeny reconstruction. Some support for the utility of this metric is provided by results that we present in Sections 3 and 4: if we get just *close* to a model tree in variational distance, then we already excluded most of the false candidates for the phylogenetic tree.

However, a simple fact provides a sharp contrast to the results mentioned above. Note that in practice we estimate the model distribution of patterns by the *observed frequency* of patterns. For *sub-exponential* sequence length, which is known to be sufficient for phylogeny reconstruction with probability $1 - o(1)$ as $0 < f \leq g < 1/2$ fixed and $f \leq p_e \leq g$, as $n \rightarrow \infty$ (see the discussion in Section 1), the variational distance between the model pattern distribution and the observed pattern distribution is near 2 with probability $1 - o(1)$. (For details, see our technical report [19].)

In other words, phylogeny reconstruction is well possible *without* convergence of the observed pattern distribution to the model pattern distribution in variational distance.

Therefore the accuracy of tree reconstruction cannot be captured by variational distance alone. This conclusion was suggested by [7] and [14], though with less explicit theoretical justification.

3. Variational distance of CFN trees is typically large.

THEOREM 3.1. *Fix $0 < f$ and $g < 1/2$. There exists a function $\epsilon(n) = \epsilon_{f,g}(n) = o(1)$ as $n \rightarrow \infty$, such that for every binary phylogenetic X -tree \mathcal{T}_1 with CFN transition mechanism \mathcal{P}_1 where $p_e \leq g$ in \mathcal{P}_1 , the following holds: For almost all (i.e. $(1 - o(1))(2n - 5)!!$ in number) binary phylogenetic X -trees \mathcal{T}_2 , equipped with an arbitrary transition mechanism \mathcal{P}_2 , where $f \leq p_e$ in \mathcal{P}_2 , we have*

$$(3.1) \quad \text{vardist}\left((\mathcal{T}_1, \mathcal{P}_1), (\mathcal{T}_2, \mathcal{P}_2)\right) \geq 2 - \epsilon(n).$$

The proof requires a number of lemmas, which we now state.

LEMMA 3.2. *For every binary phylogenetic X -tree \mathcal{T} on $n \geq 4$ leaves, there are at least $n/4$ disjoint pairs of leaves a_i, b_i , such that for every i :*

- (i) a_i and b_i are separated by a distance of 2 or 3;
- (ii) for $i \neq j$, the $a_i b_i$ and the $a_j b_j$ paths in \mathcal{T} are edge disjoint.

PROOF. The claim is true for $4 \leq n \leq 8$, since then any longest path ends in two disjoint cherries. This is the basis for an induction proof on n . It is easy to see that, for $n \geq 9$, there exists a longest path in \mathcal{T} , for which one end must be a leaf in a cherry that lies at the top portion of the tree given by one of the four cases shown in Fig. 1 (the other end of the path lies in the bottom part of the tree, represented by a circle). In each of the four cases truncate the tree T_i as indicated by the dashed curve to obtain T'_i . For $i = 1, 2, 3, 4$, T'_i has $n - 2$ (resp. $n - 2, n - 3, n - 4$) leaves, and the induction hypothesis applies to T'_i . In all four cases it is easy to add two new close vertex pairs to create the required set of them for T_i , while destroying at most one which pre-existed in T'_i . \square





FIG 1. Ending of a longest path in a binary tree.

FIG 2. Binary tree on $4t + 9$ leaves, with only $t + 3$ close leaf pairs.

REMARK 3.3. As Fig. 2 shows, the conclusion of Lemma 3.2 is essentially the best possible.

LEMMA 3.4. Tree-chopping lemma [Steel, Goldstein, and Waterman [17] Lemma 3]

Let \mathcal{T} be an arbitrary binary X -tree and $q \geq 2$ integer. Then edges can be deleted from \mathcal{T} such that a forest results with the following properties:

- (i) The number of leaves from X in any tree of the forest is at most $2q - 2$.
- (ii) The number of leaves from X in any tree of the forest is at least q , except possibly for one tree. (We shall call this exceptional tree degenerate.)

erate.)

Recall the Azuma–Hoeffding inequality (see [1]):

LEMMA 3.5. *Suppose $\mathbf{X} = (X_1, X_2, \dots, X_k)$ are independent random variables taking values in any set S , and $L : S^k \rightarrow \mathbb{R}$ is any function that satisfies the condition: $|L(\mathbf{u}) - L(\mathbf{v})| \leq t$ whenever \mathbf{u} and \mathbf{v} differ at just one coordinate. Then,*

$$(3.2) \quad \mathbb{P}[|L(\mathbf{X}) - \mathbb{E}[L(\mathbf{X})]| \geq \lambda] \leq 2 \exp\left(-\frac{\lambda^2}{2t^2k}\right). \quad \square$$

The following lemma is obvious.

LEMMA 3.6. *Let \mathcal{F} denote a fixed phylogenetic X -tree, with $|X| = n$, and let $\tau = [\mathcal{F}]$ (the corresponding unlabelled tree). Let π be a randomly selected permutation of X under the uniform distribution. Let $\pi(\mathcal{F})$ denote the phylogenetic X -tree that we obtain from \mathcal{F} by changing all leaf labels from v to $\pi(v)$ simultaneously. Then $\pi(\mathcal{F})$ represents a random uniform selection from those binary phylogenetic X -trees whose underlying unlabelled tree is τ . \square*

From now on, for notational convenience, we pretend that 4 divides n .

LEMMA 3.7. *For an X with $|X| = n$, and $n/4$ disjoint a_i, b_i ordered pairs from X , there exist functions $m(n) \rightarrow \infty$, $h(n) \rightarrow \infty$, and $g(n) \rightarrow \infty$, such that the following holds. For every unlabelled binary tree τ with n leaves, for all but a $\frac{1}{g(n)}$ fraction of binary phylogenetic X -trees \mathcal{T} with property $[\mathcal{T}] = \tau$, there is an index set I such that $|I| = m(n)$ and*

- (i) $d_{\mathcal{T}}(a_i, b_i) \geq h(n)$ for all $i \in I$; and
- (ii) for $i, j \in I$, $i \neq j$, $\text{path}_{\mathcal{T}}(a_i, b_i)$ and $\text{path}_{\mathcal{T}}(a_j, b_j)$ are edge disjoint.

PROOF. Let \mathcal{F} denote a fixed binary phylogenetic X -tree such that $[\mathcal{F}] = \tau$, with $|X| = n$. Apply Lemma 3.4 to \mathcal{F} with $q = \lceil \log^2 n \rceil$. Let L_1, L_2, \dots, L_s denote the leaf sets that the non-degenerate trees contain from X . From

the lemma, $q \leq |L_i| \leq 2q - 2$, and at most $q - 1$ element of X are not in some L_i . Let π be a randomly selected permutation of X under the uniform distribution. Let $\pi(\mathcal{F})$ denote the phylogenetic X -tree that we obtain from \mathcal{F} by changing all leaf labels from v to $\pi(v)$ simultaneously. According to Lemma 3.6, $\pi(\mathcal{F})$ represents a random uniform selection from those binary phylogenetic X -trees whose underlying unlabelled tree is τ . The previous application of Lemma 3.4 still partitions $\pi(\mathcal{F})$, the leaf sets of the non-degenerate trees intersect X in $\pi(L_1), \pi(L_2), \dots, \pi(L_s)$, and we still have $q \leq |\pi(L_i)| \leq 2q - 2$. Therefore, for $i \neq j$, if $path_{\pi(\mathcal{F})}^i$ (respectively, $path_{\pi(\mathcal{F})}^j$) connects an arbitrary pair of vertices of L_i (respectively L_j) in the tree $\pi(\mathcal{F})$, then

$$(3.3) \quad path_{\pi(\mathcal{F})}^i \text{ is edge disjoint from } path_{\pi(\mathcal{F})}^j.$$

Set $h(n) = \log \log n$ and $m(n) = \frac{n}{4q-2} - \frac{1}{2}$. Observe from Lemma 3.4 and the choice of q that $n \leq (s+1)(2q-2)$, and therefore

$$(3.4) \quad m(n) \leq \frac{s}{2}.$$

We are going to find an appropriate $g(n)$ for this choice. We call a leaf set $Y \subset X$ *infected*, if there is a $1 \leq j \leq n/4$, such that *both* $a_j, b_j \in Y$. Let E denote the event that for our fixed τ and \mathcal{F} , $\pi(\mathcal{F})$ has the property that for all $j = 1, 2, \dots, s$, $\pi(L_j)$ is infected; and let F denote the event that in addition to E , for at least half of the indices $j = 1, 2, \dots, s$, one finds some i_j , such that both $a_{i_j}, b_{i_j} \in \pi(L_j)$ (i.e. they do infect $\pi(L_j)$) and $d_{\pi(\mathcal{F})}(a_{i_j}, b_{i_j}) \geq h(n)$. In view of (3.3), the a_{i_j}, b_{i_j} paths in $\pi(\mathcal{F})$ are pairwise edge disjoint for $j = 1, 2, \dots, s$.

Observe that

$$(3.5) \quad \mathbb{P}[\pi(L_j) \text{ not infected}] = \frac{\sum_{u=0}^{|L_j|} \binom{n/4}{u} 2^u \binom{n/2}{|L_j|-u}}{\binom{n}{|L_j|}}.$$

(A non-infected L_j can have zero or one element from every (a_i, b_i) pair, for $i = 1, 2, \dots, n/4$. The case analysis is based on the number $u = |\pi(L_j) \cap \{a_i, b_i : i = 1, 2, \dots, n/4\}|$. There are $\binom{n/4}{u}$ to select a subset of u indices from $\{1, 2, \dots, n/4\}$, and then 2^u ways to tell if a_i or b_i selected for the particular index set into L_j . There are $\binom{n/2}{|L_j|-u}$ ways to make L_j complete using $|L_j| - u$ elements not belonging to $\{a_i, b_i : i = 1, 2, \dots, n/4\}$.)

Comparison of consecutive terms show that the largest term in the numerator of the RHS of (3.5) is $u = |L_j|$. Using the usual notation $(x)_m$ for

the m^{th} falling factorial, it follows that

$$(3.6) \quad \mathbb{P}[\pi(L_j) \text{ not infected}] \leq \frac{(|L_j| + 1)2^{|L_j|} \binom{n/4}{|L_j|}}{\binom{n}{|L_j|}}$$

$$(3.7) \quad = \frac{(n/4)^{|L_j|}}{(n)^{|L_j|}} (|L_j| + 1)2^{|L_j|} \leq \frac{n^{|L_j|}}{4^{|L_j|} (n - |L_j|)^{|L_j|}} (|L_j| + 1)2^{|L_j|}$$

$$(3.8) \quad \leq (1 + o(1))2^{-|L_j|} (|L_j| + 1) \leq (1 + o(1))2^{-q} (2q - 1) \leq 2^{-.01 \log^2 n},$$

and from (3.6-3.7-3.8),

$$(3.9) \quad \mathbb{P}[\exists j : \pi(L_j) \text{ not infected}] \leq \frac{n}{q} 2^{-.01 \log^2 n}.$$

By (3.9), we showed that

$$(3.10) \quad \mathbb{P}[E] > 1 - n2^{-.01 \log^2 n}.$$

Call the ordered s -tuple of pairwise disjoint sets $Y_1, Y_2, \dots, Y_s \subset X$ *feasible*, if $|Y_i| = |L_i|$ and Y_i is infected for $i = 1, 2, \dots, s$. Now we turn to the conditional probability $\mathbb{P}[F|E]$. Observe

$$(3.11) \quad \mathbb{P}[F|E] = \sum_{Y_1, Y_2, \dots, Y_s \text{ feasible}} \mathbb{P}\left[F|\forall i : \pi(L_i) = Y_i\right] \mathbb{P}\left[\forall i : \pi(L_i) = Y_i\right]$$

$$(3.12) \quad \leq \max_{Y_1, Y_2, \dots, Y_s \text{ feasible}} \mathbb{P}\left[F|\forall i : \pi(L_i) = Y_i\right].$$

Assume now that an arbitrary feasible Y_1, Y_2, \dots, Y_s is *fixed*. A π that satisfies the condition in (3.12) is nothing else but the juxtaposition of $\pi_i : L_i \rightarrow Y_i$ bijections for $i = 1, 2, \dots, s + 1$. Therefore a uniform random π satisfying the condition in (3.12) can be realized by a sequence of *independent* uniform random choices of bijections π_i from L_i to Y_i , $i = 1, 2, \dots, s + 1$.

Let $\pi_i : L_i \rightarrow Y_i$ denote a uniform random bijection for $i = 1, 2, \dots, s + 1$. Conditional on E , for every $i = 1, 2, \dots, s$, fix an a_{i_j}, b_{i_j} leaf pair that infects Y_i . Observe that the conditional event

$$F|\forall i : \pi(L_i) = Y_i$$

is implied, if for at least half of the indices $1 \leq i \leq s$, we have $d_{\pi(\mathcal{F})}(a_{i_j}, b_{i_j}) \geq h(n)$. Also observe that notwithstanding the notation $d_{\pi(\mathcal{F})}$, this distance

depends *only* on the *single* π_i under consideration. No matter what is the value of $\pi_i^{-1}(a_{i_j})$, at most $2^{h(n)}$ vertices of L_i can be closer than $h(n)$ to $\pi_i^{-1}(a_{i_j})$ in the binary tree \mathcal{F} . Those at most $2^{h(n)}$ vertices can be pre-images of b_{i_j} under π_i (and π as well), if $d_{\pi(\mathcal{F})}(a_{i_j}, b_{i_j}) < h(n)$. Therefore,

$$\mathbb{P}\left[d_{\pi(\mathcal{F})}(a_{i_j}, b_{i_j}) \geq h(n)\right] \geq 1 - \frac{2^{h(n)}}{|L_i|} = 1 - \frac{2^{\log \log n}}{\log^2 n} = 1 - \frac{1}{\log^{2-\log 2} n}.$$

Hence, a lower bound for $\mathbb{P}[F|E]$ is the probability of at least $s/2$ successes in a sequence of s independent Bernoulli trials, each with probability of success $p = 1 - \frac{1}{\log^{2-\log 2} n}$. Not having at least $m(n)$ successes implies not having at least $s/2$ successes by (3.4), and probability of the latter event can easily be bounded from above by Lemma 3.5 ($t = 1$, $k = s$, $\lambda = s/3$), as soon as $\frac{1}{\log^{2-\log 2} n} < 1/6$, by

$$(3.13) \quad 2e^{-s/18}.$$

Finally, using (3.10) and (3.13), we have

$$(3.14) \quad 1 - \mathbb{P}[F] = 1 - \mathbb{P}[E] + \mathbb{P}[E](1 - \mathbb{P}[F|E]) \leq n2^{-.01 \log^2 n} + 2e^{-n/(64 \log^2 n)},$$

and since the RHS of (3.14) is $o(n)$, we can take for $g(n)$ its reciprocal. \square

Proof of Theorem 3.1 Specify now $n/4$ leaf pairs $\{a_i, b_i\}$ of \mathcal{T}_1 according to Lemma 3.2—for notational convenience we assume again that n is a multiple of 4. We set $m(n)$, $h(n)$, $g(n)$, and I according to the statement of Lemma 3.7. We are going to show that for every fixed $(\mathcal{T}_1, \mathcal{P}_1)$ and fixed unlabelled tree τ , if $[\mathcal{T}_2] = \tau$ and \mathcal{T}_2 is not in the exceptional set of trees described in Lemma 3.7, then the variational distance between $(\mathcal{T}_1, \mathcal{P}_1)$ and $(\mathcal{T}_2, \mathcal{P}_2)$ differs from 2 by at most an quantity that is $o(1)$ as a function of n . Recall that $state(x)$ denotes the state of leaf $x \in X$ in a CFN tree. Consider the random indicator variable Z_i , which is 1, if $state(a_i) = state(b_i)$, and 0 otherwise, and $Z = \sum_{i \in I} Z_i$, which depends on the distribution of leaf colorations of the CFN tree. We will speak about $Z_i^{(1)}$, $Z^{(1)}$ and $Z_i^{(2)}$, $Z^{(2)}$ as the CFN tree is $(\mathcal{T}_1, \mathcal{P}_1)$ or $(\mathcal{T}_2, \mathcal{P}_2)$, and similarly about $state_1$ and $state_2$, and will drop the superscript if the argument applies to both.

By the linearity of expectation

$$(3.15) \quad \mathbb{E}[Z] = \sum_{i \in I} \mathbb{E}[Z_i] = \sum_{i \in I} \mathbb{P}[state(a_i) = state(b_i)].$$

In $(\mathcal{T}_1, \mathcal{P}_1)$, we have $\mathbb{P}[\text{state}_1(a_i) \neq \text{state}_1(b_i)] \leq \frac{1}{2} \left(1 - (1 - 2g)^3\right)$, by (1.2), and hence

$$(3.16) \quad 1 - 3g + 6g^2 - 4g^3 \leq \mathbb{P}[\text{state}_1(a_i) = \text{state}_1(b_i)]$$

Formula (3.15) and inequality (3.16) imply that

$$(3.17) \quad \mathbb{E}[Z^{(1)}] \geq (1 - 3g + 6g^2 - 4g^3)m(n)$$

In $(\mathcal{T}_2, \mathcal{P}_2)$, by a similar argument, we have

$$(3.18) \quad \mathbb{P}[\text{state}_2(a_i) = \text{state}_2(b_i)] \leq 1 - \frac{1}{2} \left(1 - (1 - 2f)^{h(n)}\right) = \frac{1}{2} + o(1)$$

by (1.2), and $h(n) \rightarrow \infty$. By linearity (3.15), we have

$$(3.19) \quad \mathbb{E}[Z^{(2)}] \leq (1 + o(1)) \frac{m(n)}{2}.$$

We are going to show that with high probability both $Z^{(1)}$ and $Z^{(2)}$ are very close to their respective expectations. This will be easy to show, since both of them are the sums of independent indicator variables. (Use Lemma 3.5 for $X_i = Z_i^{(1)}$ (resp. $Z_i^{(2)}$), $k = m(n)$, $t = 1$, $\lambda = m(n)^{2/3}$.)

It is easy to see that for $0 < g < 1/2$, we have

$$(3.20) \quad 1/2 < 1 - 3g + 6g^2 - 4g^3,$$

and therefore, using (3.17) and (3.19), $\mathbb{E}[Z^{(1)}]$ and $\mathbb{E}[Z^{(2)}]$ are separated by a linear function of $m(n)$, for example $l(n) = \frac{1}{2}(1 - 3g + 6g^2 - 4g^3 + \frac{1}{2})m(n)$. Consider now the event H : “ $Z > l(n)$ ”. In $(\mathcal{T}_1, \mathcal{P}_1)$, event H has probability $1 - o(1)$, while in $(\mathcal{T}_2, \mathcal{P}_2)$, the complement of event H has probability $1 - o(1)$. This implies that the variational distance of $(\mathcal{T}_1, \mathcal{P}_1)$ and $(\mathcal{T}_2, \mathcal{P}_2)$ is $2 - o(1)$.

□

4. Variational distance in more general models. In this section we provide a result (Theorem 4.1) that is a three-fold generalization of Theorem 3.1. The three extensions allow (i) more general probability distributions on trees (‘permutation-invariant measures’), (ii) more general transition models than the CFN model (‘conservative, separable processes’) and (iii) a weakening of the constraints on the parameters of the model.

Permutation-invariant measures on trees. Let us call a measure μ on the set of $(2n-5)!!$ binary phylogenetic X -trees *permutation invariant*, if for every π permutation of X and any phylogenetic X -tree \mathcal{F} , $\mu(\mathcal{F}) = \mu(\pi(\mathcal{F}))$. Note that Lemma 3.6 stated that the uniform distribution (or counting measure) on binary phylogenetic X -trees is permutation invariant. A practitioner may argue that Theorem 3.1 has limited relevance, since the uniform distribution of trees is just one particular prior distribution on trees. However, any relevant distribution of trees is permutation invariant and it is easy to see that the stronger Theorem 3.1 holds with basically the same proof. A non-uniform, phylogenetically relevant permutation invariant distribution on phylogenetic X -trees is the *unrooted Yule-Harding distribution* [6].

More general transition processes (conservative, separable processes). The restriction of the CFN to two states and symmetric transition probabilities is convenient for description and proofs. However much of the argument used in the proof of Theorem 3.1 can be generalized to models that are much closer to those used in modern molecular biology. We identify two key properties that are used in the proof, and that both apply to a range of substitution models.

Suppose we have a set S of $q \geq 2$ states. A *pattern* will now refer to a state assignment function $\chi : X \rightarrow S$, where X is the leaf set of \mathcal{T} . Assume that we have a probability distribution on the patterns of a binary phylogenetic X -tree, where \mathcal{P}_χ denotes the probability of pattern χ . Selecting a random pattern according to the distribution, we can observe a random *state* of any particular leaf. For a pair of leaves a, b let $E(a, b)$ be the event that $state(a) = state(b)$. Let us be given a strictly decreasing function $H : [0, \infty) \rightarrow (c, 1]$ with $H(0) = 1$, and a $c > 0$ constant, such that $\lim_{x \rightarrow \infty} H(x) = c$. We assume that H and c are fixed and do not depend on n . We say that a probability distribution on patterns is *conservative* if

- (C) there exists an assignment of $t(e) > 0$ to each edge e of \mathcal{T} ,
so that the following condition holds: For each pair $a, b \in X$,
 $\mathbb{P}[E(a, b)] = H(\sum_{e \in \text{path}(a, b)} t(e))$.

The CFN model satisfies condition (C), as can easily be seen from (1.2) by taking $t(e) = -\frac{1}{2} \log(1 - 2p_e)$, $H(x) = \frac{1}{2}(1 + \exp(-2x))$, and $c = \frac{1}{2}$. More generally, condition (C) is satisfied by any tree-based Markov process that can be realised by a stationary, reversible, continuous-time Markov process operating on each edge e of \mathcal{T} for a duration (corresponding to $t(e)$) (this is

Theorem 4(2) of [18]; for more details on such models see [16]).

Next, we say that a probability distribution on patterns is *separable* if it satisfies the following property:

- (S) Whenever $(a_1, b_1), (a_1, b_2), \dots, (a_m, b_m)$ are pairs of leaves whose connecting paths are pairwise edge-disjoint then $\{E(a_i, b_i), i = 1, \dots, m\}$ are independent events.

It is easily seen that the CFN model is separable. Moreover, any group-based model satisfies the separation condition (S) (Theorem 10 of [21], generalizing [11]); briefly, ‘group-based models’ are defined in the same way as the CFN model, but over an arbitrary finite abelian group, rather than the particular group $(\{0, 1\}, +_{\text{mod}2})$ (for more details see [16]).

We will call a model that satisfies conditions (C) and (S) a *conservative, separable process*. Examples of such models include the CFN model, and more generally the symmetric q -state model, for which, when a transition occurs, one of the remaining states is selected uniformly at random. For this model we have $c = \frac{1}{q}$ in condition (C), and this model is well-known in a variety of field, including physics, broadcasting and molecular biology, where it is referred to as the ‘ q -state Potts model’, the ‘ q -ary symmetric channel’, and the ‘Neyman q -state model’, respectively (and, in the special case when $q = 4$, as the ‘Jukes-Cantor model’); for more details see [13]. A further example of a conservative, separable process in molecular biology is the Kimura 3ST model (for details see [16]).

Weakened constraints. In Theorem 3.1 we imposed the condition $f \leq p_e$ for a fixed $f > 0$ for the transition mechanism \mathcal{P}_2 . In fact an inspection of the proof reveals that $0 < f = f(n)$ may depend on n , as far as we have $\lim_{n \rightarrow \infty} h(n)f(n) = \infty$, where $h(n)$ is any function satisfying the statement of Lemma 3.7. (The present proof of Lemma 3.7 allows $f(n) \rightarrow 0$ “very slowly”, but the truth is likely just “slowly”.)

The result allowing these three types of extension is the following.

THEOREM 4.1. *Fix $0 < t_+ < \infty$, and allow $t_- = t_-(n) > 0$ to vary with n if still $\lim_{n \rightarrow \infty} h(n)t_-(n) = \infty$, where $h(n)$ is any function satisfying the statement of Lemma 3.7. For every binary phylogenetic X -tree \mathcal{T}_1 with a conservative, separable process \mathcal{P}_1 where $t(e) \leq t_+$ in \mathcal{P}_1 , and any μ permutation invariant measure on phylogenetic X -trees, the following holds for*

a function $\epsilon(n) = o(1)$. The set of binary phylogenetic X -trees of measure $1 - o(1)$ has the property, that any of them equipped with an arbitrary conservative, separable process \mathcal{P}_2 , with $t(e) \geq t_-$ in \mathcal{P}_2 (assuming \mathcal{P}_2 has the same H and c as \mathcal{P}_1) has

$$(4.1) \quad \text{vardist}\left((\mathcal{T}_1, \mathcal{P}_1), (\mathcal{T}_2, \mathcal{P}_2)\right) \geq 2 - \epsilon(n).$$

PROOF. We need a straightforward modification of the proof of Theorem 3.1. Leaving out the subscript from the notation for the generic leaf pair (a_i, b_i) , formula (3.16) can be substituted by

$$(4.2) \quad H(3t_+) \leq H\left(d_{\mathcal{T}_1}(a, b)t_+\right) \leq \mathbb{P}[\text{state}_1(a) = \text{state}_1(b)];$$

(3.18) can be substituted by

$$(4.3) \quad \mathbb{P}[\text{state}_2(a) = \text{state}_2(b)] \leq H\left(d_{\mathcal{T}_2}(a, b)t_-\right) \leq H\left(h(n)t_-\right) < c + \epsilon$$

for any fixed $\epsilon > 0$ as $n \rightarrow \infty$. For a sufficiently small $\epsilon > 0$, we have

$$(4.4) \quad c + \epsilon < H(3t_+)$$

(this follows from the assumptions on H and c), and thus inequality (4.4) substitutes for (3.20). \square

Acknowledgement. We thank Éva Czabarka for her careful reading of earlier versions of this manuscript, and the referees for helpful comments.

REFERENCES

- [1] ALON, N. and SPENCER, J. H. (1992). *The Probabilistic Method*, John Wiley and Sons, New York. [MR1140703](#)
- [2] AMBIANIS, A., DESPER, R., FARACH, M. and KANNAN, S., (1997) Nearly tight bounds on the learnability of evolution, *Proc. 38th Annual Symposium on Foundations of Computer Science (FOCS'97)* 524–533.
- [3] BININDA-ÉMONDS, O. R. P., BRADY, S. G., KIM, J. and SANDERSON, M. J. (2001) Scaling of accuracy in extremely large phylogenetic trees. *Pacific Symposium on Bio-computing* **6** 547–558.
- [4] BOLLOBÁS, B. *Random graphs*. Second edition. Cambridge Studies in Advanced Mathematics, 73. Cambridge University Press, Cambridge, 2001. [MR1864966](#)

- [5] DASKALAKIS, C., MOSSEL, E., ROCH, S. (2005) Optimal Phylogenetic Reconstruction Manuscript (ArXiv math.PR/0509575)
- [6] ERDŐS, P. L., STEEL, M. A., SZÉKELY, L. A. and WARNOW, T. J. (1999) A few logs suffice to build (almost) all trees I, *Random Structures and Algorithms* **14** (2) 153–184. [MR1667319](#)
- [7] ERDŐS, P.L., STEEL, M.A., SZÉKELY, L.A. and WARNOW, T. A few logs suffice to build (almost) all trees (Part II) *Theoretical Computer Science* **221** 77–118. [MR1700821](#)
- [8] EVANS, W., KENYON, C., PERES, Y. and SCHULMAN, L. J. (2000) Broadcasting on trees and the Ising model. *Adv. Appl. Prob.* **10** 410–433. [MR1768240](#)
- [9] FARACH, M. and KANNAN, S. (1996) Efficient algorithms for inverting evolution, *Proceedings of the ACM Symposium on the Foundations of Computer Science*, 230–236. [MR1427518](#)
- [10] FARACH, M. and KANNAN, S. (1999) Efficient algorithms for inverting evolution, *J. ACM* **46**(4) 437–449. [MR1812126](#)
- [11] FU, Y.-X. and LI, W.-H. (1991) Necessary and sufficient conditions for the existence of certain quadratic invariants under a phylogenetic tree. *Math. Biosci.* **105** 229–238.
- [12] MOSSEL, E. (2004) Phase transitions in Phylogeny Mossel, E. *Trans. Amer. Math. Soc.* **356**(6) 2379–2404.
- [13] MOSSEL, E. and PERES, Y. (2003) Information flow on trees. *Annals of Applied Probability* **13** (3) 817–844. [MR1994038](#)
- [14] RICE, K. and WARNOW, T. Parsimony is hard to beat, *COCOON'97. (Computing and Combinatorics, Third Annual International Conference)*, Shanghai, August 20–22, 1997, Tao Jiang and D.T. Lee, (Eds.). Lecture Notes in Computer Science Vol. **1276**, Springer-Verlag 124–133. [MR1616310](#)
- [15] ROKAS, A. and CARROLL, S. B. (2005) More Genes or More Taxa? The Relative Contribution of Gene Number and Taxon Number to Phylogenetic Accuracy. *Mol. Biol. Evol.* **22**(5) 1337–1344.
- [16] SEMPLE, C. and STEEL, M. (2003) *Phylogenetics*, Oxford University Press. [MR2060009](#)
- [17] STEEL, M. A., GOLDSTEIN L. and WATERMAN, M. S. A central limit theorem for the parsimony length of trees, (1996) *Adv. Appl. Prob.* **28**, 1051–1071. [MR1418246](#)
- [18] STEEL, M. A., HENDY, M. D. and PENNY, D. (1998) Reconstructing phylogenies from nucleotide pattern frequencies - a survey and some new results, *Discrete Applied Mathematics* **88** 367–396. [MR1658533](#)
- [19] STEEL, M. and SZÉKELY, L. A. On the variational distance of two trees, Technical Report 05:09, Industrial Mathematics Institute, University of South Carolina, <http://www.math.sc.edu/~IMI/technical/tech05.html>)
- [20] STEEL, M. A. and SZÉKELY, L. A. (2002) Inverting random functions II: explicit bounds for the discrete maximum likelihood estimation, with applications, *SIAM J. Discrete Math.* **15**(4) 562–575. [MR1935839](#)
- [21] TUFFLEY, C. and STEEL, M.A. (1997) Modelling the covarion hypothesis of nucleotide substitution, *Mathematical Biosciences* **147** 63–91. [MR1604518](#)

M. A. STEEL
 BIOMATHEMATICS RESEARCH CENTRE,
 DEPARTMENT OF MATHEMATICS AND STATISTICS,
 UNIVERSITY OF CANTERBURY,
 CHRISTCHURCH, NEW ZEALAND.
 E-MAIL: m.steel@math.canterbury.ac.nz

L. A. SZÉKELY
 DEPARTMENT OF MATHEMATICS,
 UNIVERSITY OF SOUTH CAROLINA
 COLUMBIA SC, USA.
 E-MAIL: szekely@math.sc.edu