

Exam 1
8 February 2013

Name: _____

Instructions:

1. There are a total of 5 problems on 3 pages. Check that your copy of the exam has all of the problems.
2. No electronic or other inanimate objects can be used during this exam. All questions have been designed with this in mind and should not involve unreasonable manual calculations.
3. Be sure you answer the questions that are asked.
4. You must show all of your work to receive full credit for a correct answer. Correct answers with no supporting work will be eligible for at most half-credit.
5. Your answers must be clearly labeled and written legibly on additional sheets of paper (that I will provide). Be sure each sheet contains your name and the work for each question is clearly labeled.
6. Check your work. If I see *clear evidence* that you checked your answer (when possible) and you *clearly indicate* that your answer is incorrect, you will be eligible for more points than if you had not checked your work.





Problem	Points	Score
1	15	
2	20	
3	32	
4	21	
5	12	
Total	100	

Good Luck!

- (15 points) Draw the direction line for the differential equation $y' = (y + 1)^2 e^{y-2}$. Use this information to determine the behavior of y as $t \rightarrow \infty$. If this behavior depends on the initial value of y at $t = 0$, describe this dependency.
- (20 points) Consider the following list of differential equations.

(i) $y' = t + y$	(ii) $y' = t - y$	(iii) $y' = 0.2t^2 + y$	(iv) $y' = y(1 - y/2)$
(v) $y' = y^2 - t^2$	(vi) $y' = t^2 - y^2$	(vii) $y' = 1 - ty$	(viii) $y' = \sin(t) \cos(t)$

Identify the differential equation that corresponds to each direction field.



3. (32 points) Find the general solution of each differential equation. If possible, solve for y . If an initial condition is given, find the solution to the initial value problem. For each differential equation that is nonlinear determine the interval on which the solution exists.

(a) $t^2y' + t(t+2)y = e^t$

(b) $\frac{dy}{dt} = e^{3t+2y}, y(0) = 0$

4. (21 points) Determine if the existence and uniqueness theorem applies. If it applies, determine the largest interval on which the solution of the initial value problem is certain to exist. If it does not apply, explain why not. *In no case should you waste any time trying to find a solution to the initial value problem.*

(a) $\frac{dy}{dt} + \frac{1}{t^2 - 6t + 8}y = \sqrt{t+1}, y(0) = 3$

(b) $\frac{dy}{dt} + \frac{1}{t^2 - 6t + 8}y = \sqrt{t+1}, y(2) = 1$

(c) $\frac{dy}{dt} + \frac{1}{t^2 - 6t + 8}y = \sqrt{t+1}, y(5) = -3$

5. (12 points) Find the value of b for which the differential equation

$$6xy^3 + \cos(y) + (3bx^2y^2 - x \sin(y))\frac{dy}{dx} = 0$$

is exact. Then solve the equation (with this value of b).