

**Complex Variables Notes for Math 703.**

**Part I**

**Updated Fall 2011**

Anton R. Schep



## Holomorphic (or Analytic) Functions

### 1. Definitions and elementary properties

In complex analysis we study functions  $f : S \rightarrow \mathbb{C}$ , where  $S \subset \mathbb{C}$ . When referring to open sets in  $\mathbb{C}$  and continuity of functions  $f$  we will always consider  $\mathbb{C}$  (and its subsets) as a metric space with respect to the metric  $d(z_1, z_2) = |z_1 - z_2|$ , where  $|\cdot|$  denotes the complex modulus, i.e.,  $|z| = \sqrt{x^2 + y^2}$  whenever  $z = x + iy$  with  $x, y \in \mathbb{R}$ . An open ball with respect this metric will be also referred to as an open disc and denoted by

$$D(a; r) = \{z \in \mathbb{C} : |z - a| < r\},$$

where  $a$  is the center and  $r > 0$  is the radius of the open ball. The closed disc with center  $a$  and radius  $r$  is denoted by  $\overline{D(a; r)}$ , so

$$\overline{D(a; r)} = \{z \in \mathbb{C} : |z - a| \leq r\}.$$

Recall that  $G \subset \mathbb{C}$  is called open if for all  $a \in G$  there exists  $r > 0$  such that  $D(a; r) \subset G$ .

If  $z = x + iy$ , then the conjugate  $\bar{z}$  of  $z$  is defined by  $\bar{z} = x - iy$ . Now  $z\bar{z} = |z|^2$ , so that  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}$  for  $z \neq 0$ . Elementary properties of complex numbers are given by:

- (1) The real part  $\operatorname{Re} z$  of  $z$  satisfies  $\operatorname{Re} z = \frac{1}{2}(z + \bar{z})$ , while the imaginary part  $\operatorname{Im} z$  of  $z$  is given by  $\operatorname{Im} z = \frac{1}{2i}(z - \bar{z})$ .
- (2) For all  $z_1, z_2 \in \mathbb{C}$  we have  $z_1 + z_2 = \bar{z}_1 + \bar{z}_2$  and  $\overline{z_1 z_2} = \bar{z}_1 \bar{z}_2$ .
- (3) For all  $z_1, z_2 \in \mathbb{C}$  we have  $|z_1 z_2| = |z_1| |z_2|$ .

### 2. Elementary transcendental functions

Recall also that if  $z = x + iy \neq 0$ , then, using polar coordinates, we can write  $z = r \cos \theta + ir \sin \theta$ . In this case we write  $\arg z = \{\theta + 2k\pi : k \in \mathbb{Z}\}$ . By  $\operatorname{Arg} z$  we will denote the principal value of the argument of  $z \neq 0$ , i.e.  $\theta = \operatorname{Arg} z \in \arg z$  if  $-\pi < \theta \leq \pi$ . Note that if  $z_1 = |z_1|(\cos \theta_1 + i \sin \theta_1)$  and  $z_2 = |z_2|(\cos \theta_2 + i \sin \theta_2)$ , then we have  $z_1 z_2 = |z_1| |z_2| (\cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2 + i(\sin \theta_1 \cos \theta_2 + \cos \theta_1 \sin \theta_2)) = |z_1 z_2| (\cos(\theta_1 + \theta_2) + i(\sin(\theta_1 + \theta_2)))$ . Hence we have  $\arg(z_1 z_2) = \arg z_1 + \arg z_2$ . Define now  $e^z = e^x(\cos y + i \sin y)$ . Then  $|e^z| = e^x$  and  $\arg e^z = y + 2k\pi$ . In particular  $e^{2\pi i} = 1$  and the function  $e^z$  is  $2\pi i$ -periodic, i.e.,  $e^{z+2\pi i} = e^z e^{2\pi i} = e^z$  for all  $z \in \mathbb{C}$ . We want now to define  $\log w$  such that  $w = e^z$  where  $z = \log w$ , but we can not define it as just the inverse of  $e^z$  as  $e^z$  is not one-to-one. Consider therefore the equation  $w = e^z$  for a given  $w$ . We must assume that  $w \neq 0$  as  $e^z \neq 0$  (and thus  $\log 0$  is not defined). Then  $|w| = |e^z| = e^x$  and  $y = \operatorname{Arg} w + 2k\pi$  ( $k \in \mathbb{Z}$ ). Hence  $\{\log |w| + i(\operatorname{Arg} w + 2k\pi) : k \in \mathbb{Z}\}$  is the set of all solutions  $z$  of  $w = e^z$ . We write  $\log w$  for any  $w$  in the set  $\{\log |w| + i(\operatorname{Arg} w + 2k\pi) : k \in \mathbb{Z}\}$ .

DEFINITION 2.1. Let  $G \subset \mathbb{C}$  be an open connected set and  $f : G \rightarrow \mathbb{C}$  a continuous function such that  $z = e^{f(z)}$  for all  $z \in G$ . Then  $f$  is called a *branch of the logarithm* on  $G$ .

It is clear that if  $f$  is a branch of the logarithm on  $G$ , then  $0 \notin G$  and  $f(z) = \log |z| + i(\text{Arg } z + 2k\pi)$  for some  $k \in \mathbb{Z}$ , where  $k$  can depend on  $z$ . Also, if  $f$  is a branch of the logarithm on  $G$ , then for fixed  $k$  also  $g(z) = f(z) + 2k\pi i$  is a branch of the logarithm on  $G$ . The converse also holds.

PROPOSITION 2.2. Let  $G \subset \mathbb{C}$  be an open connected set and  $f : G \rightarrow \mathbb{C}$  a branch of the logarithm on  $G$ . Then every other branch of the logarithm on  $G$  is of the form  $f + 2k\pi i$  for some fixed  $k \in \mathbb{Z}$ .

PROOF. Suppose  $g$  is another branch of the logarithm on  $G$ . Then define  $h = \frac{1}{2\pi i}(f - g)$ . Then  $h$  is continuous on  $G$ ,  $h(G) \subset \mathbb{Z}$ , and  $G$  connected implies that  $h(G) = \{k\}$  for some  $k \in \mathbb{Z}$ .  $\square$

To find a branch of  $\log z$  for a given open and connected set  $G$  requires finding (as  $\log |z|$  is continuous on  $\mathbb{C} \setminus \{0\}$ ) a continuous selection of  $\arg z$  in  $\{\text{Arg } z + 2k\pi\}$ . As  $G$  is connected, the range of this continuous selection has to be an interval of length at most  $2\pi$ , but such a selection does not always exist! This happens e.g. in case  $G = \mathbb{C} \setminus \{0\}$ , then  $G$  is open and connected, but there does not exist a branch of  $\log z$  on  $G$ , i.e.,  $\text{Arg } z$  is discontinuous on the negative  $x$ -axis. In the next examples we construct some branches of  $\log z$ .

- EXAMPLE 2.3. (i) Let  $G = \mathbb{C} \setminus \{z \in \mathbb{R} : z \leq 0\}$ . Then  $\text{Arg } z$  is continuous on  $G$ , so  $f(z) = \log |z| + i\text{Arg } z$  is a branch of  $\log z$  on  $G$ . This branch is called the *principal branch* of  $\log z$  and denoted by  $\text{Log } z$ .
- (ii) Let  $G = \mathbb{C} \setminus \{z \in \mathbb{R} : z \geq 0\}$ . Let  $\theta(z)$  denote the unique value of  $\arg z$  such that  $0 < \theta(z) < 2\pi$ . Then  $f(z) = \log |z| + i\theta(z)$  is a branch of  $\log z$  on  $G$ .

### 3. Differentiable functions

DEFINITION 3.1. Let  $G \subset \mathbb{C}$  be an open set and  $f : G \rightarrow \mathbb{C}$ . Then  $f$  is *differentiable* at  $z \in G$  if

$$\lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h}$$

exists. When this limit exists we denote it by  $f'(z)$  and call it the (complex) derivative of  $f$  at  $z$ . If  $f'(z)$  exists at every point of  $G$ , then we call  $f$  *analytic* or *holomorphic* on  $G$ .

NOTATION.  $H(G) = \{f : g \rightarrow \mathbb{C}; f \text{ holomorphic in } G\}$ .

If  $S \subset \mathbb{C}$  is any set, then we say that  $f$  is holomorphic in  $S$  if  $f \in H(G)$  for some open set  $G \supset S$ .

REMARKS 3.2.

1. The function  $f$  is differentiable at  $z \in G$ , if for  $|h|$  small enough we can write  $f(z+h) = f(z) + f'(z)h + \epsilon(h)h$ , where  $\epsilon(h) \rightarrow 0$  as  $h \rightarrow 0$ . From this it follows directly that if  $f$  is differentiable at  $z$ , then  $f$  is continuous at  $z$ .

2. Note that  $f$  is differentiable at  $z_0 \in G$  with derivative equal to  $f'(z_0)$  is equivalent to saying that for all  $\epsilon > 0$  there exists a  $\delta > 0$  such that

$$\left| \frac{f(z+h) - f(z)}{h} - f'(z_0) \right| < \epsilon$$

for all  $h \in \mathbb{C}$  with  $0 < |h| < \delta$ . In particular we can take  $h = x$  with  $x$  real and  $0 < |x| < \delta$  or  $h = iy$  with  $y$  real and  $0 < |y| < \delta$ . This fact will be exploited in the proof of the next theorem.

**THEOREM 3.3.** (*Cauchy–Riemann equations*) Let  $G \subset \mathbb{C}$  be an open set and  $f : G \rightarrow \mathbb{C}$  be differentiable at  $z = x + iy \in G$ . Let  $f(z) = u(x, y) + iv(x, y)$ , where  $u$  and  $v$  are real valued functions on  $G$ . Then the first order partials  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$  and  $\frac{\partial v}{\partial y}$  exist at  $(x, y)$  and satisfy the Cauchy–Riemann equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

at the point  $(x, y)$ .

**PROOF.** In the definition of the derivative we can restrict ourselves first to real valued  $h \rightarrow 0$ . We get then that

$$f'(z) = \lim_{h \rightarrow 0, h \in \mathbb{R}} \left\{ \frac{u(x+h, y) - u(x, y)}{h} + i \frac{v(x+h, y) - v(x, y)}{h} \right\} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

exists at  $z = x + iy$  and similarly by restricting to  $h = ik$  with  $k$  real valued and  $k \rightarrow 0$ , we get

$$f'(z) = \lim_{k \rightarrow 0, k \in \mathbb{R}} \left\{ \frac{u(x, y+k) - u(x, y)}{ik} + i \frac{v(x, y+k) - v(x, y)}{ik} \right\} = -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y}.$$

Equating the two expressions for  $f'(z)$  we get that

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

at the point  $(x, y)$ . □

**EXAMPLE 3.4.**

- (i) Let  $f(z) = \bar{z}z = x^2 + y^2$ . Then  $\frac{\partial u}{\partial x} = 2x$ ,  $\frac{\partial v}{\partial y} = 0$ ,  $\frac{\partial u}{\partial y} = 2y$  and  $\frac{\partial v}{\partial x} = 0$ . Hence the Cauchy–Riemann equations hold if and only if  $(x, y) = (0, 0)$ . At  $z = 0$  we have

$$\frac{f(0+h) - f(0)}{h} = \bar{h} \rightarrow 0$$

as  $h \rightarrow 0$ . Hence  $f$  is differentiable only at  $z = 0$  and thus nowhere holomorphic as there exists no open set  $G$  containing 0 on which  $f$  is differentiable.

- (ii) Let  $f(z) = c$ , where  $c \in \mathbb{C}$  is a constant. Then  $f'(z) = 0$  for all  $z \in \mathbb{C}$ , so  $f \in H(\mathbb{C})$ . Similarly if  $g(z) = z$ , then  $g'(z) = 1$  for all  $z \in \mathbb{C}$ , so  $g \in H(\mathbb{C})$   
 (iii) Let  $f(z) = 1/z$  on  $\mathbb{C} \setminus \{0\}$ . Then

$$\frac{f(z+h) - f(z)}{h} = \frac{-1}{z(z+h)} \rightarrow \frac{-1}{z^2}$$

for all  $z \neq 0$ , so that  $f$  is holomorphic on  $\mathbb{C} \setminus \{0\}$ .

DEFINITION 3.5. A function  $f : \mathbb{C} \rightarrow \mathbb{C}$  is called *entire* if  $f$  is holomorphic on  $\mathbb{C}$ .

The above example shows that  $f(z) = c$  and  $f(z) = z$  are entire functions. To get additional examples of holomorphic and entire functions we first observe that analogously to the rules of differentiation of real valued functions one can prove the following proposition.

PROPOSITION 3.6. *Let  $G$  be a nonempty open subset of  $\mathbb{C}$ . Then the following holds.*

- (1) *If  $f, g$  holomorphic on  $G$  and  $\lambda \in \mathbb{C}$ , then so are  $f + g$ ,  $\lambda f$ , and  $fg$ .*
- (2) *If  $f(G) \subset G_1$ , where  $G_1$  is open and  $g \in H(G_1)$ , then  $h = g \circ f$  is holomorphic on  $G$  and  $h'(z) = g'(f(z))f'(z)$  for all  $z \in G$ .*

PROOF. We will only prove ???. Let  $z \in G$  and put  $w = f(z)$ . Then  $f$  being holomorphic at  $z$  implies that we can write

$$f(z+h) - f(z) = [f'(z) + \epsilon_1(h)]h,$$

where  $\epsilon_1(h) \rightarrow 0$  as  $h \rightarrow 0$ . Similarly

$$g(w+k) - g(w) = [g'(w) + \epsilon_2(k)]k,$$

where  $\epsilon_2(k) \rightarrow 0$  as  $k \rightarrow 0$ . Putting  $k = f(z+h) - f(z)$  we get

$$\begin{aligned} \frac{g(f(z+h)) - g(f(z))}{h} &= (g'(f(z)) + \epsilon_2(f(z+h) - f(z)))(f'(z) + \epsilon_1(h)) \\ &\rightarrow g'(f(z))f'(z) \end{aligned}$$

as  $h \rightarrow 0$ . □

- COROLLARY 3.7. (1) *Any polynomial  $p(z) = a_0 + a_1z + \dots + a_nz^n$  is entire.*  
 (2) *Any rational function  $f(z) = \frac{p(z)}{q(z)}$ , where  $p$  and  $q$  are polynomials, is holomorphic on  $\mathbb{C} \setminus \{z \in \mathbb{C} : q(z) = 0\}$ .*

We will now compare complex differentiability of  $f = u + iv$  with the real differentiability of the map  $(u, v) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ . Recall first the definition of real differentiability of a vector valued mapping.

DEFINITION 3.8. Let  $G \subset \mathbb{R}^m$  an open set and  $F : G \rightarrow \mathbb{R}^n$ . Then  $F$  is real differentiable at  $c \in G$  if there exist a linear mapping  $DF(c) : \mathbb{R}^m \rightarrow \mathbb{R}^n$  such that

$$\lim_{h \rightarrow 0} \frac{\|F(c+h) - F(c) - DF(c)h\|}{\|h\|} = 0.$$

Writing  $F = (F_1, \dots, F_n)$ , where  $F_i : \mathbb{R}^m \rightarrow \mathbb{R}$ , then real differentiability of  $F$  at  $c \in G$  is equivalent with the real differentiability of each  $F_i$  and  $DF_i(c)h = \nabla F_i(c) \cdot h$ , where  $\nabla F_i$  denotes the gradient of  $F_i$  and thus  $DF(c)$  is the linear map given by the Jacobian matrix of  $F$ . We now take  $m = n = 2$  to compare complex differentiability of  $f = u + iv$  at  $z_0 = x_0 + iy_0$  with real differentiability of  $F = (u, v)$  at  $c = (x_0, y_0)$ . We first deal with the special case of a linear map.

LEMMA 3.9. *Let  $A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a real linear map, given by the matrix  $[a_{i,j}]$ . Then  $A = (u, v)$  where  $f = u + iv$  is a complex linear map from  $\mathbb{C}$  to  $\mathbb{C}$  if and only if  $a_{1,1} = a_{2,2}$  and  $a_{1,2} = -a_{2,1}$ .*

PROOF. Assume first that  $f(z) = Cz$  for some  $C = c_1 + ic_2$ . Then  $u(x, y) = (c_1x - c_2y)$  and  $v(x, y) = (c_2x + c_1y)$ , which implies immediately that  $A = (u, v)$  is a linear map with matrix  $[a_{i,j}]$ , where  $a_{1,1} = a_{2,2} = c_1$  and  $a_{1,2} = -a_{2,1} = -c_2$ . Conversely, if  $a_{1,1} = a_{2,2} = c_1$  and  $a_{1,2} = -a_{2,1} = -c_2$ , then it is straightforward to check that  $f(z) = Cz$  with  $C = c_1 + ic_2$ .  $\square$

REMARK 3.10. Note that the condition on the matrix  $A$  are the ones imposed by the Cauchy-Riemann equations for  $f(z) = Cz = u + iv$ . As the real derivative  $DF(c)$  of a linear map  $F : \mathbb{C} \rightarrow \mathbb{C}$  is  $F(c)$  this says that a linear map from  $\mathbb{R}^2 \rightarrow \mathbb{R}^2$  corresponds to a complex differentiable map from  $\mathbb{C}$  to  $\mathbb{C}$  if and only if it is complex linear.

An immediate consequence of the Lemma is the following theorem.

THEOREM 3.11. *Let  $G \subset \mathbb{C}$  be an open set and  $f : G \rightarrow \mathbb{C}$ , where  $f(z) = u(x, y) + iv(x, y)$ . Let  $z_0 = x_0 + iy_0 \in G$ . then the following are equivalent.*

- (1)  $f$  is complex differentiable at  $z_0$ .
- (2)  $F = (u, v)$  is real differentiable at  $(x_0, y_0)$  and the derivative  $DF(x_0, y_0)$  is complex linear.
- (3)  $F = (u, v)$  is real differentiable at  $(x_0, y_0)$  and the Cauchy-Riemann equations hold at  $(x_0, y_0)$ .

To prove a theorem about complex differentiability when the Cauchy-Riemann equations hold, we need first a result from vector calculus.

LEMMA 3.12. *Let  $G$  be an open subset of  $\mathbb{R}^2$  and  $u : G \rightarrow \mathbb{R}^2$  a function which has partial derivatives on  $G$ , which are continuous at  $(x_0, y_0) \in G$ . Then there exist  $\epsilon_1(h)$ , and  $\epsilon_2(h)$  in a neighborhood of  $(0, 0)$  with  $\epsilon_1(h) \rightarrow 0$  and  $\epsilon_2(h) \rightarrow 0$  as  $h = (h_1, h_2) \rightarrow (0, 0)$  such that*

$$u(x_0 + h_1, y_0 + h_2) = u(x_0, y_0) + \frac{\partial u}{\partial x}(x_0, y_0)h_1 + \frac{\partial u}{\partial y}(x_0, y_0)h_2 + \epsilon_1(h)h_1 + \epsilon_2(h)h_2.$$

PROOF. Let  $r > 0$  such that for  $h = (h_1, h_2)$  with  $\|h\| < r$  we have that  $(x_0 + h_1, y_0 + h_2) \in G$ . Let  $\|h\| < r$ . Then by the Mean Value theorem there exist  $k_1$  between  $x_0$  and  $x_0 + h_1$  and  $k_2$  between  $y_0$  and  $y_0 + h_2$  such that

$$\begin{aligned} u(x_0 + h_1, y_0 + h_2) - u(x_0, y_0) &= u(x_0 + h_1, y_0 + h_2) - u(x_0, y_0 + h_2) \\ &\quad + u(x_0, y_0 + h_2) - u(x_0, y_0) \\ &= \frac{\partial u}{\partial x}(k_1, y_0 + h_2)h_1 + \frac{\partial u}{\partial y}(x_0, k_2)h_2. \end{aligned}$$

The proof now follows if we put  $\epsilon_1(h) = \frac{\partial u}{\partial x}(k_1, y_0 + h_2) - \frac{\partial u}{\partial x}(x_0, y_0)$  and  $\epsilon_2(h) = \frac{\partial u}{\partial y}(x_0, k_2) - \frac{\partial u}{\partial y}(x_0, y_0)$ .  $\square$

THEOREM 3.13. *Let  $G \subset \mathbb{C}$  be an open set and  $f : G \rightarrow \mathbb{C}$ . Let  $f(z) = u(x, y) + iv(x, y)$ , where  $u$  and  $v$  are real valued functions on  $G$ . Assume that the first order partials  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$  and  $\frac{\partial v}{\partial y}$  exist on  $G$ , are continuous at  $(x, y)$  and satisfy the Cauchy-Riemann equations*

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

*at the point  $(x, y)$ . Then  $f$  is complex differentiable at  $z = x + iy$ .*

PROOF. Identifying  $\mathbb{C}$  with  $\mathbb{R}^2$  we can find by the above lemma  $\epsilon_j(h)$  with  $\epsilon_j(h) \rightarrow 0$  as  $h = h_1 + ih_2 \rightarrow 0$  for  $j = 1, \dots, 4$  such that

$$\begin{aligned} \frac{f(z+h) - f(z)}{h} &= \frac{\partial u}{\partial x}(x, y) \frac{h_1}{h} + \frac{\partial u}{\partial y}(x, y) \frac{h_2}{h} + \epsilon_1(h) \frac{h_1}{h} + \epsilon_2(h) \frac{h_2}{h} \\ &\quad + i \left( \frac{\partial v}{\partial x}(x, y) \frac{h_1}{h} + \frac{\partial v}{\partial y}(x, y) \frac{h_2}{h} + \epsilon_3(h) \frac{h_1}{h} + \epsilon_4(h) \frac{h_2}{h} \right) \\ &= \frac{\partial u}{\partial x}(x, y) + i \frac{\partial v}{\partial x}(x, y) + \epsilon_1(h) \frac{h_1}{h} + \epsilon_2(h) \frac{h_2}{h} \\ &\quad + i \epsilon_3(h) \frac{h_1}{h} + i \epsilon_4(h) \frac{h_2}{h} \\ &\rightarrow \frac{\partial u}{\partial x}(x, y) + i \frac{\partial v}{\partial x}(x, y) \end{aligned}$$

as  $h \rightarrow 0$ , since  $|\frac{h_1}{h}| \leq 1$  and  $|\frac{h_2}{h}| \leq 1$ .  $\square$

COROLLARY 3.14. *Let  $f(z) = e^z$ . Then  $f$  is entire and  $f'(z) = e^z$  for all  $z \in \mathbb{C}$ .*

PROOF. If  $f = u + iv$ , then  $u(x, y) = e^x \cos y$  and  $v(x, y) = e^x \sin y$ . Now  $\frac{\partial u}{\partial x}(x, y) = e^x \cos y$ ,  $\frac{\partial v}{\partial x}(x, y) = e^x \sin y$ ,  $\frac{\partial u}{\partial y}(x, y) = -e^x \sin y$ , and  $\frac{\partial v}{\partial y}(x, y) = e^x \cos y$ . Hence the Cauchy-Riemann equations hold for all  $(x, y)$  and, as the partials are continuous, it follows from the above theorem that  $f$  is holomorphic at all  $z \in \mathbb{C}$ . Moreover  $f'(z) = \frac{\partial u}{\partial x}(x, y) + i \frac{\partial v}{\partial x}(x, y) = e^z$ .  $\square$

PROPOSITION 3.15. *Let  $G_1, G_2 \subset \mathbb{C}$  be open sets and let  $f : G_1 \rightarrow G_2$ ,  $g : G_2 \rightarrow G_1$  be continuous mappings such that  $g(f(z)) = z$  for all  $z \in G_1$ . If  $g$  is holomorphic on  $G_2$  and  $g'(z) \neq 0$  for all  $z \in G_2$ , then  $f$  is holomorphic on  $G_1$  and  $f'(z) = \frac{1}{g'(f(z))}$  for all  $z \in G_1$ .*

PROOF. Let  $z \in G_1$ . Then for  $h \neq 0$  but small enough we have  $z+h \in G_1$  and  $f(z+h) \neq f(z)$ , since  $g(f(z)) = z \neq (z+h) = g(f(z+h))$ . Now

$$1 = \frac{g(f(z+h)) - g(f(z))}{f(z+h) - f(z)} \frac{f(z+h) - f(z)}{h}$$

implies that  $f$  is differentiable at  $z$  and  $1 = g'(f(z))f'(z)$ .  $\square$

COROLLARY 3.16. *Let  $G \subset \mathbb{C}$  be an open connected set and  $f : G \rightarrow \mathbb{C}$  a branch of the logarithm on  $G$ . Then  $f$  is holomorphic on  $G$  and  $f'(z) = \frac{1}{z}$  for all  $z \in G$ .*

PROOF. Take  $g(z) = e^z$  in the above proposition.  $\square$

We conclude this section with some remarks about harmonic functions. Recall that if  $G \subset \mathbb{R}^2$  is open and  $u : G \rightarrow \mathbb{R}$  satisfies the Laplace equation  $\Delta u = \frac{\partial^2 u}{\partial x^2}(x, y) + \frac{\partial^2 u}{\partial y^2}(x, y) = 0$  on  $G$ . Let now  $f \in H(G)$ , let  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$ . Assume that  $u$  and  $v$  have continuous second order partials (an assumption which we will show later on to be always true). Then  $\Delta u = \frac{\partial^2 u}{\partial x^2}(x, y) + \frac{\partial^2 u}{\partial y^2}(x, y) = \frac{\partial^2 v}{\partial x \partial y}(x, y) + \frac{\partial^2 v}{\partial y \partial x}(x, y) = 0$ . Hence  $u$  is harmonic on  $G$ . Similarly  $v$  is harmonic on  $G$ . Two harmonic functions  $u$ , and  $v$  are called conjugate harmonic functions, when  $f = u + iv$  is holomorphic on  $G$ . Another consequence of the Cauchy-Riemann equations is that the inner product of the gradients  $\nabla u$  and  $\nabla v$  satisfy  $\nabla u \cdot \nabla v = 0$ , i.e., the level curves  $u(x, y) = c_1$  and  $v(x, y) = c_2$  intersect orthogonally.



#### 4. Power series

In this section we will see how one can use power series to get a large class of examples of holomorphic functions. In fact, in a later chapter we will see that locally every holomorphic function can be so obtained. We start by recalling some basic facts concerning series. Recall that if  $\langle a_n \rangle_{n \geq 0}$  is a sequence of complex numbers, then the series  $\sum_{n=0}^{\infty} a_n$  converges to  $s \in \mathbb{C}$  if  $|s - s_n| \rightarrow 0$  as  $n \rightarrow \infty$ , where  $s_n = a_0 + \dots + a_n$ . The number  $s$  is then called the sum of the series. The series is said to diverge, if it does not converge to any  $s \in \mathbb{C}$ . As in the real variable case we have:

- (1) If  $\sum_{n=0}^{\infty} a_n$  converges, then  $a_n \rightarrow 0$  as  $n \rightarrow \infty$ .
- (2) If  $\sum_{n=0}^{\infty} |a_n|$  converges, then  $\sum_{n=0}^{\infty} a_n$  converges.

A power series is a series of the form  $\sum_{n=0}^{\infty} c_n(z-a)^n$ . Usually we will treat  $z$  as a variable and the  $c_n$ 's and  $a$  as constants in this expression.

EXAMPLE 4.1. Consider the geometric series  $\sum_{n=0}^{\infty} z^n$ . The partial sums  $s_n$  are in this case given by  $s_n = 1 + \dots + z^n = \frac{1-z^{n+1}}{1-z}$  for all  $z \neq 1$ . Hence for  $|z| < 1$  the series  $\sum_{n=0}^{\infty} z^n$  converges and has sum equal to  $\frac{1}{1-z}$ , while if  $|z| \geq 1$  the series diverges, since in that case it is not true that  $z^n \rightarrow 0$  as  $n \rightarrow \infty$ .

The following simple result turns out to be a useful tool in studying the convergence of power series.

THEOREM 4.2. (*Weierstrass M-test*) Let  $G \subset \mathbb{C}$  and  $u_n : G \rightarrow \mathbb{C}$  such that  $|u_n(z)| \leq M_n$  on  $G$ , where  $\sum_0^{\infty} M_n < \infty$ . Then  $\sum_0^{\infty} u_n(z)$  converges uniformly on  $G$ .

PROOF. For fixed  $z \in G$  we have that  $\sum_0^{\infty} |u_n(z)| \leq \sum_0^{\infty} M_n < \infty$ . Hence the series  $\sum_0^{\infty} u_n(z)$  converges for all  $z \in G$ . Let  $f(z) = \sum_0^{\infty} u_n(z)$  for  $z \in G$  denote the sum of the series and let  $\epsilon > 0$ . Then there exists  $N$  such that  $\sum_{k=N+1}^{\infty} M_k < \epsilon$ . Then we have for all  $z \in G$  and all  $n \geq N$  that

$$\left| f(z) - \sum_{k=0}^n u_k(z) \right| = \left| \sum_{k=n+1}^{\infty} u_k(z) \right| \leq \sum_{k=n+1}^{\infty} |u_k(z)| \leq \sum_{k=n+1}^{\infty} M_k < \epsilon$$

for all  $n \geq N$  and all  $z \in G$  and thus the series  $\sum_0^{\infty} u_n(z)$  converges uniformly to  $f(z)$  on  $G$ .  $\square$

For a given power series  $\sum_{n=0}^{\infty} c_n(z-a)^n$  we define the *radius of convergence*  $R$ ,  $0 \leq R \leq \infty$ , by  $\frac{1}{R} = \overline{\lim} \sqrt[n]{|c_n|}$ . The circle  $\{z \in \mathbb{C} : |z-a| = R\}$  is called the *circle of convergence* of the power series.

THEOREM 4.3. (*Cauchy Root test*) Let  $\sum_{n=0}^{\infty} c_n(z-a)^n$  be a power series with radius of convergence  $R$ . Then the following holds.

- (1)  $\sum_{n=0}^{\infty} c_n(z-a)^n$  converges absolutely for  $|z-a| < R$ .
- (2)  $\sum_{n=0}^{\infty} c_n(z-a)^n$  diverges for  $|z-a| > R$ .
- (3) If  $0 < r < R$ , then  $\sum_{n=0}^{\infty} c_n(z-a)^n$  converges uniformly on  $|z-a| \leq r$ .

PROOF. Let  $|z-a| < r < R$ . Then  $\frac{1}{r} > \frac{1}{R}$  implies that there exists  $N$  such that  $|c_n|^{\frac{1}{n}} < \frac{1}{r}$  for all  $n \geq N$ . It follows that  $|c_n(z-a)^n| \leq \left(\frac{|z-a|}{r}\right)^n$  for all  $n \geq N$ .

Since  $\frac{|z-a|}{r} < 1$ , it follows that  $\sum_{n=0}^{\infty} c_n(z-a)^n$  converges absolutely for  $|z-a| < r$  for any  $r < R$  and thus 1. holds. Let now  $|z-a| > r > R$ . Then there exist infinitely many  $n$  such that  $|c_n|^{\frac{1}{n}} > \frac{1}{r}$ . Hence  $|c_n(z-a)^n| \geq \left(\frac{|z-a|}{r}\right)^n > 1$  for infinitely many  $n$ , i.e., the series  $\sum_{n=0}^{\infty} c_n(z-a)^n$  diverges for  $|z-a| > r$  for any  $r > R$  and thus 2. holds. To prove 3. let  $0 < r < s < R$ . Then as above there exists  $N$  such that  $|c_n|^{\frac{1}{n}} < \frac{1}{s}$  for all  $n \geq N$ . It follows that  $|c_n(z-a)^n| \leq \left(\frac{r}{s}\right)^n$  for all  $n \geq N$  and all  $|z-a| < r$ . Since  $\frac{r}{s} < 1$ , it follows that  $\sum_{n=0}^{\infty} c_n(z-a)^n$  converges uniformly on  $|z-a| \leq r$  by the Weierstrass M-test.  $\square$

In dealing with power series with coefficients involving factorials, it is often easier to use the following result.

**THEOREM 4.4. (Ratio test)** *Let  $\sum_{n=0}^{\infty} c_n(z-a)^n$  be a power series with radius of convergence  $R$ . Assume  $c_n \neq 0$  for all  $n$ . Then*

$$\underline{\lim} \left| \frac{c_{n+1}}{c_n} \right| \leq \frac{1}{R} \leq \overline{\lim} \left| \frac{c_{n+1}}{c_n} \right|.$$

*In particular, if  $\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right|$  exists, then  $\frac{1}{R} = \lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right|$ .*

**PROOF.** Exercise  $\square$

A power series can converge or diverge at any point of its circle of convergence as can be seen from the following examples.

**EXAMPLE 4.5.**

- (i) The series  $\sum_{n=0}^{\infty} \frac{(z+1)^n}{2^{n+1}}$  has  $R = 2$ , as  $\overline{\lim} \sqrt[n]{\frac{1}{2^{n+1}}} = \frac{1}{2}$ . Note that the sum of series equals  $\frac{1}{1-z}$  for all  $|z+1| < 2$ , since  $\frac{1}{1-z} = \frac{1}{2-(z+1)} = \frac{1}{2} \frac{1}{1-\frac{z+1}{2}} = \frac{1}{2} \sum_{n=0}^{\infty} \left(\frac{z+1}{2}\right)^n$  for  $\frac{|z+1|}{2} < 1$ .
- (ii) The series  $\sum_{n=1}^{\infty} \frac{z^n}{n^2}$  has  $R = 1$  (e.g. by the Ratio test), and the series converges absolutely for any  $z$  on the circle of convergence as  $\sum_{n=1}^{\infty} \frac{1}{n^2} < \infty$ .
- (iii) The series  $\sum_{n=1}^{\infty} \frac{z^n}{n}$  has  $R = 1$  (e.g. by the Ratio test), but it does not converge absolutely for any  $z$  on the circle of convergence as  $\sum_{n=1}^{\infty} \frac{1}{n} = \infty$ . In particular it diverges for  $z = 1$ . One can show however (but this is not completely trivial) that it converges for any  $z \neq 1$  with  $|z| = 1$  (for  $z = -1$  this follows e.g. from the so-called alternating series test).
- (iv) The series  $\sum_{n=0}^{\infty} \frac{z^n}{n!}$  has  $R = \infty$  (e.g. by the Ratio test). We will see after the next theorem that  $e^z$  equals the sum of this series.
- (v) The series  $\sum_{n=1}^{\infty} n!z^n$  has  $R = 0$  (e.g. by the Ratio test). Hence it converges only for  $z = 0$ .

**PROPOSITION 4.6.** *Let  $R$  be the radius of convergence of  $\sum_{n=0}^{\infty} c_n(z-a)^n$ . Then  $R$  is also the radius of convergence of the power series  $\sum_{n=1}^{\infty} n c_n(z-a)^{n-1} = \sum_{n=0}^{\infty} (n+1)c_{n+1}(z-a)^n$ .*

**PROOF.** From calculus we know that  $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$ . Hence

$$\overline{\lim} \sqrt[n+1]{(n+1)|c_{n+1}|} = \overline{\lim} \left( \sqrt[n+1]{(n+1)|c_{n+1}|} \right)^{\frac{n+1}{n}} = \frac{1}{R}.$$

$\square$

Note, if we apply the above proposition twice, we get that  $\sum_{n=2}^{\infty} n(n-1)z^{n-2}$  converges absolutely for  $|z-a| < R$ .

The following theorem says that inside the circle of convergence the sum of the power series is a holomorphic function.

**THEOREM 4.7.** *Let  $\sum_{n=0}^{\infty} c_n(z-a)^n$  have radius of convergence  $R \neq 0$  and define  $f(z) = \sum_{n=0}^{\infty} c_n(z-a)^n$  for  $|z-a| < R$ . Then  $f \in H(D(a; R))$  and  $f'(z) = \sum_{n=1}^{\infty} n c_n(z-a)^{n-1}$  for  $|z-a| < R$ .*

**PROOF.** It follows from the above corollary that  $g(z) = \sum_{n=1}^{\infty} n c_n(z-a)^{n-1}$  also converges in  $D(a; R)$ . Remains to show that  $f'(z) = g(z)$  on  $|z-a| < R$ . W.l.o.g. we can assume that  $a = 0$ . In the argument below we will use that  $(z+h)^n - z^n = h \sum_{k=1}^n (z+h)^{k-1} z^{n-k}$ . Let  $z, z+h \in D(0; r)$ , where  $0 < r < R$ . Then we have

$$\begin{aligned} \left| \frac{f(z+h) - f(z)}{h} - g(z) \right| &= \left| \sum_{n=1}^{\infty} c_n \left\{ \frac{(z+h)^n - z^n}{h} - n z^{n-1} \right\} \right| \\ &= \left| \sum_{n=2}^{\infty} c_n \sum_{k=1}^n \{ (z+h)^{k-1} z^{n-k} - z^{n-1} \} \right| \\ &\leq \sum_{n=2}^{\infty} |c_n| \sum_{k=2}^n |z^{n-k} ((z+h)^{k-1} - z^{k-1})| \\ &\leq |h| \sum_{n=2}^{\infty} |c_n| \left( \sum_{k=2}^n |z^{n-k}| \left( \sum_{l=1}^{k-1} |z+h|^{l-1} |z|^{k-1-l} \right) \right) \\ &\leq |h| \sum_{n=2}^{\infty} |c_n| \sum_{k=2}^n (k-1) r^{n-k} r^{k-2} \\ &= |h| \sum_{n=2}^{\infty} |c_n| \frac{1}{2} n(n-1) r^{n-2} \rightarrow 0 \end{aligned}$$

as  $h \rightarrow 0$ , since, by the above proposition,  $\sum_{n=2}^{\infty} |c_n| \frac{1}{2} n(n-1) r^{n-2} < \infty$  as  $r < R$ . Hence  $f'(z) = g(z)$  on  $|z| < r$  for any  $r < R$  and the proof is complete.  $\square$

**COROLLARY 4.8.** *Let  $f(z) = \sum_{n=0}^{\infty} c_n(z-a)^n$  have radius of convergence  $R \neq 0$ . Then  $f^{(k)}(z)$  exists on  $D(a; R)$  for all  $k \geq 1$  and thus  $f^{(k)} \in H(D(a; R))$  and*

$$f^{(k)}(z) = \sum_{n=k}^{\infty} c_n n(n-1) \dots (n-k+1) (z-a)^{n-k}$$

for all  $k \geq 1$  and all  $|z-a| < R$ . In particular  $k!c_k = f^{(k)}(a)$  and thus the coefficients  $c_k$  of the power series are unique.

**EXAMPLE 4.9.** Let  $f(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$ . Then by the above theorem

$$f'(z) = \sum_{n=1}^{\infty} \frac{n z^{n-1}}{n!} = \sum_{n=1}^{\infty} \frac{z^{n-1}}{(n-1)!} = f(z)$$

for all  $z \in \mathbb{C}$ . Let  $h(z) = e^{-z} f(z)$ . Then  $h'(z) = -e^{-z} f(z) + e^{-z} f'(z) = 0$  for all  $z \in \mathbb{C}$ . From the next proposition it follows that  $h(z) = h(0) = 1$  for all  $z$ , i.e.,  $f(z) = e^z$  for all  $z$ .

PROPOSITION 4.10. *Let  $G \subset \mathbb{C}$  be an open and connected set. Assume  $f \in H(G)$  such that  $f'(z) = 0$  for all  $z \in G$ . Then  $f$  is constant on  $G$ .*

PROOF. Let  $z_0 \in G$  and put  $A = \{z \in G : f(z) = f(z_0)\}$ . Then the continuity of  $f$  implies that  $A$  is closed. Let now  $a \in A$ . Then there exists  $\epsilon > 0$  such that  $D(a; \epsilon) \subset G$ . Let  $z \in D(a; \epsilon)$  and put  $g(t) = f(tz + (1-t)a)$  for  $0 \leq t \leq 1$ . Then by the chain rule  $g'(t) = f'(tz + (1-t)a)(z - a) = 0$  for  $0 < t < 1$ , so  $g$  is constant on  $0 \leq t \leq 1$ . Hence  $f(z) = g(0) = g(1) = f(z_0)$ , and thus  $D(a; \epsilon) \subset A$ . It follows that  $A$  is nonempty open and closed subset of  $G$ , thus  $A = G$ .  $\square$

## Integration over contours

### 1. Curves and Contours

A *curve* is a continuous map  $\gamma : [a, b] \rightarrow \mathbb{C}$ . We call  $\gamma(a)$  the initial point and  $\gamma(b)$  the end point of the curve  $\gamma$ , and  $[a, b]$  is called the parameter interval of  $\gamma$ . If  $\gamma(a) = \gamma(b)$ , then  $\gamma$  is called a *closed* curve. Denote by  $\gamma^*$  the range of  $\gamma$ . The curve  $\gamma$  induces an orientation of  $\gamma^*$ , namely the direction in which  $\gamma(t)$  traces  $\gamma^*$  as  $t$  increases from  $a$  to  $b$ . Often we will specify a curve by its range together with an orientation indicating how (and possibly how often) the range is traversed. Given a curve  $\gamma$  we can find an oriented curve  $-\gamma$ , with identical range, but with opposite orientation, e.g.,

$$(-\gamma)(t) = \gamma(a + b - t) \text{ where } a \leq t \leq b$$

as a parametrization of the curve  $-\gamma$ . If  $\gamma_1$  and  $\gamma_2$  are two curves with parameter intervals  $[a_1, b_1]$ ,  $[a_2, b_2]$  respectively such that  $\gamma_1(b_1) = \gamma_2(a_2)$ , then we can join the two curves to get the curve  $\gamma = \gamma_1 \cup \gamma_2$  by taking

$$\gamma(t) = \begin{cases} \gamma_1(t) & a_1 \leq t \leq b_1 \\ \gamma_2(t + a_2 - b_1) & b_1 \leq t \leq b_1 + b_2 - a_2. \end{cases}$$

A curve  $\gamma$  is called *smooth*, if  $\gamma'(t)$  exists and is continuous for all  $a \leq t \leq b$  (with one-sided derivatives at  $a$  and  $b$ ). Note if we write  $\gamma(t) = x(t) + iy(t)$ , then  $\gamma'(t)$  exists if and only if  $x'(t)$  and  $y'(t)$  exist. From multi-variable calculus we know that  $\gamma'(t)$  represents a tangent vector to the curve  $\gamma$ .

A *path or contour*  $\gamma$  is a piecewise smooth curve, i.e.,  $\gamma : [a, b] \rightarrow \mathbb{C}$  such that there exist  $a = t_0 < t_1 < \dots < t_n = b$  where  $\gamma$  restricted to  $[t_{i-1}, t_i]$  is smooth for  $i = 1, \dots, n$ . Note that  $\gamma$  can have corners at the points  $\gamma(t_i)$ , i.e., the right and left hand derivatives of  $\gamma(t)$  at  $t_i$  can differ.

A path  $\gamma$  is called *simple* if  $\gamma : [a, b] \rightarrow \mathbb{C}$  is such that  $\gamma(s) \neq \gamma(t)$  for all  $a \leq s < t \leq b$ , except possibly for  $s = a$  and  $t = b$ . The path  $\gamma$  is *closed* if  $\gamma(a) = \gamma(b)$ .

EXAMPLE 1.1.

- (i) The directed line segment  $C$  from  $z_1$  to  $z_2$  is the range of a smooth curve. A parametrization of  $C$  is given  $\gamma : [0, 1] \rightarrow \mathbb{C}$  defined by  $\gamma(t) = (1-t)z_1 + tz_2$ . We will denote this curve by  $[z_1, z_2]$ .
- (ii) A circular arc oriented counterclockwise is the range of an curve. Suppose the arc is part of the circle with center  $z_0$  and radius  $r$ , then  $\gamma(t) = z_0 + re^{it}$  with  $\theta_1 \leq t \leq \theta_2$  will trace a circular arc counterclockwise. If  $\theta_2 - \theta_1 = 2\pi$  the curve will be the complete circle. Note the curve is simple if and only if  $\theta_2 - \theta_1 \leq 2\pi$ .

**1.1. Conformal mappings.** Let  $f$  be a holomorphic function on an open set  $G \subset \mathbb{C}$ . Let  $z_0 \in G$  be a fixed point and let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a smooth curve in  $G$  passing through  $z_0$  with non-zero tangent, i.e.,  $\gamma(t_0) = z_0$  for some  $t_0 \in (a, b)$  and  $\gamma'(t_0) \neq 0$ . Then  $\gamma_1 = f \circ \gamma$  is a curve passing through  $f(z_0)$  and  $\gamma_1'(t_0) = f'(z_0)\gamma'(t_0)$ . If now  $f'(z_0) \neq 0$ , we see that  $\arg \gamma_1'(t_0) = \arg f'(z_0) + \arg \gamma'(t_0)$  and  $|\gamma_1'(t_0)| = |f'(z_0)||\gamma'(t_0)|$ . Thus the tangent vector  $\gamma'(t_0)$  to the curve  $\gamma$  at  $z_0$  is under the mapping  $f$  rotated over an angle  $\theta \in \arg f'(z_0)$  and stretched by a factor  $|f'(z_0)|$ . Applying this to two curves passing through  $z_0$  we see that under the mapping  $f$  the angle between the two curves is preserved (including the direction they are measured), while their tangent vectors are stretched by the same amount. Mappings which preserve angles (including the direction they are measured) between smooth curves are called *conformal*. Thus we have proved:

**THEOREM 1.2.** *Let  $f$  be a holomorphic function on an open set  $G \subset \mathbb{C}$ . Assume  $f'(z) \neq 0$  for all  $z \in G$ . Then  $f$  is conformal on  $G$ .*

We will now see that in fact the converse is true too. To do so we will introduce some additional notation. Let  $f = u + iv$  as usual. Then we define  $\frac{\partial f}{\partial z} = \frac{\partial f}{\partial x} - i\frac{\partial f}{\partial y}$  and  $\frac{\partial f}{\partial \bar{z}} = \frac{\partial f}{\partial x} + i\frac{\partial f}{\partial y}$ . It is now a routine calculation to show that  $u$  and  $v$  satisfy the Cauchy-Riemann equations if and only if  $\frac{\partial f}{\partial \bar{z}} = 0$ .

**THEOREM 1.3.** *Let  $f = u + iv$  be a function on an open set  $G \subset \mathbb{C}$  with continuous partials. Assume  $f$  is conformal on  $G$ . Then  $f$  is holomorphic on  $G$  and  $f'(z) \neq 0$  for all  $z \in G$ .*

**PROOF.** Let  $\gamma$  be a smooth curve with non-zero tangent passing through  $z_0 \in G$ . Let  $\gamma_1(t) = f(\gamma(t))$ . Write  $\gamma(t) = x(t) + iy(t)$ . Then  $\gamma_1' = \frac{\partial u}{\partial x}x' + \frac{\partial u}{\partial y}y' + i\frac{\partial v}{\partial x}x' + i\frac{\partial v}{\partial y}y' = \frac{\partial f}{\partial z}x' + \frac{\partial f}{\partial \bar{z}}y' = \frac{\partial f}{\partial z}\gamma' + \frac{\partial f}{\partial \bar{z}}\overline{\gamma'}$ . Let  $\gamma(t_0) = z_0$ . Then

$$\frac{\gamma_1'(t_0)}{\gamma'(t_0)} = \frac{\partial f}{\partial z} + \frac{\partial f}{\partial \bar{z}} \frac{\overline{\gamma'(t_0)}}{\gamma'(t_0)}.$$

Now  $f$  conformal implies that the argument of the left hand side of this equation is constant modulo  $2\pi$ . This implies that  $\frac{\partial f}{\partial \bar{z}}(z_0) = 0$ , since the argument of  $\frac{\overline{\gamma'(t_0)}}{\gamma'(t_0)}$  is not constant modulo  $2\pi$ , when we take e.g.  $\gamma(t) = z_0 + te^{i\theta}$ . Hence  $u$  and  $v$  satisfy the Cauchy-Riemann equations at  $z_0$  and thus  $f'(z_0)$  exists and  $f'(z_0) = \frac{\partial f}{\partial z}(z_0) = \frac{\gamma_1'(t_0)}{\gamma'(t_0)} \neq 0$ . □

## 2. Contour integrals

**DEFINITION 2.1.** A curve  $\gamma : [a, b] \rightarrow \mathbb{C}$  is called *rectifiable* if  $\gamma$  is of bounded variation, i.e., if

$$\ell(\gamma) = \sup \left\{ \sum_{i=1}^n |\gamma(t_i) - \gamma(t_{i-1})| : a = t_0 < \dots < t_n = b \right\} < \infty.$$

In this case the *length* of  $\gamma$  is defined to be  $\ell(\gamma)$ . Given a continuous  $\gamma : [a, b] \rightarrow \mathbb{C}$  we define  $\int_a^b \gamma(t) dt = \int_a^b \operatorname{Re} \gamma(t) dt + i \int_a^b \operatorname{Im} \gamma(t) dt$ . In case  $\gamma$  is (piecewise)

smooth we have by the Fundamental Theorem of Calculus for real integrals that  $\int_a^b \gamma'(t) dt = \gamma(b) - \gamma(a)$ .

LEMMA 2.2. *Let  $f : [a, b] \rightarrow \mathbb{C}$  be a continuous function. Then*

$$\left| \int_a^b f(t) dt \right| \leq \int_a^b |f(t)| dt.$$

PROOF. Let  $\alpha = \int_a^b f(t) dt$ . If  $\alpha = 0$ , then the inequality is trivial. Assume  $\alpha \neq 0$ . Then we can write  $\alpha = r e^{i\theta}$ , where  $r = |\int_a^b f(t) dt|$ . Now we have

$$r = \operatorname{Re}(e^{-i\theta} \alpha) = \int_a^b \operatorname{Re}(e^{-i\theta} f(t)) dt \leq \int_a^b |f(t)| dt.$$

□

THEOREM 2.3. *Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve. Then  $\gamma$  is rectifiable and*

$$\ell(\gamma) = \int_a^b |\gamma'(t)| dt.$$

PROOF. Without loss of generality we can assume that  $\gamma$  is smooth. Let  $a = t_0 < \dots < t_n = b$  be a partition of  $[a, b]$ . Then by the Fundamental Theorem of Calculus and the above lemma we have

$$|\gamma(t_i) - \gamma(t_{i-1})| = \left| \int_{t_{i-1}}^{t_i} \gamma'(t) dt \right| \leq \int_{t_{i-1}}^{t_i} |\gamma'(t)| dt.$$

This implies that  $\gamma$  is rectifiable and  $\ell(\gamma) \leq \int_a^b |\gamma'(t)| dt$ . For the reverse inequality, let  $\epsilon > 0$ . Then  $\gamma'$  is uniformly continuous on  $[a, b]$ , so there exists  $\delta > 0$  such that  $|\gamma'(t) - \gamma'(s)| < \epsilon$  whenever  $|t - s| < \delta$ . Now there exists a partition  $a = t_0 < \dots < t_n = b$  with  $\Delta t_i = t_i - t_{i-1} < \delta$  such that

$$\left| \int_a^b |\gamma'(t)| dt - \sum_{i=1}^n |\gamma'(t_i)| \Delta t_i \right| < \epsilon.$$

For  $1 \leq i \leq n$  we have now that

$$\begin{aligned} \left| |\gamma(t_i) - \gamma(t_{i-1})| - |\gamma'(t_i)| \Delta t_i \right| &\leq |\gamma(t_i) - \gamma(t_{i-1}) - \gamma'(t_i) \Delta t_i| \\ &= \left| \int_{t_{i-1}}^{t_i} \gamma'(t) - \gamma'(t_i) dt \right| \\ &\leq \int_{t_{i-1}}^{t_i} |\gamma'(t) - \gamma'(t_i)| dt < \epsilon \Delta t_i. \end{aligned}$$

Combining the last two estimates we get

$$\begin{aligned} \int_a^b |\gamma'(t)| dt &\leq \sum_{i=1}^n |\gamma'(t_i)| \Delta t_i + \epsilon \\ &\leq \sum_{i=1}^n (|\gamma(t_i) - \gamma(t_{i-1})| + \epsilon \Delta t_i) + \epsilon \\ &\leq \ell(\gamma) + \epsilon(b-a) + \epsilon \end{aligned}$$

for all  $\epsilon > 0$ . Hence  $\int_a^b |\gamma'(t)| dt \leq \ell(\gamma)$ .  $\square$

Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve and let  $f : \gamma^* \rightarrow \mathbb{C}$  be continuous. Then we define  $\int_\gamma f(z) dz = \int_a^b f(\gamma(t)) \gamma'(t) dt$ .

EXAMPLE 2.4. Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve. Then  $\int_\gamma 1 dz = \gamma(b) - \gamma(a)$ . This is immediate from the definition and the Fundamental Theorem of Calculus.

PROPOSITION 2.5. Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve and let  $f : \gamma^* \rightarrow \mathbb{C}$  be a continuous function. Then the following hold.

- (i)  $\int_{-\gamma} f(z) dz = -\int_\gamma f(z) dz$ , where  $-\gamma(t) = \gamma(a+b-t)$ .
- (ii) If  $\gamma = \gamma_1 \cup \gamma_2$ , then

$$\int_\gamma f(z) dz = \int_{\gamma_1} f(z) dz + \int_{\gamma_2} f(z) dz.$$

- (iii) If  $|f(z)| \leq M$  on  $\gamma^*$ , then  $|\int_\gamma f(z) dz| \leq M\ell(\gamma)$ .
- (iv) (“Independence of parametrization”) Let  $\tau : [a_1, b_1] \rightarrow [a, b]$  be a smooth onto function with  $\tau' > 0$ . Then for  $\gamma_1 = \gamma \circ \tau$  we have

$$\int_{\gamma_1} f(z) dz = \int_\gamma f(z) dz.$$

- (v) If also  $g : \gamma^* \rightarrow \mathbb{C}$  continuous and  $\alpha, \beta \in \mathbb{C}$ , then  $\int_\gamma \alpha f(z) + \beta g(z) dz = \alpha \int_\gamma f(z) dz + \beta \int_\gamma g(z) dz$ .

PROOF. Let  $-\gamma(t) = \gamma(a+b-t)$ . Then  $-\gamma : [a, b] \rightarrow \mathbb{C}$  is piecewise smooth and  $(-\gamma)'(t) = -\gamma'(a+b-t)$  except possibly finitely many points, from which (i) follows directly. Part (ii) is an immediate consequence of the definition. Part (iii) follows from

$$\left| \int_\gamma f(z) dz \right| \leq \int_a^b |f(\gamma(t)) \gamma'(t)| dt \leq M \int_a^b |\gamma'(t)| dt = M\ell(\gamma).$$

Part (iv) follows from the chain rule  $\gamma_1'(t) = \gamma'(\tau(t))\tau'(t)$  and the change of variable rules for real integrals

$$\begin{aligned} \int_{\gamma_1} f(z) dz &= \int_{a_1}^{b_1} f(\gamma_1(t)) \gamma_1'(t) dt = \int_{a_1}^{b_1} f(\gamma(\tau(t))) \gamma'(\tau(t)) \tau'(t) dt \\ &= \int_{\tau(a_1)}^{\tau(b_1)} f(\gamma(s)) \gamma'(s) ds = \int_\gamma f(z) dz. \end{aligned}$$



Part (v) is immediate from the definition and the corresponding property of real integrals.  $\square$

**COROLLARY 2.6.** *Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve and let  $f_n : \gamma^* \rightarrow \mathbb{C}$  be continuous functions which converge uniformly to  $f$  on  $\gamma^*$ . Then*

$$\int_{\gamma} f_n(z) dz \rightarrow \int_{\gamma} f(z) dz$$

as  $n \rightarrow \infty$ .

**PROOF.** Note first that  $f$  is also continuous on  $\gamma^*$  as it is the uniform limit of a sequence of continuous functions. Let  $M_n = \sup_{z \in \gamma^*} |f_n(z) - f(z)|$ . Then by assumption  $M_n \rightarrow 0$  as  $n \rightarrow \infty$ . From (iii) and (v) above we have now

$$\left| \int_{\gamma} f_n(z) dz - \int_{\gamma} f(z) dz \right| = \left| \int_{\gamma} f_n(z) - f(z) dz \right| \leq M_n \ell(\gamma) \rightarrow 0$$

as  $n \rightarrow \infty$ .  $\square$

The following example is important for the development of the theory.

**EXAMPLE 2.7.** Let  $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$  be given by  $\gamma(t) = a + re^{it}$ , i.e.,  $\gamma$  is the circle with center  $a$  and radius  $r$  traversed counterclockwise. We will show that

$$(2.1) \quad \int_{\gamma} (z - a)^n dz = \begin{cases} 0 & \text{if } n \in \mathbb{Z} \setminus \{-1\} \\ 2\pi i & \text{if } n = -1 \end{cases}$$

Since  $\gamma$  is smooth we can write

$$\begin{aligned} \int_{\gamma} (z - a)^n dz &= \int_0^{2\pi} (re^{it})^n ire^{it} dt \\ &= ir^{n+1} \int_0^{2\pi} e^{i(n+1)t} dt \\ &= \begin{cases} ir^{n+1} \left( \frac{1}{i(n+1)} e^{i(n+1)t} \Big|_0^{2\pi} \right) = 0 & \text{if } n \in \mathbb{Z} \setminus \{-1\} \\ 2\pi i & \text{if } n = -1, \end{cases} \end{aligned}$$

which proves the formula. Note that this integral does not depend on  $r$ .

The following Theorem will allow us to extend this example, in case  $n \neq -1$ , to arbitrary closed contours  $\gamma$  with  $a \notin \gamma^*$ .

**THEOREM 2.8.** *Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve and assume  $F$  is holomorphic on (an open set containing)  $\gamma^*$  with  $F'$  continuous on  $\gamma^*$ . Then*

$$\int_{\gamma} F'(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

In particular, if  $\gamma$  is a closed contour, then  $\int_{\gamma} F'(z) dz = 0$ .

**PROOF.** Assume first that  $\gamma$  is smooth. Then by the chainrule  $(F \circ \gamma)'(t) = F'(\gamma(t))\gamma'(t)$  for all  $a \leq t \leq b$ . Hence  $\int_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t))\gamma'(t) dt = \int_{\gamma} (F \circ \gamma)'(t) dz = F(\gamma(b)) - F(\gamma(a))$ , which proves the theorem for the special case of a smooth curve. In the general case, choose  $a = s_0 < s_1 < \dots < s_n = b$  such that

$\gamma_i = \gamma|_{[s_{i-1}, s_i]}$  is smooth. Then  $\int_{\gamma} F'(z) dz = \sum_{i=1}^n \int_{\gamma_i} F'(z) dz = \sum_{i=1}^n F(\gamma(s_i)) - F(\gamma(s_{i-1})) = F(\gamma(b)) - F(\gamma(a))$ .  $\square$

**COROLLARY 2.9.** *Let  $\gamma$  be any closed contour. Then  $\int_{\gamma} (z - a)^n dz = 0$  for all  $n \geq 0$  and if in addition  $a \notin \gamma^*$ , then also  $\int_{\gamma} (z - a)^n dz = 0$  for all  $n \leq -2$ .*

**PROOF.** Take  $F(z) = \frac{1}{n+1}(z - a)^{n+1}$  in the above theorem.  $\square$

Let now  $\{a, b, c\}$  be an ordered triple of complex numbers. Then  $\Delta = \Delta(a, b, c)$  denotes the triangle with vertices  $a, b$ , and  $c$ . By  $\partial\Delta$  we denote curve obtained by joining the line segments  $[a, b]$ ,  $[b, c]$  and  $[c, a]$ , i.e.,  $\partial\Delta$  denotes the boundary of  $\Delta(a, b, c)$  traversed counterclockwise. Hence

$$\int_{\partial\Delta} f(z) dz = \int_{[a,b]} f(z) dz + \int_{[b,c]} f(z) dz + \int_{[c,a]} f(z) dz$$

for any continuous  $f$  on  $\partial\Delta^*$ .

**THEOREM 2.10.** (*Cauchy's Theorem for a Triangle*) *Let  $G \subset \mathbb{C}$  be an open set and assume  $\Delta = \Delta(a, b, c) \subset G$ . Let  $p \in G$  and  $f : G \rightarrow \mathbb{C}$  such that  $f$  is continuous on  $G$  and holomorphic on  $G \setminus \{p\}$ . Then*

$$\int_{\partial\Delta} f(z) dz = 0.$$

**REMARK.** If  $f$  satisfies the above hypotheses, then we shall see later that  $f$  is actually holomorphic on  $G$ .

**PROOF.** Assume first that  $p \notin \Delta = \Delta(a, b, c)$ . Let  $\{a_1, b_1, c_1\}$  be the midpoints of  $[b, c]$ ,  $[c, a]$ , and  $[a, b]$  respectively. Consider the four triangles  $\Delta_1, \Delta_2, \Delta_3$ , and  $\Delta_4$  formed by the triples  $\{a, c_1, b_1\}$ ,  $\{c_1, b, a_1\}$ ,  $\{a_1, b_1, c_1\}$  and  $\{a_1, c, b_1\}$  (see Figure ??). Put  $I = \int_{\partial\Delta} f(z) dz$ . Then

$$I = \sum_{j=1}^4 \int_{\partial\Delta_j} f(z) dz.$$

Now  $|\int_{\partial\Delta_j} f(z) dz| \geq \frac{|I|}{4}$  for at least one  $j$ . By relabeling we can assume that




FIGURE 1.  $\Delta = \Delta_1 \cup \Delta_2 \cup \Delta_3 \cup \Delta_4$

$$\left| \int_{\partial\Delta_1} f(z) dz \right| \geq \frac{|I|}{4}.$$

Dividing similarly  $\Delta_1$  into four triangles by means of the midpoints of the edges and repeating this process, we get a sequence of triangles  $\Delta \supset \Delta_1 \supset \Delta_2 \supset \dots$  such that  $\ell(\partial\Delta_n) = \frac{1}{2^n}L$ , where  $L = \ell(\partial\Delta)$ , and such that

$$(2.2) \quad \left| \int_{\partial\Delta_n} f(z) dz \right| \geq \frac{|I|}{4^n}.$$

Since  $\Delta$  is compact and  $\{\Delta_n\}$  has the finite intersection property, it follows that there exists  $z_0 \in \bigcap_n \Delta_n$ . As  $p \notin \Delta$ , we have that  $z_0 \neq p$  and thus  $f$  is differentiable at  $z_0$ . Let  $\epsilon > 0$ . Then there exists  $r > 0$  such that

$$|f(z) - f(z_0) - f'(z_0)(z - z_0)| \leq \epsilon|z - z_0|$$

for all  $z$  with  $|z - z_0| < r$ . Now  $\ell(\partial\Delta_n) \rightarrow 0$  implies that there exists  $N$  such that  $\Delta_N \subset D(z_0; r)$ . This implies that  $|z - z_0| < \ell(\partial\Delta_N) = \frac{1}{2^N}L$  for all  $z \in \Delta_N$ . By Corollary ?? we know that

$$\int_{\partial\Delta_N} f(z) dz = \int_{\partial\Delta_N} f(z) - f(z_0) - f'(z_0)(z - z_0) dz.$$

This implies that

$$\left| \int_{\partial\Delta_N} f(z) dz \right| \leq (\epsilon 2^{-N}L) (2^{-N}L) = \epsilon (2^{-N})^2 L^2.$$

From the inequality ?? it follows that  $|I| \leq \epsilon L^2$  for all  $\epsilon > 0$  and thus  $I = 0$ . This completes the proof in case  $p \notin \Delta$ . Assume next that  $p$  is a vertex of the




FIGURE 2. The case  $a = p$

triangle  $\Delta(a, b, c)$ , say  $p = a$ . Then pick  $x \in [a, b]$  and  $y \in [a, c]$ . Then by the above  $\int_{\Delta(x, b, y)} f(z) dz = \int_{\Delta(y, b, c)} f(z) dz = 0$  and thus

$$\int_{\partial\Delta} f(z) dz = \int_{\partial\Delta(a, x, y)} f(z) dz \rightarrow 0$$

as  $x, y \rightarrow a$ , since  $\ell(\partial\Delta(a, x, y)) \rightarrow 0$  and  $f$  is bounded on  $\Delta(a, x, y)$ . Hence  $\int_{\partial\Delta} f(z) dz = 0$  also in the case that  $p$  is a vertex of  $\Delta$ . It remains the case that  $p \in \Delta \setminus \{a, b, c\}$ . In that case apply the above to the triangles  $\Delta(a, b, p)$ ,  $\Delta(b, c, p)$  and  $\Delta(c, a, p)$  to get the desired result.  $\square$

DEFINITION 2.11. A set  $S \subset \mathbb{C}$  is called *starlike* if there exists  $a \in S$  such that the line segment  $[a, z] \subset S$  for all  $z \in S$ . The point  $a$  is called a star center of  $S$  in this case.

Recall that a set  $S \subset \mathbb{C}$  is called *convex* if for  $z_1, z_2 \in S$  we have that  $[z_1, z_2] \subset S$ , i.e., a convex set is a starlike set such that every point of  $S$  is a star center of  $S$ .

THEOREM 2.12. (*Cauchy's Theorem for starlike sets*) Let  $G \subset \mathbb{C}$  be an open starlike set. Let  $p \in G$  and  $f : G \rightarrow \mathbb{C}$  such that  $f$  is continuous on  $G$  and holomorphic on  $G \setminus \{p\}$ . Then  $f = F'$  for some holomorphic  $F$  on  $G$ . In particular

$$(2.3) \quad \int_{\gamma} f(z) dz = 0$$

for every closed contour  $\gamma$  in  $G$ .

PROOF. Let  $a \in G$  be a star center of  $G$ . Then the line segment  $[a, z] \subset G$  for all  $z \in G$ . Now define

$$F(z) = \int_{[a, z]} f(w) dw.$$

Let  $z_0 \in G$ . Then there exists  $r > 0$  such that  $D(z_0, r) \subset G$ . Now for any  $z \in D(z_0, r)$  the triangle  $\Delta(a, z_0, z) \subset G$ , so by Theorem ?? we have

$$\int_{\partial\Delta(a, z_0, z)} f(w) dw = 0,$$

and thus

$$F(z) - F(z_0) = \int_{[a, z]} f(w) dw - \int_{[a, z_0]} f(w) dw = \int_{[z_0, z]} f(w) dw.$$

Fixing  $z_0$  we get for all  $z \neq z_0$  in  $G$ , since  $\int_{[z_0, z]} 1 dw = z - z_0$ , that

$$\begin{aligned} \left| \frac{F(z) - F(z_0)}{z - z_0} - f(z_0) \right| &= \left| \frac{1}{z - z_0} \int_{[z_0, z]} f(w) - f(z_0) dw \right| \\ &\leq \frac{1}{|z - z_0|} \left( \sup_{w \in [z_0, z]} |f(w) - f(z_0)| \right) |z - z_0| \\ &= \sup_{w \in [z_0, z]} |f(w) - f(z_0)| \rightarrow 0 \end{aligned}$$

as  $z \rightarrow z_0$ , by the continuity of  $f$  at  $z_0$ . This proves that  $f(z_0) = F'(z_0)$  for all  $z_0 \in G$  and thus  $F$  is holomorphic on  $G$ . Now equation ?? follows from Theorem ??  $\square$

DEFINITION 2.13. Let  $\gamma$  be a closed piecewise smooth curve in  $\mathbb{C}$  and let  $a \in G = \mathbb{C} \setminus \gamma^*$ . Then

$$\text{Ind}_{\gamma}(a) = \frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - a}$$

is called the *index* of  $\gamma$  with respect to  $a$  or *winding number* of  $\gamma$  around  $a$ .

**THEOREM 2.14.** (*Cauchy's Integral Formula for starlike sets*) Let  $G \subset \mathbb{C}$  be an open starlike set and let  $\gamma$  be a closed contour in  $G$ . Let  $f$  be holomorphic on  $G$  and  $z_0 \in G \setminus \gamma^*$ . Then

$$f(z_0) \cdot \text{Ind}_\gamma(z_0) = \frac{1}{2\pi i} \int_\gamma \frac{f(z)}{z - z_0} dz.$$

**PROOF.** Let  $z \in G \setminus \gamma^*$  and define

$$g(z) = \begin{cases} \frac{f(z) - f(z_0)}{z - z_0} & \text{if } z \in G \setminus \{z_0\} \\ f'(z_0) & \text{if } z = z_0 \end{cases}$$

Then  $g$  satisfies the hypotheses of Theorem ??, so

$$\frac{1}{2\pi i} \int_\gamma g(z) dz = 0.$$

Hence

$$\begin{aligned} \frac{1}{2\pi i} \int_\gamma \frac{f(z)}{z - z_0} dz &= \frac{1}{2\pi i} \int_\gamma \frac{f(z_0)}{z - z_0} dz \\ &= f(z_0) \frac{1}{2\pi i} \int_\gamma \frac{1}{z - z_0} dz \\ &= f(z_0) \cdot \text{Ind}_\gamma(z_0), \end{aligned}$$

and thus the proof of the theorem is complete.  $\square$

**REMARK 2.15.** The above theorem is used most often for the case that  $\text{Ind}_\gamma(a) = 1$ . We will see e.g. that  $\text{Ind}_\gamma(z_0) = 1$ , when  $\gamma$  is a circle containing  $z_0$ , traversed counter clockwise once.

**THEOREM 2.16.** (*Fundamental Theorem of Algebra*) Let  $p(z)$  be a polynomial of degree  $m \geq 1$ . Then  $p$  has exactly  $m$  zeros in  $\mathbb{C}$ , counting each zero according to its multiplicity.

**PROOF.** Assume  $p(z) \neq 0$  for all  $z \in \mathbb{C}$ . Then  $f(z) = \frac{1}{p(z)}$  is an entire function. We can assume that  $p(z) = z^m + \dots + a_1 z + a_0$ . Now

$$\begin{aligned} |p(z)| &= |z|^m \left| 1 + \dots + \frac{a_1}{z^{m-1}} + \frac{a_0}{z^m} \right| \\ &\geq |z|^m \left| 1 - \dots - \frac{|a_1|}{|z|^{m-1}} - \frac{|a_0|}{|z|^m} \right| \geq \frac{1}{2} |z|^m \geq \frac{1}{2} R^m \end{aligned}$$

for  $|z| \geq R$  for  $R$  large enough. Now applying Cauchy's Integral formula to  $f(z)$  and  $\gamma_R = Re^{it}$  with  $0 \leq t \leq 2\pi$ , we get

$$\int_{\gamma_R} \frac{f(z)}{z} dz = 2\pi i f(0) = \frac{2\pi i}{p(0)} \neq 0,$$

while

$$\left| \int_{\gamma_R} \frac{f(z)}{z} dz \right| \leq 2\pi \max_{|z|=R} \left| \frac{1}{p(z)} \right| \leq 2\pi \frac{2}{R^m} \rightarrow 0,$$

as  $R \rightarrow \infty$ , which is a contradiction. Hence there exists  $z_1 \in \mathbb{C}$  such that  $p(z_1) = 0$ . Now factor  $p(z) = (z - z_1)p_1(z)$  and repeat the above argument.  $\square$

To apply the Cauchy's Integral formula, we need to be able to compute the index of a curve. We will derive a number of properties of the index, which will facilitate this.

**PROPOSITION 2.17.** *Let  $\gamma$  be a closed contour and let  $G = \mathbb{C} \setminus \gamma^*$ . Then  $\text{Ind}_\gamma(a)$  is an integer for all  $a \in G$ .*

**PROOF.** Let  $\gamma : [b, c] \rightarrow \mathbb{C}$  be piecewise smooth such that  $\gamma(b) = \gamma(c)$ . Then

$$\frac{1}{2\pi i} \int_\gamma \frac{dz}{z-a} = \frac{1}{2\pi i} \int_b^c \frac{\gamma'(s)}{\gamma(s)-a} ds.$$

Let

$$g(t) = \int_b^t \frac{\gamma'(s)}{\gamma(s)-a} ds.$$

Then  $g(b) = 0$  and  $g'(t) = \frac{\gamma'(t)}{\gamma(t)-a}$ , except possibly on the finite set  $S$  where  $\gamma$  is not differentiable. Now  $e^{-g(t)}(\gamma(t)-a)$  is a continuous function such that

$$\begin{aligned} \frac{d}{dt} e^{-g(t)}(\gamma(t)-a) &= e^{-g(t)}\gamma'(t) - g'(t)e^{-g(t)}(\gamma(t)-a) \\ &= e^{-g(t)}\{\gamma'(t) - g'(t)(\gamma(t)-a)\} = 0, \end{aligned}$$

except on the finite set  $S$ . This implies that  $e^{-g(t)}(\gamma(t)-a)$  is constant on  $[b, c]$ . Evaluating this function at  $t = b$  and  $t = c$  gives then

$$e^{-g(b)}(\gamma(b)-a) = \gamma(b)-a = e^{-g(c)}(\gamma(c)-a),$$

which implies  $e^{-g(c)} = 1$ , since  $\gamma(b) = \gamma(c)$ . Hence  $g(c) = 2\pi im$  for some integer  $m$ , and thus  $\frac{1}{2\pi i} \int_\gamma \frac{dz}{z-a} = m$  which completes the proof of the theorem.  $\square$

By the above proposition the index of a closed contour is an integer  $m$ . Intuitively this integer measures how many times the contour  $\gamma$  winds around the point  $a$  and in what direction. From the properties of contour integrals we have immediately that the following properties hold.

- (1)  $\text{Ind}_{-\gamma}(a) = -\text{Ind}_\gamma(a)$ .
- (2) If  $\gamma$  is obtained by joining the closed contours  $\gamma_1$  and  $\gamma_2$ , then

$$\text{Ind}_\gamma(a) = \text{Ind}_{\gamma_1}(a) + \text{Ind}_{\gamma_2}(a).$$

We shall prove that the index of a closed contour depends continuously on the point  $a$  and that therefore the index is constant on each connected component of  $\mathbb{C} \setminus \gamma^*$ . We recall first the relevant definitions. Let  $S \subset \mathbb{C}$ . Then  $S_1$  is called a (connected) *component* of  $S$ , if  $S_1$  is a maximal connected subset of  $S$ . One can show that if  $S_1$  is a connected subset of  $S$ , then so is the relative closure of  $S_1$ . Hence connected components of a set  $S$  are always relatively closed.

**PROPOSITION 2.18.** *Let  $G$  be an open set in  $\mathbb{C}$ . Then every connected component of  $G$  is also open and thus  $G$  is a countable disjoint union of open and relatively closed components.*

**PROOF.** Let  $C$  denote a component of  $G$  and let  $z_0 \in C$ . Let  $\epsilon > 0$  such that  $D(z_0; \epsilon) \subset G$ . Then  $C \cup D(z_0; \epsilon)$  is a connected subset of  $G$  and thus  $C = C \cup D(z_0; \epsilon)$ , i.e.,  $D(z_0; \epsilon) \subset C$ . Hence  $C$  is open. In each component we can pick a different  $a + bi$  with  $a, b \in \mathbb{Q}$ , so there are countably many components.  $\square$

REMARK 2.19. If  $G = \mathbb{C} \setminus K$ , where  $K$  is a compact set, then  $G$  has exactly one unbounded component. In particular, when  $G = \mathbb{C} \setminus \gamma^*$  for a closed contour  $\gamma$ , then  $G$  has one unbounded component.

THEOREM 2.20. *Let  $\gamma$  be a closed contour and let  $G = \mathbb{C} \setminus \gamma^*$ . Then  $\text{Ind}_\gamma$  is constant on each component of  $G$  and  $\text{Ind}_\gamma(a) = 0$  for all  $a$  in the unbounded component of  $G$ .*

PROOF. Define  $f(w) = \text{Ind}_\gamma(w)$  for  $w \in G$ . We first show that  $f : G \rightarrow \mathbb{C}$  is continuous. Let  $w \in G$ . Then  $r = \text{dist}(w, \gamma^*) > 0$ , since  $\gamma^*$  is compact. Let  $\epsilon > 0$  and then take  $0 < \delta < \min\{\frac{r}{2}, \frac{\epsilon\pi r^2}{L}\}$ , where  $L = \ell(\gamma)$ . Then for  $|w_1 - w| < \delta$  we have

$$|f(w) - f(w_1)| = \frac{1}{2\pi} \left| \int_\gamma \frac{(w - w_1)}{(z - w)(z - w_1)} dz \right|.$$

For  $z \in \gamma^*$  we have  $|z - w| \geq r$  and  $|z - w_1| \geq |z - w| - |w - w_1| > \frac{r}{2}$ . Hence

$$|f(w) - f(w_1)| < \frac{\delta}{\pi r^2} L < \epsilon.$$

It follows that  $f$  is continuous. If now  $C \subset G$  is a component, then  $f(C)$  is a connected subset of  $\mathbb{C}$ . On the other hand  $f(C) \subset \mathbb{Z}$  and thus  $f(C)$  consists of a single point. To see that  $\text{Ind}_\gamma(a) = 0$  for all  $a$  in the unbounded component of  $G$ , let  $R > 0$  such that  $\{z : |z| > R\}$  is contained in the unbounded component of  $G$ . Then find  $a \in \mathbb{C}$  with  $|a| > R$  such that  $|z - a| > \frac{L}{\pi}$  for all  $z \in \gamma^*$ . Then

$$|\text{Ind}_\gamma(a)| \leq \frac{1}{2\pi} \frac{\pi}{L} L = \frac{1}{2},$$

and thus  $\text{Ind}_\gamma(a) = 0$ . As  $\text{Ind}_\gamma(a)$  is constant on the unbounded component it follows that this holds for all  $a$  in the unbounded component of  $G$ .  $\square$

EXAMPLE 2.21.

- (i) Let  $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$  be defined by  $\gamma(t) = z_0 + Re^{it}$ . Then  $\gamma$  traces the circle  $|z - z_0| = R$  once counterclockwise. In this case  $\text{Ind}_\gamma(a) = 1$  for  $|a - z_0| < R$  and  $\text{Ind}_\gamma(a) = 0$  for  $|a - z_0| > R$ , since  $\text{Ind}_\gamma(z_0) = 0$  and the component of  $G \setminus \gamma$  containing  $z_0$  equals  $|z - z_0| < R$ .
- (ii) Let  $\gamma : [0, 4\pi] \rightarrow \mathbb{C}$  be defined by  $\gamma(t) = z_0 + Re^{-it}$ . Then  $\gamma$  traces the circle  $|z - z_0| = R$  twice clockwise. In this case  $\text{Ind}_\gamma(a) = -2$  for  $|a - z_0| < R$  and  $\text{Ind}_\gamma(a) = 0$  for  $|a - z_0| > R$ .

The following proposition provides the index for practically every curve encountered in applications.

PROPOSITION 2.22. *Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  a closed curve. Assume there exists  $z_0 \in \mathbb{C} \setminus \gamma^*$ ,  $t_0 \in (a, b)$  and  $\epsilon > 0$  so that the rays  $R_t = \{z_0 + s(\gamma(t) - z_0) : s \geq 0\}$  have the following properties.*

- (1)  $R_t \cap \gamma^* = \{\gamma(t)\}$  for all  $t \in (t_0 - \epsilon, t_0 + \epsilon)$
- (2) The part of  $R_t$  with  $s > 1$  lies in the unbounded component of  $\mathbb{C} \setminus \gamma^*$  and the part with  $0 < s < 1$  lies in a bounded component of  $\mathbb{C} \setminus \gamma^*$ .
- (3)  $\gamma$  traces  $\gamma^* \cap \{\gamma(t) : t \in (t_0 - \epsilon, t_0 + \epsilon)\}$  once counter clockwise.

Then  $\text{Ind}_\gamma(z_0) = 1$ .

PROOF. Let  $f(z) = \log|z - z_0| + i \arg^*(z - z_0)$  be a branch of  $\log(z - z_0)$  with domain  $\mathbb{C} \setminus R_{t_0}$ . Denote by  $\gamma_\epsilon$  the part of the curve  $\gamma$  in  $\mathbb{C} \setminus R_{t_0}$  with initial point  $\gamma(t_0 + \epsilon)$  and terminal point  $\gamma(t_0 - \epsilon)$ . Then

$$\int_{\gamma_\epsilon} \frac{1}{z - z_0} dz = f(\gamma(t_0 - \epsilon)) - f(\gamma(t_0 + \epsilon)) \rightarrow 2\pi i$$

as  $\epsilon \rightarrow 0$ . On the other hand

$$\int_{\gamma_\epsilon} \frac{1}{z - z_0} dz \rightarrow \int_\gamma \frac{1}{z - z_0} dz$$

as  $\epsilon \rightarrow 0$  and thus  $\text{Ind}_\gamma(z_0) = 1$ . □

**THEOREM 2.23.** (*Power series expansion of holomorphic functions*) Let  $G \subset \mathbb{C}$  and let  $f$  be holomorphic on  $G$ . Then for all  $a \in G$  and all  $R > 0$  such that  $D(a; R) \subset G$  there exists (unique)  $c_n$  such that

$$f(z) = \sum_{n=0}^{\infty} c_n (z - a)^n$$

for all  $z \in D(a; R)$ .

PROOF. Let  $0 < r < R$  and define  $\gamma : [0, 2\pi] \rightarrow D(a; R)$  by  $\gamma(t) = a + re^{it}$ . Then  $\text{Ind}_\gamma(z) = 1$  for all  $z \in D(a; r)$ . Hence by the Cauchy's Integral formula (applied to the open set  $D(a; R)$ ) we have

$$f(z) = \frac{1}{2\pi i} \int_\gamma \frac{f(\zeta)}{\zeta - z} d\zeta$$

Now  $\left| \frac{z-a}{\zeta-a} \right| = \frac{|z-a|}{r} < 1$  for all  $z \in D(a; r)$  and all  $\zeta \in \gamma^*$ . Hence the geometric series

$$\sum_{n=0}^{\infty} \frac{(z-a)^n}{(\zeta-a)^{n+1}} = \frac{1}{\zeta-a} \left\{ \frac{1}{1 - \frac{z-a}{\zeta-a}} \right\} = \frac{1}{\zeta-z}$$

converges uniformly in  $\zeta$  on  $\gamma^*$  for each  $z \in D(a; r)$ . Hence

$$\begin{aligned} f(z) &= \frac{1}{2\pi i} \int_\gamma f(\zeta) \sum_{n=0}^{\infty} \frac{(z-a)^n}{(\zeta-a)^{n+1}} d\zeta \\ &= \sum_{n=0}^{\infty} \left( \frac{1}{2\pi i} \int_\gamma \frac{f(\zeta)}{(\zeta-a)^{n+1}} d\zeta \right) (z-a)^n \\ &= \sum_{n=0}^{\infty} c_n (z-a)^n \end{aligned}$$

where

$$c_n = \frac{1}{2\pi i} \int_\gamma \frac{f(\zeta)}{(\zeta-a)^{n+1}} d\zeta.$$

The uniqueness follows from Corollary ?? in Chapter??, where it was shown that  $c_n = \frac{f^{(n)}(a)}{n!}$ . □



COROLLARY 2.24. Let  $G \subset \mathbb{C}$  be an open set and assume  $f : G \rightarrow \mathbb{C}$  is holomorphic. Then  $f'$  is holomorphic on  $G$  and thus  $f^{(n)}$  exists for all  $n \geq 1$  on  $G$ . Moreover, if  $D(a; R) \subset G$  and  $|f(z)| \leq M$  on  $D(a; R)$ , then

$$|f^{(n)}(a)| \leq \frac{n!M}{R^n} \text{ (Cauchy Estimates).}$$

PROOF. The fact that  $f'$  is holomorphic on  $G$  follows immediately from the above theorem and Theorem ???. From Corollary ??? we get

$$f^{(n)}(a) = \frac{n!}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{(\zeta - a)^{n+1}} d\zeta,$$

where  $\gamma(t) = a + re^{it}$ ,  $0 \leq t \leq 2\pi$ ,  $0 < r < R$  and thus

$$|f^{(n)}(a)| \leq \frac{n!}{2\pi} 2\pi r \frac{M}{r^{n+1}} = \frac{n!M}{r^n}.$$

As this holds for all  $0 < r < R$  the proof is complete.  $\square$

THEOREM 2.25. (Morera's Theorem) Let  $G \subset \mathbb{C}$  be an open set and  $f : G \rightarrow \mathbb{C}$  a continuous function such that

$$\int_{\partial\Delta} f(z) dz = 0$$

for all triangles  $\Delta \subset G$ . Then  $f$  is holomorphic on  $G$ .

PROOF. Let  $D(a; R) \subset G$  for  $a \in G$ . Then as in the proof of Theorem ??? we can find  $F$  holomorphic on  $D(a; R)$  such that  $F' = f$  on  $D(a; R)$ . From the above corollary we now conclude that  $f$  is holomorphic on  $D(a; R)$ . As this holds for all  $D(a; R) \subset G$  we conclude that  $f$  is holomorphic on  $G$ .  $\square$

THEOREM 2.26. (Liouville's Theorem) Let  $f$  be an entire function. Assume that  $f$  is bounded on  $\mathbb{C}$ . Then  $f$  is constant.

PROOF. Let  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  be the power series expansion around  $z = 0$ . Since  $f$  is entire, this series has radius of convergence equal to  $\infty$ . Let  $M$  be such that  $|f(z)| \leq M$  for all  $z \in \mathbb{C}$ . Then for all  $R > 0$  we have for  $n \geq 1$  that  $|f^{(n)}(0)| \leq \frac{n!M}{R^n} \rightarrow 0$  as  $R \rightarrow \infty$ . Hence  $f^{(n)}(0) = 0$  for all  $n \geq 1$ , and thus also  $a_n = 0$  for all  $n \geq 1$ . Therefore  $f(z) = a_0$  for all  $z \in \mathbb{C}$ .  $\square$