

An Overview of Number Field Counting

Frank Thorne

University of South Carolina

Québec-Vermont Number Theory Seminar

Definition

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

Definition

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

and for each transitive subgroup $G \subseteq S_d$,

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X, \text{Gal}(K^c/\mathbb{Q}) = G\},$$

Definition

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

and for each transitive subgroup $G \subseteq S_d$,

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X, \text{Gal}(K^c/\mathbb{Q}) = G\},$$

so that

$$N_d(X) = \sum_{\substack{G \subseteq S_d \\ \text{transitive}}} N_d(X, G).$$

Definition

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

and for each transitive subgroup $G \subseteq S_d$,

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X, \text{Gal}(K^c/\mathbb{Q}) = G\},$$

so that

$$N_d(X) = \sum_{\substack{G \subseteq S_d \\ \text{transitive}}} N_d(X, G).$$

Theorem (Finiteness – Hermite)

For each d and X , $N_d(X)$ is finite.

Theorem (Finiteness – Hermite)

For each d and X , $N_d(X)$ is finite.

Theorem (Minkowski's Lower Bound)

If $[K : \mathbb{Q}] = d$, then

$$|\text{Disc}(K)| \geq \left(\frac{d^d}{d!}\right)^2 \left(\frac{\pi}{4}\right)^d.$$

Theorem (Finiteness – Hermite)

For each d and X , $N_d(X)$ is finite.

Theorem (Minkowski's Lower Bound)

If $[K : \mathbb{Q}] = d$, then

$$|\text{Disc}(K)| \geq \left(\frac{d^d}{d!}\right)^2 \left(\frac{\pi}{4}\right)^d.$$

In other words,

$$N_d(X) = 0 \text{ for } X < (5.803 \cdots + o(1))^d.$$

The Inverse Galois Problem

Conjecture

For every d and transitive subgroup $G \subseteq S_d$,

$$X \text{ big enough} \implies N_d(X, G) \neq 0.$$

The Inverse Galois Problem

Conjecture

For every d and transitive subgroup $G \subseteq S_d$,

$$X \text{ big enough} \implies N_d(X, G) \neq 0.$$

Proof.

(Your Name Here)



Malle's Conjecture

Conjecture

In fact we have

$$N_d(X, G) \sim c(G)X^{1/a(G)}(\log X)^{b(G)},$$

where $a(G) \geq 1$ and $b(G) \geq 0$ are **explicitly described** integers.

Malle's Conjecture

Conjecture

In fact we have

$$N_d(X, G) \sim c(G)X^{1/a(G)}(\log X)^{b(G)},$$

where $a(G) \geq 1$ and $b(G) \geq 0$ are **explicitly described** integers.

Four Methods to Count Number Fields

- ▶ Generator methods (Schmidt, Ellenberg-Venkatesh, ...)

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)
Use **class field theory** and/or **Kummer theory**.

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)
Use **class field theory** and/or **Kummer theory**.
Get good results when G is abelian.

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)
Use **class field theory** and/or **Kummer theory**.
Get good results when G is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)
Use **class field theory** and/or **Kummer theory**.
Get good results when G is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)
Count **lattice points** in **prehomogeneous representations**.

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)
Use **class field theory** and/or **Kummer theory**.
Get good results when G is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)
Count **lattice points** in **prehomogeneous representations**.
Brilliant when it works.

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)
Use **class field theory** and/or **Kummer theory**.
Get good results when G is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)
Count **lattice points** in **prehomogeneous representations**.
Brilliant when it works.
- ▶ **Inductive methods** (Klüners, Cohen-Diaz-Olivier, ...)

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)
Use **class field theory** and/or **Kummer theory**.
Get good results when G is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)
Count **lattice points** in **prehomogeneous representations**.
Brilliant when it works.
- ▶ **Inductive methods** (Klüners, Cohen-Diaz-Olivier, ...)
Obtain **old results** from **new**.

Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)
Count in terms of appropriate **algebraic integers**.
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)
Use **class field theory** and/or **Kummer theory**.
Get good results when G is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)
Count **lattice points** in **prehomogeneous representations**.
Brilliant when it works.
- ▶ **Inductive methods** (Klüners, Cohen-Diaz-Olivier, ...)
Obtain **old results** from **new**.
Expand the scope of existing methods.

If $\alpha \in \mathcal{O}_K$ is a generator of K/\mathbb{Q} , then $\mathbb{Z}[\alpha] \subseteq \mathcal{O}_K$ and

$$\begin{aligned} |\text{Disc}(\mathcal{O}_K)| &= \text{Disc}(\mathbb{Z}[\alpha]) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2} \\ &= \text{Disc}(\text{minpoly}_\alpha) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2}. \end{aligned}$$

If $\alpha \in \mathcal{O}_K$ is a generator of K/\mathbb{Q} , then $\mathbb{Z}[\alpha] \subseteq \mathcal{O}_K$ and

$$\begin{aligned} |\text{Disc}(\mathcal{O}_K)| &= \text{Disc}(\mathbb{Z}[\alpha]) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2} \\ &= \text{Disc}(\text{minpoly}_\alpha) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2}. \end{aligned}$$

Theorem (Schmidt)

For each d we have

$$N_d(X) \ll X^{\frac{d+2}{4}}.$$

Schmidt's proof

- ▶ By **Minkowski's theory**, there exists $\alpha \in \mathcal{O}_K$ with **trace 0** and $\|\alpha\|_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$ for all embeddings $\sigma : K \mapsto \mathbb{C}$.

Schmidt's proof

- ▶ By Minkowski's theory, there exists $\alpha \in \mathcal{O}_K$ with trace 0 and $\|\alpha\|_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$ for all embeddings $\sigma : K \mapsto \mathbb{C}$.
- ▶ Assume that $\mathbb{Q}(\alpha) = K$. (If not, induct.)

Schmidt's proof

- ▶ By Minkowski's theory, there exists $\alpha \in \mathcal{O}_K$ with trace 0 and $\|\alpha\|_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$ for all embeddings $\sigma : K \mapsto \mathbb{C}$.
- ▶ Assume that $\mathbb{Q}(\alpha) = K$. (If not, induct.)
- ▶ The minimal polynomial of α is

$$\text{minpoly}_\alpha(x) = \prod_{\sigma} (x - \sigma(\alpha)) = x^n + a_2(\alpha)x^{n-2} + \cdots + a_n(\alpha),$$

with $a_i(\alpha) \in \mathbb{Z}$, $|a_i(\alpha)| \ll |\text{Disc}(K)|^{\frac{i}{2n-2}}$.

Schmidt's proof

- ▶ By Minkowski's theory, there exists $\alpha \in \mathcal{O}_K$ with trace 0 and $\|\alpha\|_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$ for all embeddings $\sigma : K \mapsto \mathbb{C}$.
- ▶ Assume that $\mathbb{Q}(\alpha) = K$. (If not, induct.)
- ▶ The minimal polynomial of α is

$$\text{minpoly}_\alpha(x) = \prod_{\sigma} (x - \sigma(\alpha)) = x^n + a_2(\alpha)x^{n-2} + \cdots + a_n(\alpha),$$

with $a_i(\alpha) \in \mathbb{Z}$, $|a_i(\alpha)| \ll |\text{Disc}(K)|^{\frac{i}{2n-2}}$.

- ▶ There are at most $O(X^{\frac{d+2}{4}})$ possibilities.

Schmidt's proof

- ▶ By Minkowski's theory, there exists $\alpha \in \mathcal{O}_K$ with trace 0 and $\|\alpha\|_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$ for all embeddings $\sigma : K \mapsto \mathbb{C}$.
- ▶ Assume that $\mathbb{Q}(\alpha) = K$. (If not, induct.)
- ▶ The minimal polynomial of α is

$$\text{minpoly}_\alpha(x) = \prod_{\sigma} (x - \sigma(\alpha)) = x^n + a_2(\alpha)x^{n-2} + \cdots + a_n(\alpha),$$

with $a_i(\alpha) \in \mathbb{Z}$, $|a_i(\alpha)| \ll |\text{Disc}(K)|^{\frac{i}{2n-2}}$.

- ▶ There are at most $O(X^{\frac{d+2}{4}})$ possibilities.

Given a base number field K .

Given a base number field K .

Theorem (Class Field Theory)

Abelian extensions L/K of degree d and conductor dividing \mathfrak{m} are in bijection with index d quotients of $\text{Cl}_{\mathfrak{m}}(L)$.

Given a base number field K .

Theorem (Class Field Theory)

Abelian extensions L/K of degree d and conductor dividing \mathfrak{m} are in bijection with index d quotients of $\text{Cl}_{\mathfrak{m}}(L)$.

Theorem (Kummer Theory)

If in addition $\mu_d \subseteq K$, then abelian extensions L/K of exponent d are in bijection with subgroups of $K^{\times}/(K^{\times})^d$.

Theorem (Cohn, 1954)

We have

$$\sum_{K \text{ cyclic cubic}} \frac{1}{\text{Disc}(K)^s} = -\frac{1}{2} + \frac{1}{2} \left(1 + \frac{1}{3^{4s}}\right) \prod_{p \equiv 1 \pmod{6}} \left(1 + \frac{2}{p^{2s}}\right).$$

Corollary

We have

$$N_3(X, C_3) \sim \frac{11\sqrt{3}}{36\pi} \prod_{p \equiv 1 \pmod{6}} \frac{(p+2)(p-1)}{p(p+1)}.$$

Theorem (Wright, Mäki, **but read Wood's treatment**)

Let G be any abelian group of order n . Then we have

$$\sum_{\text{Gal}(K/\mathbb{Q}) \simeq G} \frac{1}{\text{Disc}(K)^s} = \text{finite sum of Euler products} .$$

Corollary

We have

$$N_{|G|}(X, G) \sim c(G)X^{1/a(G)}(\log X)^{b(G)},$$


where $a(G)$ and $b(G)$ are explicit and $c(G)$ is 'explicit'.

Prime degree (Cohen, Diaz y Diaz, Olivier 2002)

"It is claimed that this constant can be explicitly computed as a finite product of local adelic integrals, but in practice this has not been done, even for the simplest Abelian groups G , except for $G = C_2$..."

Prime degree (Cohen, Diaz y Diaz, Olivier 2002)

"It is claimed that this constant can be explicitly computed as a finite product of local adelic integrals, but in practice this has not been done, even for the simplest Abelian groups G , except for $G = C_2 \dots$ "



Cohen, Diaz y Diaz and Olivier, Cyclic extensions 171

Theorem 1.1. *Let K be a number field of signature (r_1, r_2) . Let \mathcal{R} (resp. \mathcal{S}) be the set of prime ideals of K which are ramified (resp. totally split) in K_ℓ/K . Then*

$$N_{K, \ell}(C_\ell, X^{\ell-1}) \sim c_1 c_2 c_3 c_4 X \log^{\ell-1} X$$

with

$$c_1 = \frac{\left(\prod_{d|d_\ell} \zeta_{K_\ell|d}(d)^{\mu(d)} \right)^{\ell-1}}{d_\ell^{\ell r_2 + r_1} \ell!},$$
$$c_2 = \prod_{\mathfrak{p} \in \mathcal{S}} \left(\left(1 + \frac{\ell-1}{\mathcal{N} \mathfrak{p}} \right) \prod_{d|d_\ell} \left(1 - \frac{1}{\mathcal{N} \mathfrak{p}^d} \right)^{(\ell-1)\mu(d)/d} \right),$$
$$c_3 = \left(\prod_{\mathfrak{p} \in \mathcal{R}} \prod_{d|d_\ell} \left(1 - \frac{1}{\mathcal{N} \mathfrak{p}^{d f(\mathfrak{p}_d/\mathfrak{p})}} \right)^{\theta(\mathfrak{p}_d/\mathfrak{p})\mu(d)} \right)^{\ell-1},$$
$$c_4 = \prod_{\mathfrak{p} | \ell, \mathfrak{p} \notin \mathcal{S}} \left(1 + \frac{\ell-1}{\mathcal{N} \mathfrak{p}} - \frac{\ell-1-r(\mathfrak{e}(\mathfrak{p}))(1-1/\mathcal{N} \mathfrak{p})}{\mathcal{N} \mathfrak{p}^{\lceil \mathfrak{e}(\mathfrak{p})/(\ell-1) \rceil}} \right),$$

where $r_2 = 0$ if $\zeta_\ell \in K$, while $r_2 = r_1 - 1$ otherwise, and by abuse of notation, for any number field L we write $\zeta_L(1)$ for the residue of the Dedekind zeta function $\zeta_L(s)$ at $s = 1$.

The parametrization method

هذه رسالة بسم الله الرحمن الرحيم من محمد بن موسى الخوارزمي المشهور بأبي الفتح الخوارزمي
 شريفاً تقسم بربع أسد وربع أسد وربع أسد وربع أسد وربع أسد وربع أسد وربع أسد وربع أسد
 على نظرية فيكون نسبة أولها إلى ثانياً كنسبة قوس إلى قوس مركز الدائرة واه نصف القطر
 نصف القوس فإنها لا تاندينا حتى نرى في العمل المار معلوم
 تركيب على ذلك الصفة فبعد ما ربع أسد وربع أسد وربع أسد وربع أسد
 ربعاً طهان على ذلك واه فانه يخرج عموداً يكون نسبة
 إليه كنسبة قوس إلى قوس وخرج عموداً فاره طسم وسم على ذلك
 بعد ان جعلنا خطاً سم مثل آه فلان نسبة آه إلى ب كنسبة قوس
 إلى قوس وسم مثل آه يكون نسبة سم إلى ب ضرب سم في قوس مساوياً ضرب ب في قوس
 بنية التدبير في معرفة الاصل ضرب سم في قوس مثل على ذلك ضرب ب في قوس مثل على
 ذلك فبكرة على ذلك مساوياً السطح في ذلك مثل على ذلك
 شراً فيكون على ذلك مساوياً سطح ذلك فان جعلنا قطعاً
 لايقا وخطاه ط ط م وخرج على نقطة كما سبقت الجوريس
 في نظرية القائل الاول من كتاب الجوهرة والشكل
 واه من القائل الثاني من هذا الكتاب اذ هذا العمل
 الاشكال لثلاثة فان ذلك القطع الريد ب على مقصود لا يحتمل كما ينبغي من شكل الاشكال
 القائل الثاني من كتاب الجوهرة ونقطة معلومة الموضع وخطاً معلوم الموضع والقطر
 ان نقطة عندنا تكسب غير معلومة الموضع لانها كانت معلومة الموضع لكانت نقطة معلومة
 الموضع لا خطاً معلومة القطر فيكون خطاً معلوم القطر ولكن ان كان الخط معلوماً وكذا

Intersections of conics

Example. Solve $x^4 - x^3 + 3x^2 - 5x + 1 = 0$.

Intersections of conics


Example. Solve $x^4 - x^3 + 3x^2 - 5x + 1 = 0$.

$$y = x^2, \quad y^2 - yx + 3y - 5x + 1 = 0.$$

Intersections of conics

Example. Solve $x^4 - x^3 + 3x^2 - 5x + 1 = 0$.

$$y = x^2, \quad y^2 - yx + 3y - 5x + 1 = 0.$$



Theorem (Bhargava, *Annals*, 2004)

There exists an explicit, discriminant preserving bijection between the following two sets:

Theorem (Bhargava, *Annals*, 2004)

There exists an explicit, discriminant preserving bijection between the following two sets:

- ▶ $GL_3(\mathbb{Z}) \times GL_2(\mathbb{Z})$ -orbits on the lattice $(\text{Sym}^2 \mathbb{Z}^3 \otimes \mathbb{Z}^2)$ of pairs of integral ternary quartic forms.

Theorem (Bhargava, *Annals*, 2004)

There exists an explicit, discriminant preserving bijection between the following two sets:

- ▶ $GL_3(\mathbb{Z}) \times GL_2(\mathbb{Z})$ -orbits on the lattice $(\text{Sym}^2 \mathbb{Z}^3 \otimes \mathbb{Z}^2)$ of **pairs of integral ternary quartic forms**.
- ▶ **Pairs (Q, R) , where Q is a quartic ring and R is a cubic resolvent of Q .**

A Bhargava-style metatheorem

Theorem

There exists an explicit, discriminant preserving bijection between the following two sets:

Theorem

There exists an explicit, discriminant preserving bijection between the following two sets:

- ▶ $G(\mathbb{Z})$ -orbits on a lattice $V(\mathbb{Z})$; where G is an **algebraic group** acting (often **prehomogeneously**) on a vector space V ;

Theorem

There exists an explicit, discriminant preserving bijection between the following two sets:

- ▶ $G(\mathbb{Z})$ -orbits on a lattice $V(\mathbb{Z})$; where G is an **algebraic group** acting (often **prehomogeneously**) on a vector space V ;
- ▶ Some nice class of arithmetic objects we want to count.

How to prove Bhargava-style theorems

How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration.

How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)

How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)
- ▶ Pick your favorite complex representation (G, V) (which should be defined over \mathbb{Z} , and for which the invariant theory should be nice).

How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)
- ▶ Pick your favorite complex representation (G, V) (which should be defined over \mathbb{Z} , and for which the invariant theory should be nice).
- ▶ Try to prove that the $G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$ parametrize something. Hope to get lucky.

Counting Low Degree Fields

Theorem (Davenport-Heilbronn, Bhargava, et al.)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3}(\log X)^{2.09}),$$

Counting Low Degree Fields

Theorem (Davenport-Heilbronn, Bhargava, et al.)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3}(\log X)^{2.09}),$$

$$N_4(X, S_4) \sim \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})X,$$

Counting Low Degree Fields

Theorem (Davenport-Heilbronn, Bhargava, et al.)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3}(\log X)^{2.09}),$$

$$N_4(X, S_4) \sim \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})X,$$

$$N_5(X) \sim \frac{13}{130} \prod_p (1 + p^{-2} - p^{-4} - p^{-5})X.$$

Counting Low Degree Fields

Theorem (Davenport-Heilbronn, Bhargava, et al.)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3}(\log X)^{2.09}),$$

$$N_4(X, S_4) \sim \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})X,$$

$$N_5(X) \sim \frac{13}{130} \prod_p (1 + p^{-2} - p^{-4} - p^{-5})X.$$

These are now **lattice point** counting problems.

Theorem (Cohen, Diaz y Diaz, Olivier)

We have

$$N_4(X, D_4) \sim X \cdot \frac{3}{\pi^2} \sum_D \frac{2^{-r_2(D)}}{D^2} \frac{L(1, D)}{L(2, D)},$$

where the sum ranges over all fundamental discriminants $\neq 1$.

Theorem (Cohen, Diaz y Diaz, Olivier)

We have

$$N_4(X, D_4) \sim X \cdot \frac{3}{\pi^2} \sum_D \frac{2^{-r_2(D)}}{D^2} \frac{L(1, D)}{L(2, D)},$$

where the sum ranges over all fundamental discriminants $\neq 1$.

Theorem (Belabas-Fouvry, Bhargava-Wood)

We have

$$N_6(X, S_3) \sim \frac{2}{9} \left(\frac{4}{3} + \frac{1}{3^{5/3}} + \frac{2}{3^{7/3}} \right) \prod_{p \neq 3} (1 + p^{-1} + p^{-4/3}) \cdot X^{1/3}.$$

Theorem (Belabas-Fouvry, Bhargava-Wood)

We have

$$N_6(X, S_3) \sim \frac{2}{9} \left(\frac{4}{3} + \frac{1}{3^{5/3}} + \frac{2}{3^{7/3}} \right) \prod_{p \neq 3} (1 + p^{-1} + p^{-4/3}) \cdot X^{1/3}.$$

Idea: If K is an S_3 -cubic with $\text{Disc}(K) = Dn^2$, then $\text{Disc}(\tilde{K}) = D^3 n^4$ apart from the 2- and 3-adic factors.

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

Idea: If K is an S_d -field and L is an A -field, then **usually** K and L are linearly disjoint with

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

Idea: If K is an S_d -field and L is an A -field, then **usually** K and L are linearly disjoint with

$$\text{Disc}(KL) = * \cdot \text{Disc}(K)^{|A|} \text{Disc}(L)^d,$$

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

Idea: If K is an S_d -field and L is an A -field, then **usually** K and L are linearly disjoint with

$$\text{Disc}(KL) = * \cdot \text{Disc}(K)^{|A|} \text{Disc}(L)^d,$$

where $*$ is divisible only primes **ramified in both K and L .**

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

Idea: If K is an S_d -field and L is an A -field, then **usually** K and L are linearly disjoint with

$$\text{Disc}(KL) = * \cdot \text{Disc}(K)^{|A|} \text{Disc}(L)^d,$$

where $*$ is divisible only primes **ramified in both K and L .**

(This doesn't happen too often.)

Theorem (Klüners + ϵ)

Assume a 'weak Malle conjecture' of the form

$$N_d(X, G) \ll X^{3/2}.$$

Theorem (Klüners $+\epsilon$)

Assume a 'weak Malle conjecture' of the form

$$N_d(X, G) \ll X^{3/2}.$$

Then we have

$$N_{2d}(X, C_2 \wr G) \sim c(C_2 \wr G)X.$$

Theorem (Klüners + ϵ)

Assume a 'weak Malle conjecture' of the form

$$N_d(X, G) \ll X^{3/2}.$$

Then we have

$$N_{2d}(X, C_2 \wr G) \sim c(C_2 \wr G)X.$$

Idea. A quadratic extension of a G -extension usually has Galois group $C_2 \wr G$.

Theorem (Klüners + ϵ)

Assume a 'weak Malle conjecture' of the form

$$N_d(X, G) \ll X^{3/2}.$$

Then we have

$$N_{2d}(X, C_2 \wr G) \sim c(C_2 \wr G)X.$$

Idea. A quadratic extension of a G -extension usually has Galois group $C_2 \wr G$.

Note. $D_4 \simeq C_2 \wr C_2$; subsumes Cohen-Diaz-Olivier as a special case.

Theorem (Alberts, 2018)

Assume that *“the m -torsion in class groups is small on average”*.
Then, for every solvable transitive subgroup $G \subseteq S_d$ we have

$$N_d(X, G) \ll X^{1/a(G)+\epsilon}.$$

Theorem (Alberts, 2018)

Assume that *“the m -torsion in class groups is small on average”*.
Then, for every solvable transitive subgroup $G \subseteq S_d$ we have

$$N_d(X, G) \ll X^{1/a(G)+\epsilon}.$$

Further development of the same family of ideas.

Theorem (Alberts, 2018)

Assume that *“the m -torsion in class groups is small on average”*.
Then, for every solvable transitive subgroup $G \subseteq S_d$ we have

$$N_d(X, G) \ll X^{1/a(G)+\epsilon}.$$

Further development of the same family of ideas.

Also: See further related works by [Altuğ](#), [Lemke Oliver](#), [Mehta](#), [Shankar](#), [Taniguchi](#), [Varma](#), [Wilson](#), and previously named authors (in various permutations).

Theorem (Lemke Oliver-T., 2020)

We have

$$N_d(X) \ll \dots$$